

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

Vertox 25 Whole Wheat Bait (purple)

Product type(s)

PT14: Rodenticides

Authorisation number: IE/BPA 70512

R4BP asset number: IE-0016189-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	Vertox 25 Whole Wheat Bait Endorats Original Rat & Mouse Killer Bait Endomice Mouse Killer Bait Rascal 25 Brodifacoum Whole Wheat Bait Endorats Rat Killer Plus The Big Cheese Ultra Power Rat & Mouse Killer
---------------	--

1.2. Authorisation holder

Name and address of the authorisation holder	Name	PelGar International Limited
	Address	18 rue des Remparts d'Ainay 69002 Lyon France
Authorisation number	IE/BPA 70512	
<i>R4BP asset number</i>	IE-0016189-0000	
Date of the authorisation	01/07/2015	
Expiry date of the authorisation	01/07/2026	

1.3. Manufacturer(s) of the product

Name of manufacturer	PelGar International Limited
Address of manufacturer	Unit 13, Newman Lane GU34 2QR Alton, Hants United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Unit 13, Newman Lane GU34 2QR Alton, Hants United Kingdom of Great Britain and Northern Ireland (the) Promedivet SRL 545500 SOVATA , str. Lunga nr. 46/ G jud. Mures, Romania

1.4. Manufacturer(s) of the active substance(s)

Active substance	Brodifacoum
Name of manufacturer	PelGar International Limited
Address of manufacturer	Unit 13, Newman Lane GU34 2QR Alton, Hants United Kingdom of Great Britain and Northern Ireland (the)
Location of manufacturing sites	Prazska 54 280 02 Kolin Czechia

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	active substance	56073-10-0	259-980-5	0,0025

2.2. Type(s) of formulation

RB Bait (ready for use)

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H373: May cause damage to organs blood through prolonged or repeated exposure .. EUH208: Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction. EUH208: Contains 2-octyl-2H-isothiazol-3-one (OIT). May produce an allergic reaction.
Precautionary statements	P260: Do not breathe dust. P314: Get medical advice if you feel unwell. P501: Dispose of contents to in accordance with national regulations.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice – general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: Adults and juveniles
Field(s) of use	indoor use Indoors
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: For mice up to 20g in bait stations. Dilution (%): 0 Number and timing of application: For mice use 5-20g of bait in tamper-resistant bait stations. Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Maximum quantity of bait per pack 50g 15, 20, 25, 30 or 50g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE/PP packs (tubs, pails or pouches) 15g 20g, 25g 30g 40g 50g 15, 20, 25, 30 or 50g paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE lined carton 15g 20g, 25g 30g 40g 50g

	<p>15, 20, 25, 30 or 50g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: fibreboard carton/cardboard 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g Loose bait of up to 20g packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid Outer Packaging: cardboard outer 10g, 15g, 20g (packed in multiples of 1/2/4) Maximum allowed 50g Loose bait of up to 20g packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: HDPE or PP bait station packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper 10g, 15g, 20g (packed in multiples of 1/2/4) Maximum allowed 50g</p>
--	---

4.1.1. Use-specific instructions

For mice use 5-20g of bait in tamper-resistant bait stations.

Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service Do not use this product for permanent or pulse-baiting.

4.1.2. Use-specific risk mitigation measures

4.1.3. Use-specific risk mitigation measures

None.

4.1.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

None

4.1.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.1.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None.

4.2. Use description

Table 2. Rats – general public – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All
Field(s) of use	indoor use Indoors
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: For rats up to 60g in bait stations. Dilution (%): 0 Number and timing of application: For rats use 10-60g of bait in covered tamper-resistant bait stations spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days,

	<p>move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.</p>
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	<p>Maximum quantity of bait per pack 150g 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE/PP packs (tubs, pails or pouches) 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g 15, 20, 25, 30, 50 or 60 paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE lined carton 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: fibreboard carton/cardboard 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g Loose bait of up to 60g packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid Outer Packaging: cardboard outer 10g, 15g, 20g,25g, 50g, 60g (packed in multiples of 2/4/8/12/15) Maximum allowed 150g Loose bait of up to 60g packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid Outer Packaging: HDPE or PP bait station packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper 10g, 15g, 20g, 25g, 50g, 60g (packed in multiples of 1, 2 or 4) Maximum allowed 150g</p>

4.2.1. Use-specific instructions

For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Do not use this product for permanent or pulse-baiting.

4.2.2. Use-specific risk mitigation measures

4.2.3. Use-specific risk mitigation measures

None

4.2.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

None

4.2.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.2.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.3. Use description

Table 3. Rats – general public – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.

Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: Other - all
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart. Dilution (%): 0 Number and timing of application: For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.
Category(ies) of users	general public (non-professional)
Pack sizes and packaging material	Maximum quantity of bait per pack 150g 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE/PP packs (tubs, pails or pouches) 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE lined carton 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: fibreboard carton/cardboard 15g 20g, 25g 30g 40g 50g, 60g, 75g, 100g, 120g 150g

	<p>Loose bait of up to 60g packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: cardboard outer 10g, 15g, 20g,25g, 50g, 60g (packed in multiples of 2/4/8/12/15) Maximum allowed 150g</p> <p>Loose bait of up to 60g packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: HDPE or PP bait station packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper 10g, 15g, 20g, 25g, 50g, 60g (packed in multiples of 1, 2 or 4) Maximum allowed 150g</p>
--	--

4.3.1. Use-specific instructions

For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Do not use this product for permanent or pulse-baiting.

4.3.2. Use-specific risk mitigation measures

4.3.3. Use-specific risk mitigation measures

None

4.3.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

None

4.3.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.3.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.4. Use description

Table 4. House mice – professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: All
Field(s) of use	indoor use
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: For mice use up to 20g in tamper-resistant bait stations. Dilution (%): 0 Number and timing of application: For mice use 5-20g of bait in tamper-resistant bait stations. Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the

	bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.
Category(ies) of users	professional
Pack sizes and packaging material	<p>Minimum pack size of 2.5 kg.</p> <p>Up to 25 kg multi-layer paper with PE moisture barrier or multi-layer paper with separate internal PE sack or woven PP with separate internal PE sack or woven PP sack with no liner. Inner packaging: N /A</p> <p>Outer Packaging: Paper, PP, PE sack, PP sack 2.5kg 3kg, 3.5kg, 4kg 4.5kg 5kg 10kg 20kg</p> <p>15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: fibreboard carton/cardboard 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>Loose bait in PE/PP packs (tubs, pails or pouches) Inner packaging: N /A</p> <p>Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg, 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>Loose bait in PE lined carton Inner packaging: N /A</p> <p>Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: cardboard outer Multiples</p> <p>10g –250</p> <p>15g - 250</p> <p>20g – 125, 144</p> <p>25g – 120, 144</p> <p>30g – 96, 120, 144</p> <p>40g – 72, 96, 120, 144</p> <p>50g – 60, 72, 96, 120, 144</p> <p>60g – 48, 60, 72, 96, 120</p>

Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid

Outer Packaging:

HDPE or PP bait station

packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Multiples

10g –250

15g - 250

20g – 125, 144

25g – 120, 144

30g – 96, 120, 144

40g – 72, 96, 120, 144

50g – 60, 72, 96, 120, 144

60g – 48, 60, 72, 96, 120

80g – 32, 48, 60, 72, 96

90g – 32, 48, 60, 72, 96

100g – 32, 48, 60, 72, 96

120g – 24, 32, 48, 60, 72

200g – 16, 24, 32, 48, 60

240g – 16, 24, 32, 48, 60

4.4.1. Use-specific instructions

For mice use 5-20g of bait in tamper-resistant bait stations.

Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Follow any additional instructions provided by the relevant code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

[If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service. Do not use this product for permanent or pulse-baiting.

4.4.2. Use-specific risk mitigation measures

4.4.3. Use-specific risk mitigation measures

Wear protective chemical resistant gloves during product handling phase (EN374).

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

4.4.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.4.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None.

4.4.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None.

4.5. Use description

Table 5. Rats – professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: All
Field(s) of use	indoor use
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: 10-60g of bait in tamper resistant baiting stations spaced 10m apart (5m apart in areas of high infestation).

	<p>Dilution (%): 0</p> <p>Number and timing of application: For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.</p>
Category(ies) of users	professional
Pack sizes and packaging material	<p>Minimum pack size of 2.5 kg. Up to 25 kg multi-layer paper with PE moisture barrier or multi-layer paper with separate internal PE sack or woven PP with separate internal PE sack or woven PP sack with no liner. Inner packaging: N /A Outer Packaging: Paper, PP, PE sack, PP sack 2.5kg 3kg, 3.5kg, 4kg 4.5kg 5kg 10kg 20kg 15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg 15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg 15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: fibreboard carton/cardboard 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg</p> <p>Loose bait in PE/PP packs (tubs, pails or pouches) Inner packaging: N /A Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg, 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg Loose bait in PE lined carton Inner packaging: N /A</p>

Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
cardboard outer Multiples
10g –250
15g - 250
20g – 125, 144
25g – 120, 144
30g – 96, 120, 144
40g – 72, 96, 120, 144
50g – 60, 72, 96, 120, 144
60g – 48, 60, 72, 96, 120

Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
HDPE or PP bait station
packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Multiples
10g –250
15g - 250
20g – 125, 144
25g – 120, 144
30g – 96, 120, 144
40g – 72, 96, 120, 144
50g – 60, 72, 96, 120, 144
60g – 48, 60, 72, 96, 120
80g – 32, 48, 60, 72, 96
90g – 32, 48, 60, 72, 96
100g – 32, 48, 60, 72, 96
120g – 24, 32, 48, 60, 72
200g – 16, 24, 32, 48, 60
240g – 16, 24, 32, 48, 60

4.5.1. Use-specific instructions

For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Follow any additional instructions provided by the relevant code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

[If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Do not use this product for permanent or pulse-baiting.

4.5.2. Use-specific risk mitigation measures

4.5.3. Use-specific risk mitigation measures

Wear protective chemical resistant gloves during product handling phase (EN374).

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

4.5.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.5.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.5.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.6. Use description

Table 6. House mice and/or rats – professionals – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> and/or <i>Rattus norvegicus</i> Common name: House mouse/and or Brown rat Development stage: All
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Application rate: For mice use up to 20g in tamper-resistant bait stations. For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart. Dilution (%): 0 Number and timing of application: For mice use 5-20g of bait in tamper-resistant bait stations. Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

Category(ies) of users	professional
Pack sizes and packaging material	<p>Minimum pack size of 2.5 kg. Up to 25 kg multi-layer paper with PE moisture barrier or multi-layer paper with separate internal PE sack or woven PP with separate internal PE sack or woven PP sack with no liner. Inner packaging: N /A</p> <p>Outer Packaging: Paper, PP, PE sack, PP sack 2.5kg 3kg, 3.5kg, 4kg 4.5kg 5kg 10kg 20kg 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg 15, 20, 25, 30, 50 or 60g paper/PE or AL/PE or paper/Al/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets</p> <p>Outer Packaging: fibreboard carton/cardboard 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg Loose bait in PE/PP packs (tubs, pails or pouches) Inner packaging: N /A</p> <p>Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg, 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg Loose bait in PE lined carton Inner packaging: N /A</p> <p>Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: cardboard outer Multiples 10g –250 15g - 250 20g – 125, 144 25g – 120, 144 30g – 96, 120, 144 40g – 72, 96, 120, 144 50g – 60, 72, 96, 120, 144 60g – 48, 60, 72, 96, 120</p> <p>Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or</p>

	<p>cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid</p> <p>Outer Packaging: HDPE or PP bait station packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Multiples 10g –250 15g - 250 20g – 125, 144 25g – 120, 144 30g – 96, 120, 144 40g – 72, 96, 120, 144 50g – 60, 72, 96, 120, 144 60g – 48, 60, 72, 96, 120 80g – 32, 48, 60, 72, 96 90g – 32, 48, 60, 72, 96 100g – 32, 48, 60, 72, 96 120g – 24, 32, 48, 60, 72 200g – 16, 24, 32, 48, 60 240g – 16, 24, 32, 48, 60</p>
--	--

4.6.1. Use-specific instructions

For mice use 5-20g of bait in tamper-resistant bait stations.

Secure 5-20g of bait in tamper-resistant bait stations spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

For rats use 10-60g of bait in tamper-resistant bait stations spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.

Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

Follow any additional instructions provided by the relevant code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected,

avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

[If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service. Do not use this product for permanent or pulse-baiting.

4.6.2. Use-specific risk mitigation measures

4.6.3. Use-specific risk mitigation measures

Wear protective chemical resistant gloves during product handling phase (EN374).

Do not apply this product directly in the burrows.

Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

4.6.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.6.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.6.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.7. Use description

Table 7. House mice and/or rats – trained professionals – indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> and/or <i>Rattus norvegicus</i> Common name: House mouse and/or Brown rat

	Development stage: All
Field(s) of use	indoor use Indoors
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in covered bait points or in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: For mice use up to 20g in tamper-resistant bait stations or covered bait points. For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart. Dilution (%): 0 Number and timing of application: For mice use 5-20g of bait in tamper-resistant bait stations or covered bait points. Secure 5-20g of bait in tamper-resistant bait stations or covered bait points spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.
Category(ies) of users	trained professional
Pack sizes and packaging material	Minimum pack size of 2.5 kg. Up to 25 kg multi-layer paper with PE moisture barrier or multi-layer paper with separate internal PE sack or woven PP with separate internal PE sack or woven PP sack with no liner. Inner packaging: N /A

Outer Packaging: Paper, PP, PE sack, PP sack 2.5kg 3kg, 3.5kg, 4kg 4.5kg 5kg 10kg 20kg
15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/AL/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/AL/PE sachets

Outer Packaging: PE/PP packs
(tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/AL/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/AL/PE sachets

Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/AL/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/AL/PE sachets

Outer Packaging: fibreboard carton/cardboard 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
Loose bait in PE/PP packs (tubs, pails or pouches) Inner packaging: N /A

Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg, 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
Loose bait in PE lined carton Inner packaging: N /A

Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg
Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
cardboard outer Multiples
10g –250
15g - 250
20g – 125, 144
25g – 120, 144
30g – 96, 120, 144
40g – 72, 96, 120, 144
50g – 60, 72, 96, 120, 144
60g – 48, 60, 72, 96, 120
Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
HDPE or PP bait station
packed in a cardboard outer or blister pack or cardboard

sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Multiples

10g –250

15g - 250

20g – 125, 144

25g – 120, 144

30g – 96, 120, 144

40g – 72, 96, 120, 144

50g – 60, 72, 96, 120, 144

60g – 48, 60, 72, 96, 120

80g – 32, 48, 60, 72, 96

90g – 32, 48, 60, 72, 96

100g – 32, 48, 60, 72, 96

120g – 24, 32, 48, 60, 72

200g – 16, 24, 32, 48, 60

240g – 16, 24, 32, 48, 60

4.7.1. Use-specific instructions

For mice use 5-20g of bait in tamper-resistant bait stations or covered bait points.

Secure 5-20g of bait in tamper-resistant bait stations or covered bait points spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Make frequent inspections of the bait points during the first 10-14 days.

Follow any additional instructions provided by the relevant code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

[If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Remove the remaining product at the end of treatment period.

Do not use this product for permanent baiting.

If used for pulsed baiting: Replace eaten bait only after 3 days and then at maximum 7 day intervals.

4.7.2. Use-specific risk mitigation measures

4.7.3. Use-specific risk mitigation measures

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign.

To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Wear protective chemical resistant gloves during product handling phase (EN374).

4.7.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.7.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.7.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

4.8. Use description

Table 8. House mice and/or rats – trained professionals – outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides.
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> and/or <i>Rattus norvegicus</i> Common name: House mouse and /or Brown rat Development stage: All
Field(s) of use	outdoor use Outdoor around buildings
Application method(s)	Method: bait application

	Detailed description: Ready-to-use bait to be used in covered bait points or in tamper-resistant bait stations
Application rate(s) and frequency	<p>Application rate: For mice use up to 20g in tamper-resistant bait stations or covered bait points. For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart.</p> <p>Dilution (%): 0</p> <p>Number and timing of application:</p> <p>For mice use 5-20g of bait in tamper-resistant bait stations or covered bait points. Secure 5-20g of bait in tamper-resistant bait stations or covered bait points spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.</p> <p>For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart. Secure 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active. Regularly check bait consumption and replace consumed or spoiled bait. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait. If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.</p>
Category(ies) of users	trained professional
Pack sizes and packaging material	<p>Minimum pack size of 2.5 kg. Up to 25 kg multi-layer paper with PE moisture barrier or multi-layer paper with separate internal PE sack or woven PP with separate internal PE sack or woven PP sack with no liner. Inner packaging: N /A Outer Packaging: Paper, PP, PE sack, PP sack 2.5kg 3kg, 3.5kg, 4kg 4.5kg 5kg 10kg 20kg 15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/Al/PE sachets in PE/PP packs (tubs, pails or pouches) Inner packaging: paper/PE, AL/PE or paper/Al/PE sachets Outer Packaging: PE/PP packs</p>

(tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg

15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/AL/PE sachets both in PE lined carton. Inner packaging: paper/PE, AL/PE or paper/AL/PE sachets

Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg

15, 20, 25, 30, 50, 60, 100g or 200g paper/PE or AL/PE or paper/AL/PE sachets, both in fibreboard carton/cardboard outers Inner packaging: paper/PE, AL/PE or paper/AL/PE sachets

Outer Packaging: fibreboard carton/cardboard 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg

Loose bait in PE/PP packs (tubs, pails or pouches) Inner packaging: N /A

Outer Packaging: PE/PP packs (tubs, pails or pouches) 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg, 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg

Loose bait in PE lined carton Inner packaging: N /A

Outer Packaging: PE lined carton 2.5kg, 3kg, 3.5kg, 4kg, 5kg, 6kg, 7kg 8kg, 9kg 10kg, 11kg, 12kg, 13kg, 14kg,15kg, 16kg, 17kg, 18kg, 19kg 20kg

Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in multiples of packed in cardboard outers Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
cardboard outer Multiples
10g –250
15g - 250
20g – 125, 144
25g – 120, 144
30g – 96, 120, 144
40g – 72, 96, 120, 144
50g – 60, 72, 96, 120, 144
60g – 48, 60, 72, 96, 120

Loose bait of up to 20g (rats and mice) and up to 60g (rats only) packed in bait trays with a heat-sealed lid packed in single or multi-use tamper-proof HDPE or PP bait station, all packed in multiples of 1, 2 or 4 in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Inner packaging: bait trays with a heat sealed lid

Outer Packaging:
HDPE or PP bait station
packed in a cardboard outer or blister pack or cardboard sleeve or heat-sealed bag or poly outer heat-sealed with a cardboard topper Multiples
10g –250
15g - 250
20g – 125, 144
25g – 120, 144
30g – 96, 120, 144

40g – 72, 96, 120, 144 50g – 60, 72, 96, 120, 144 60g – 48, 60, 72, 96, 120 80g – 32, 48, 60, 72, 96 90g – 32, 48, 60, 72, 96 100g – 32, 48, 60, 72, 96 120g – 24, 32, 48, 60, 72 200g – 16, 24, 32, 48, 60 240g – 16, 24, 32, 48, 60

4.8.1. Use-specific instructions

For mice use up to 20g in tamper-resistant bait stations or covered bait points.

For mice use 5-20g of bait in tamper-resistant bait stations or covered bait points.

Secure 5-20g of bait in tamper-resistant bait stations or covered bait points spaced 5m apart (2m apart in areas of high infestation) in areas where mice are active. Mice are very inquisitive and it may help the control program to move baits every 2-3 days at the time when bait points are inspected or topped up. Make frequent inspections of the bait points during the first 10-14 days and replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt. If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

For rats use 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart.

Secure 10-60g of bait in tamper-resistant bait stations or covered bait points spaced 10m apart (5m apart in areas of high infestation) in areas where rats are active.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Do not move or disturb bait points for several days after laying bait.

If no signs of rat activity are seen near the bait after 7-10 days, move the bait to an area of higher rat activity.

If all the bait has been eaten from certain areas, increase the quantity of bait by placing more bait points.

Do not increase the bait point size. Replace any bait eaten by rodents or that has been damaged by water or contaminated by dirt.

Regularly check bait consumption and replace consumed or spoilt bait until consumption has stopped. Repeat treatment in situations where there is evidence of new infestation (e.g. fresh tracks or droppings). Make frequent inspections of the bait points during the first 10-14 days.

The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary

Follow any additional instructions provided by the relevant code of best practice.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

[If national policy or legislation requires it] When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.

Remove the remaining product at the end of treatment period.

Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.

If used for pulsed baiting: Replace eaten bait only after 3 days and then at maximum 7 day intervals.

4.8.2. Use-specific risk mitigation measures

4.8.3. Use-specific risk mitigation measures

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign

To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

When used in burrows: Baits must be placed to minimise the exposure to non-target species and children. Cover or block the entrances of baited burrows to reduce the risks of bait being rejected and spilled.

Wear protective chemical resistant gloves during product handling phase (EN374).

4.8.4. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.8.5. Where specific to the use, the instructions for safe disposal of the product and its packaging

None

4.8.6. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

None

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

Do not open the sachets containing the bait (Trained professionals - For non-emptiable sachets only).

Bait stations should be placed in the immediate vicinity where rodent activity has been observed.

Where possible, bait stations must be fixed to the ground or other structures.

Do not place bait stations near water drainage systems where they can come into contact with water.

Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

Professionals & Trained Professionals: If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodents so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Professionals & trained Professionals: -Loose grains: Place the bait in the bait station by using a dosage devise. Specify the methods to minimise dust (e.g. wet wiping).

Remove the remaining bait or the bait stations at the end of the treatment period.

5.2. Risk mitigation measures

Do not use brodifacoum containing products as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity).

Dispose of dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.

Using this product should eliminate rodents within 35 days. The product information (i.e. label and/or leaflet) shall clearly recommend that in case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

[For products to be authorised for professional users:] The product information (i.e. label and/or leaflet) shall clearly show that the product shall not be supplied to the general public (e.g. "for professionals only").

[For products to be authorised for trained professional users:] The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of: Dermal exposure, wash skin with water and then with water and soap.

Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.

Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Contact a veterinary surgeon in case of ingestion by a pet.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call the National Poisons Information Centre (01) 809 2166".

Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. Use of gloves is recommended.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Shelf-life: 24 months

Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

Keep only in original container.

6. OTHER INFORMATION

Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

This product contains a bittering agent and a dye.