


REGULATION (EU) NO 528/2012 CONCERNING THE MAKING
AVAILABLE ON THE MARKET AND USE OF BIOCIDAL
PRODUCTS

Assessment of ACTIVE SUBSTANCES

ASSESSMENT REPORT



FORMIC ACID

Product type 3
"Veterinary hygiene"

EC Number : 200-579-1

CAS Number : 64-18-6

Applicant : Formic Acid Task Force (BASF SE, Kemira Oyj)

Contact details of evaluating CA: BELGIUM

Date : 15/10/2022

TABLE OF CONTENTS

TABLE OF CONTENTS	2
1 STATEMENT OF SUBJECT MATTER AND PURPOSE	8
2 ASSESSMENT REPORT	10
SUMMARY	10
1 PRESENTATION OF THE ACTIVE SUBSTANCE	10
1.1 IDENTITY OF THE ACTIVE SUBSTANCE	10
1.2 INTENDED USES AND EFFECTIVENESS	10
1.3 CLASSIFICATION AND LABELLING	12
1.3.1 <i>Classification and labelling for the active substance</i>	12
1.3.1.1 Current classification and labelling	13
1.3.1.2 Proposed classification and labelling.....	14
1.3.2 <i>Classification and labelling for the representative product(s)</i>	16
1.3.2.1 Proposed classification and labelling.....	16
1.3.2.2 Packaging of the biocidal product	17
2 SUMMARY OF THE HUMAN HEALTH RISK ASSESSMENT	18
2.1 SUMMARY OF THE ASSESMENT OF EFFECTS ON HUMAN HEALTH	18
2.2 REFERENCE VALUES	20
2.3 RISK CHARACTERISATION	21
3 SUMMARY OF THE ENVIRONMENTAL RISK ASSESSMENT	26
3.1 FATE AND BEHAVIOUR IN THE ENVIRONMENT	26
3.2 EFFECTS ASSESSMENT	27
3.3 EXPOSURE ASSESSMENT	27
3.4 RISK CHARACTERIZATION	27
4 ASSESSMENT OF EXCLUSION, SUBSTITUTION CRITERIA AND POP	29
PART A : ASSESSMENT OF INTRINSIC PROPERTIES AND EFFECTS OF THE ACTIVE SUBSTANCE	30
1 GENERAL SUBSTANCE INFORMATION	30
1.1 IDENTIFICATION OF THE SUBSTANCE	30
1.2 COMPOSITION OF THE SUBSTANCE (REFERENCE SPECIFICATIONS)	31
1.3 PHYSICAL AND CHEMICAL PROPERTIES OF THE ACTIVE SUBSTANCE	32
1.4 PHYSICAL HAZARDS AND RESPECTIVE CHARACTERISTICS	36
1.5 HAZARD IDENTIFICATION FOR PHYSICO-CHEMICAL PROPERTIES	39
1.6 ANALYTICAL METHODS FOR DETECTION AND IDENTIFICATION	40
2 EFFECTS AGAINST TARGET ORGANISMS	47
2.1 FUNCTION AND FIELD OF USE ENVISAGED	47
2.2 INTENDED USES	48
2.3 SUMMARY ON EFFICACY	49
2.3.1 <i>Efficacy</i>	49
2.3.2 <i>Mode of action</i>	49
2.3.3 <i>Resistance</i>	50
2.4 CONCLUSION ON EFFICACY.....	50
3 ASSESSMENT OF EFFECTS ON HUMAN HEALTH	51
3.1 TOXICOKINETICS	51
3.1.1 <i>Short summary of the toxicokinetic information</i>	64
3.1.2 <i>Values and conclusions used for the risk assessment</i>	66

3.2	ACUTE TOXICITY	69
3.2.1	Acute oral toxicity.....	69
3.2.2	Acute dermal toxicity.....	71
3.2.3	Acute inhalation toxicity.....	72
3.2.4	Overall conclusion on acute toxicity	74
3.3	IRRITATION AND CORROSION.....	76
3.3.1	Skin corrosion and irritation	76
3.3.2	Eye irritation	80
3.3.3	Respiratory tract irritation.....	81
3.3.4	Overall conclusion on corrosion and irritation.....	85
3.4	SENSITISATION	87
3.4.1	Skin sensitisation	87
3.4.2	Respiratory sensitisation.....	92
3.4.3	Overall conclusion on sensitisation.....	92
3.5	SHORT TERM REPEATED DOSE TOXICITY	93
3.5.1	Short-term oral toxicity.....	93
3.5.2	Short-term dermal toxicity.....	93
3.5.3	Short-term inhalation toxicity.....	94
3.5.4	Overall conclusion on short-term repeated dose toxicity	95
3.6	SUB-CHRONIC REPEATED DOSE TOXICITY	97
3.6.1	Sub-chronic oral toxicity	97
3.6.2	Sub-chronic dermal toxicity	102
3.6.3	Sub-chronic inhalation toxicity	103
3.6.4	Overall conclusion on sub-chronic repeated dose toxicity.....	109
3.7	LONG-TERM REPEATED DOSE TOXICITY	111
3.7.1	Long-term oral toxicity	111
3.7.2	Long-term dermal toxicity	119
3.7.3	Long-term inhalation toxicity	120
3.7.4	Overall conclusion on long-term repeated dose toxicity	121
3.8	GENOTOXICITY	123
3.8.1	In vitro.....	123
3.8.2	In vivo.....	128
3.8.3	Overall conclusion on genotoxicity	129
3.9	CARCINOGENICITY.....	130
3.10	REPRODUCTIVE TOXICITY	136
3.10.1	Developmental toxicity.....	136
3.10.2	Fertility.....	139
3.10.3	Effects on or via lactation	143
3.10.4	Overall conclusion on reproductive toxicity.....	144
3.11	NEUROTOXICITY	146
3.12	IMMUNOTOXICITY.....	153
3.13	DISRUPTION OF THE ENDOCRINE SYSTEM	154
	STEP 1 - Gathering of all relevant information	154
	STEP 2 - Assemble and assess lines of evidence for endocrine activity and adversity	166
	STEP 3 - Sufficiency of the dataset.....	181
	STEP 4 - Initial analysis of the evidence	182
3.14	FURTHER HUMAN DATA.....	183
3.15	OTHER DATA.....	196
4	ENVIRONMENTAL EFFECT ASSESSMENT.....	201
4.1	FATE AND DISTRIBUTION IN THE ENVIRONMENT	201
4.1.1	Degradation.....	201
4.1.1.1	Abiotic degradation	201
4.1.1.1.1	Hydrolysis	201
4.1.1.1.2	Phototransformation in water.....	202
4.1.1.1.3	Estimated photo-oxidation in air	203
4.1.1.2	Biotic degradation	206

4.1.1.2.1	Biodegradability (ready/inherent)	206
4.1.1.3	Rate and route of degradation including identification of metabolites and degradation products	210
4.1.1.3.1	Biological sewage treatment	210
4.1.1.3.1.1	Aerobic biodegradation	210
4.1.1.3.1.2	Anaerobic biodegradation	210
4.1.1.3.1.3	STP simulation test	212
4.1.1.3.2	Biodegradation in freshwater	212
4.1.1.3.2.1	Aerobic aquatic degradation	212
4.1.1.3.2.2	Water/sediment degradation test	212
4.1.1.3.3	Biodegradation in seawater	213
4.1.1.3.3.1	Seawater degradation study	213
4.1.1.3.4	Higher tier degradation studies in water or sediment	214
4.1.1.3.5	Biodegradation during manure storage	214
4.1.1.3.6	Biotic degradation in soil	217
4.1.2	<i>Distribution</i>	223
4.1.2.1	Adsorption onto/desorption from soils	223
4.1.2.2	Higher tier soil adsorption studies	225
4.1.3	<i>Bioaccumulation</i>	226
4.1.3.1	Measured aquatic bioconcentration	226
4.1.3.2	Estimated aquatic bioconcentration	226
4.1.3.3	Measured terrestrial bioconcentration	227
4.1.3.4	Estimated terrestrial bioconcentration	228
4.1.4	<i>Monitoring data</i>	228
4.2	EFFECTS ON ENVIRONMENTAL ORGANISMS	229
4.2.1	<i>Atmosphere</i>	229
4.2.2	<i>Sewage treatment plant (STP)</i>	229
4.2.3	<i>Aquatic compartment</i>	232
4.2.3.1	Freshwater compartment	232
4.2.3.1.1	Acute toxicity (freshwater)	232
4.2.3.1.1.1	Fish	232
4.2.3.1.1.2	Invertebrates (daphnia magna)	233
4.2.3.1.1.3	Algae	235
4.2.3.1.1.3.1	Green algae	235
4.2.3.1.1.3.2	Cyanobacteria or diatoms	236
4.2.3.1.2	Chronic toxicity (freshwater)	240
4.2.3.2	Sediment compartment	242
4.2.3.2.1	Acute toxicity (freshwater sediment)	242
4.2.3.2.2	Chronic toxicity (freshwater sediment)	242
4.2.3.3	Marine compartment	243
4.2.3.3.1	Acute toxicity (seawater)	243
4.2.3.3.1.1	Fish	243
4.2.3.3.1.2	Invertebrates (other species)	243
4.2.3.3.1.3	Algae (diatoms)	244
4.2.3.3.2	Chronic toxicity (seawater)	246
4.2.3.4	Sea sediment compartment	246
4.2.3.4.1	Acute toxicity (sea sediment)	246
4.2.3.4.2	Chronic toxicity (sea sediment)	247
4.2.3.5	Higher tier studies on aquatic organisms	247
4.2.4	<i>Terrestrial compartment</i>	248
4.2.5	<i>Groundwater</i>	248
4.2.6	<i>Birds and mammals</i>	249
4.2.7	<i>Primary and secondary poisoning</i>	251
4.2.7.1	Primary poisoning	251
4.2.7.2	Secondary poisoning	251
4.3	ENDOCRINE DISRUPTING PROPERTIES	252
4.4	DERIVATION OF PNECS	255
5	ASSESSMENT OF EXCLUSION CRITERIA, SUBSTITUTION CRITERIA AND POP	258
5.1	EXCLUSION CRITERIA	258
5.1.1	<i>Assessment of CMR properties</i>	258
5.1.2	<i>Assessment of endocrine disrupting properties</i>	259

5.1.3	PBT Assessment (following Annex XIII to Regulation (EC) No 1907/2006)	260
5.1.3.1	Assessment of persistence	260
5.1.3.1.1	Screening	260
5.1.3.2	Assessment	260
5.1.3.3	Assessment of bioaccumulation	261
5.1.3.3.1	Screening	261
5.1.3.3.2	Assessment	261
5.1.3.4	Assessment of toxicity	261
5.1.3.4.1	Screening	261
5.1.3.4.2	Assessment	262
5.1.3.5	Summary and overall conclusions on PBT or vPvB properties	262
5.1.3.5.1	Summary	262
5.1.3.5.2	Overall conclusion:	262
5.2	SUBSTITUTION CRITERIA	263
5.3	ASSESSMENT OF LONG-RANGE ENVIRONMENTAL TRANSPORTATION AND IMPACT ON ENVIRONMENTAL COMPARTMENTS	264
PART B : EXPOSURE ASSESSMENT AND EFFECTS OF THE ACTIVE SUBSTANCE IN THE BIOCIDAL PRODUCT(S) 265		
6	GENERAL PRODUCT INFORMATION	265
6.1	IDENTIFICATION OF THE PRODUCT	265
6.2	COMPLETE QUALITATIVE AND QUANTITATIVE COMPOSITION OF THE BIOCIDAL PRODUCT	265
6.3	PHYSICAL, CHEMICAL AND TECHNICAL PROPERTIES	266
6.4	HAZARD IDENTIFICATION FOR PHYSICAL AND CHEMICAL PROPERTIES	274
6.5	ANALYTICAL METHODS FOR DETECTION AND IDENTIFICATION	275
7	EFFICACY	290
7.1	EFFICACY	291
7.2	MODE OF ACTION	293
7.3	RESISTANCE	294
7.4	CONCLUSION ON EFFICACY	294
8	HUMAN EXPOSURE ASSESSMENT	295
8.1	IDENTIFICATION OF MAIN PATHS OF HUMAN EXPOSURE TOWARDS ACTIVE SUBSTANCE FROM ITS USE IN BIOCIDAL PRODUCT	297
8.2	LIST OF SCENARIOS	298
8.3	INDUSTRIAL EXPOSURE	299
8.4	PROFESSIONAL EXPOSURE	299
8.4.1	Scenario 1: animal house disinfection by fogging	300
8.4.2	Scenario 2: disinfection of footwear in dipping troughs	306
8.4.3	Scenario 3: animal feet disinfection and disinfection of transport vehicles	313
8.4.4	Scenario 4: restocking of animal housing after disinfection	319
8.4.5	Summary tables: systemic and local exposure from professional uses	322
8.4.6	Combined scenarios	324
8.5	NON-PROFESSIONAL EXPOSURE	325
8.6	SECONDARY EXPOSURE OF THE GENERAL PUBLIC EXCLUDING DIETARY EXPOSURE	325
8.7	DIETARY EXPOSURE	325
8.7.1	Information of non-biocidal use of the active substance	326
8.7.2	Estimating transfer of biocidal active substances into foods as a result of professional and/or industrial application(s)	326
8.7.3	Estimating transfer of biocidal active substances into foods as a result of non-professional use	326
8.8	EXPOSURE ASSOCIATED WITH PRODUCTION, FORMULATION AND DISPOSAL OF THE BIOCIDAL PRODUCT	327
9	ENVIRONMENTAL EXPOSURE ASSESSMENT	328
9.1	EMISSION ESTIMATION	330
9.1.1	Scenario 1: Disinfection of footwear	330
9.1.2	Scenario 2: Disinfection of animal's feet	333
9.1.3	Scenario 3: Disinfection of animal housing by means of fogging	335
9.2	FATE AND DISTRIBUTION IN EXPOSED ENVIRONMENTAL COMPARTMENTS	338
9.3	CALCULATED PEC VALUES	341

9.4	PRIMARY AND SECONDARY POISONING	342
9.4.1	<i>Primary poisoning</i>	342
9.4.2	<i>Secondary poisoning</i>	342
10	ASSESSMENT OF EFFECTS ON HUMAN HEALTH FOR THE PRODUCT	343
10.1	PRODUCT(S).....	343
10.2	DERMAL ABSORPTION.....	343
10.3	ACUTE TOXICITY	345
10.3.1	<i>Overall conclusion on acute toxicity</i>	345
10.4	CORROSION AND IRRITATION.....	346
10.4.1	<i>Skin corrosion and irritation</i>	347
10.4.2	<i>Serious eye damage and eye irritation</i>	347
10.4.3	<i>Respiratory tract irritation</i>	347
10.4.4	<i>Overall conclusion on corrosion and irritation</i>	347
10.5	SENSITISATION	349
10.5.1	<i>Skin sensitisation</i>	349
10.5.2	<i>Respiratory sensitisation</i>	349
10.5.3	<i>Overall conclusion on sensitisation</i>	350
10.6	OTHER	351
11	ENVIRONMENTAL EFFECTS ASSESSMENT FOR THE PRODUCT	352
11.1	ATMOSPHERE	352
11.2	STP	352
11.3	AQUATIC COMPARTMENT.....	352
11.4	TERRESTRIAL COMPARTMENT	352
11.5	PRIMARY AND SECONDARY POISONING	352
PART C : RISK CHARACTERISATION OF THE BIOCIDAL PRODUCT(S)		353
12	RISK CHARACTERISATION FOR HUMAN HEALTH	353
12.1	CRITICAL ENDPOINTS	353
12.1.1	<i>Systemic effects</i>	353
12.1.2	<i>Local effects</i>	357
12.1.3	<i>Absorption</i>	358
12.2	REFERENCE VALUES.....	358
12.2.1	<i>Uncertainties and assessment factors</i>	358
12.2.2	<i>AEL setting</i>	360
12.2.3	<i>Reference values to be used in Risk Characterisation</i>	366
12.2.4	<i>Maximum residue limits or equivalent</i>	367
12.3	INDUSTRIAL USES.....	368
12.4	PROFESSIONAL USES.....	369
12.4.1	<i>Systemic effects</i>	369
12.4.1.1	Combined scenarios.....	370
12.4.2	<i>Local effects</i>	371
12.4.3	<i>Conclusion</i>	379
12.5	NON-PROFESSIONAL USERS.....	382
12.6	SECONDARY (INDIRECT) EXPOSURE AS A RESULT OF USE	383
12.6.1	<i>Systemic effects</i>	383
12.6.1.1	Combined scenarios.....	383
12.6.2	<i>Local effects</i>	383
12.6.3	<i>Conclusion</i>	383
12.7	INDIRECT EXPOSURE VIA FOOD.....	384
12.8	PRODUCTION / FORMULATION OF ACTIVE SUBSTANCE	385
13	RISK CHARACTERISATION FOR THE ENVIRONMENT	386
13.1	ATMOSPHERE	386
13.2	SEWAGE TREATMENT PLANT (STP)	386

13.3	AQUATIC COMPARTMENT.....	386
13.4	TERRESTRIAL COMPARTMENT.....	387
13.5	GROUNDWATER.....	387
13.6	PRIMARY AND SECONDARY POISONING.....	388
13.6.1	<i>Primary poisoning</i>	388
13.6.2	<i>Secondary poisoning</i>	388
13.7	AGGREGATED EXPOSURE (COMBINED FOR RELEVANT EMISSION SOURCES).....	388
13.7.1	<i>STP route</i>	389
13.7.2	<i>Manure route</i>	389
13.7.2.1	Emission estimation.....	389
13.7.2.2	Fate and distribution in exposed environmental compartments.....	389
13.7.2.3	Calculated aggregated Σ PEC values.....	390
13.7.2.3.1	Manure route: refinement of the exposure calculation.....	391
13.7.2.4	Aggregated risk characterisation.....	393
13.8	SUMMARY OF THE RISK ASSESSMENT FOR THE ENVIRONMENT.....	394
14	RISK CHARACTERISATION FOR THE PHYSICO-CHEMICAL PROPERTIES.....	395
15	MEASURES TO PROTECT MAN, ANIMALS AND THE ENVIRONMENT.....	395
	PART D : APPENDICES.....	397
	APPENDIX I: LIST OF ENDPOINTS.....	397
	APPENDIX II: HUMAN EXPOSURE CALCULATIONS.....	412
	APPENDIX III: ENVIRONMENTAL EMISSION (AND EXPOSURE) CALCULATIONS.....	424
	APPENDIX IV: LIST OF TERMS AND ABBREVIATIONS.....	425
	APPENDIX V: OVERALL REFERENCE LIST.....	426
	APPENDIX VI: CONFIDENTIAL INFORMATION.....	446
	APPENDIX VII: HUMAN HEALTH – READ-ACROSS JUSTIFICATION.....	446

1 STATEMENT OF SUBJECT MATTER AND PURPOSE

This assessment report has been established as a result of the evaluation of the active substance **FORMIC ACID** in PT 3 "*Veterinary hygiene*", carried out in the context of the working programme for the review of existing active substances provided for in Article 89 of Regulation (EU) No 528/2012, with a view to the possible approval of this substance.

In accordance with the provisions of Article 7(1) of Commission Regulation (EC) No 1451/2007, Belgium was designated as Evaluator Member State to carry out the assessment on the basis of the submitted dossier.

FORMIC ACID (CAS N° 64-18-6) was notified as an existing active substance by BASF SE and KEMIRA OYJ. This notification was intended to encompass both FORMIC ACID/FA and PERFORMIC ACID/PFA *in situ*-generated (from formic acid and hydrogen peroxide) in which FA was considered the active substance and PFA the representative product.

- In the period 2007 to 2009, the BE eCA received the dossier and numerous updates from the two applicants (BASF SE and Kemira Oyj).

- In March 2015, it was decided according to the CA-March15-Doc.5.1 that the original review programme entry (37) for Formic Acid was to be split in two separate entries for Formic Acid and Performic Acid (generated from formic acid and hydrogen peroxide).

Subsequently, a resubmission of the dossier was necessary, since the original dossier consisted of a tightly interwoven dossier between the now two distinct substances.

In September 2015, a new dossier for Formic Acid was submitted by both applicants, who had now started working together in a Formic Acid Task Force (BASF SE, Kemira Oyj), following numerous updates in the periods 2015 to 2021.

On November 21st 2016 the BE eCA submitted a CLH dossier to ECHA. ECHA provided their accordance check on the CLH report on February 9th 2017, concluding that revisions and clarifications were required.

Before submitting the CAR to ECHA, the applicants were given the opportunity to provide written comments in line with Article 8(1) of Regulation (EU) No 528/2012.

On September 15st 2021, the BE eCA submitted to ECHA a copy of the assessment report containing the conclusions of the assessment, hereafter referred to as the competent authority report (CAR).

By the time of submitting this new CAR, according to the biocides Review Program Regulation/Biocides working procedure, a revised CLH report (addressing hazard classes that should be included to the already existing C&L) is duly submitted.

After ECHA Accordance Check, a peer-review by technical experts from all Member States of the draft CAR is organised by ECHA. The CAR is presented at the Biocidal Products Committee (and its Working Groups meetings) and thereafter amended according to the revisions agreed upon the comments received.

The aim of the assessment report is to support the opinion of the Biocidal Products Committee and a decision on the approval of **Formic Acid** for PT 3 and, should it be approved, to facilitate

the authorisation of individual biocidal products. In the evaluation of applications for product authorisation, the provisions of Regulation (EU) No 528/2012 shall be applied, in particular the provisions of Chapter IV, as well as the common principles laid down in Annex VI.

For the implementation of the common principles of Annex VI, the content and conclusions of the assessment report, which is available from the web-site of ECHA shall be taken into account.


However, where conclusions of this assessment report are based on data protected under the provisions of Regulation (EU) No 528/2012, such conclusions may not be used to the benefit of another applicant, unless access to these data for that purpose has been granted to that applicant.

2 ASSESSMENT REPORT

SUMMARY

1 PRESENTATION OF THE ACTIVE SUBSTANCE

1.1 IDENTITY OF THE ACTIVE SUBSTANCE

Main constituent(s)	
ISO name	Formic Acid
IUPAC or EC name	Methanoic Acid
EC number	200-579-1
CAS number	64-18-6
Index number in Annex VI of CLP	607-001-00-0
Minimum purity / content	Min. 99%
Structural formula	

Relevant impurities and additives		
IUPAC name or chemical name or EC name	Maximum concentration in % (w/w)	Index number in Annex VI of CLP
n.a.	n.a.	n.a.

1.2 INTENDED USES AND EFFECTIVENESS

Use of the active substance	
Product type	<p>PT2 "Disinfectants and algacides not intended for direct application to humans or animals"</p> <p>PT3 "Veterinary hygiene"</p> <p>PT4 "Food and feed area"</p> <p>PT5 "Drinking water"</p> <p>PT6 "Preservatives for products during storage"</p>
Intended use pattern(s)	<p>Formic Acid-based Biocidal products are intended to be used for :</p> <ul style="list-style-type: none"> - Disinfection of industrial and institutional premises and machinery, bathroom surfaces, toilets and sanitary ware in the domestic and institutional environment, - Disinfection of waters (including bathing and waste waters) - Disinfection of areas in which animals are housed, kept and transported.

	<ul style="list-style-type: none"> – Disinfection of working areas and production surfaces including food preparation and consumption areas. – Disinfection of drinking water for both humans and animals. – Preservation of industrial, consumer, household and institutional products.
Users	Industrial, professional and general public - depending on the product.

Effectiveness of the active substance	
Function	Disinfectant Preservative
Organisms to be controlled	To kill microorganisms in general : bacteria, yeasts and fungi
Limitation of efficacy including resistance	<ul style="list-style-type: none"> – Avoid formulating with, or combining with, ingredients with a strongly alkali pH value : The antimicrobial effectiveness of Formic Acid-based Biocidal products is reduced with increasing system pH and users should take this into account, particularly at pH above 4.5. – Resistance against the mode of action is unlikely to occur, i.e. there is no adaptation to cope with acidic pH values or denaturated proteins, nor is there a mechanism known to exist that a sub-lethal energy supply, due to an incomplete cytochrome C oxidase inhibition, would lead to undesired side-effects or resistance against this inhibitor. – To prevent potential development of resistance or tolerance the use of sub-lethal dosing levels should be avoided. <p>No incidence of resistance to formic acid has been recorded until now.</p>
Mode of action	Two different modes of action are reasonably considered to contribute to the biocidal activity, i.e. acidulant action and corrosion which causes enzyme denaturation and inhibition, cellular structure disruption, and impairment of cellular metabolic pathways. This mode of action is considered to depend on the low pH-value. Secondly, formic acid does inhibit cytochrome C oxidase and thus impairs cellular energy supply. Organisms and tissues with a high energy demand are specifically susceptible.

1.3 CLASSIFICATION AND LABELLING

1.3.1 Classification and labelling for the active substance

Hazard class/ property	Proposed classification
Physical hazards	
Explosives	The active substance is not an explosive
Flammable gases	Not applicable as the active substance is a liquid
Flammable aerosols	Not applicable as the active substance is a liquid
Oxidising gases	Not applicable as the active substance is a liquid
Gases under pressure	Not applicable as the active substance is a liquid
Flammable liquids	Classified as Flam liquid 3 due to the flash point being under 60°C (49°C)
Flammable solids	Not applicable as the active substance is a liquid
Self-reactive substances	The substance is not self-reactive.
Pyrophoric liquids	Not pyrophoric liquid based on auto-ignition temperature and experience in manufacture and handling.
Pyrophoric solids	Not applicable as the active substance is a liquid
Self-heating substances and mixtures	Not applicable
Substances which in contact with water emit flammable gases	Not applicable since formic acid can be diluted in water
Oxidising liquids	Not applicable. Formic acid contains oxygen but is chemically bonded only to carbon and hydrogen.
Oxidising solids	Not applicable as the active substance is a liquid
Organic peroxides	Not applicable as formic acid is not an organic peroxides as it does not contain the bivalent -O-O structure.

Corrosive to metals	Corrosive to steel. Not corrosive to aluminium.
Human health hazards	
Acute toxicity via oral route	Acute Tox. 4, H302 Harmful if swallowed
Acute toxicity via dermal route	Data lacking
Acute toxicity via inhalation route	Acute Tox. 3, H331 Toxic if inhaled EUH071
Skin corrosion/irritation	Skin Corr. 1A, H314
Serious eye damage/eye irritation	Eye Dam. 1, H318 Causes serious eye damage
Respiratory sensitisation	Conclusive but not sufficient for classification
Skin sensitisation	Conclusive but not sufficient for classification
Germ cell mutagenicity	Conclusive but not sufficient for classification
Carcinogenicity	Conclusive but not sufficient for classification
Reproductive toxicity	Conclusive but not sufficient for classification
Specific target organ toxicity-single exposure	Conclusive but not sufficient for classification
Specific target organ toxicity-repeated exposure	Conclusive but not sufficient for classification
Aspiration hazard	Conclusive but not sufficient for classification
Environmental hazards	
Hazardous to the aquatic environment	Conclusive but not sufficient for classification
Hazardous to the ozone layer	Hazard class not assessed

1.3.1.1 **CURRENT CLASSIFICATION AND LABELLING**

Current Classification and Labelling according to Regulation (EC) No 1272/2008		
Classification	Labelling	

Hazard Class and Category	Hazard statements	Pictograms	Signal word	Hazard statements	Suppl. Hazard statements	Precautionary statements	SCLs and M-factors
Skin Corr. 1A	H314	GHS05	danger	H314	-	(-)	<u>Skin Corr. 1B;</u> <u>H314:</u> 10% ≤ C < 90% <u>Skin Corr. 1A;</u> <u>H314:</u> C ≥ 90% <u>Skin Irrit. 2;</u> <u>H315:</u> 2% ≤ C < 10% <u>Eye Irrit. 2;</u> <u>H319:</u> 2% ≤ C < 10%

1.3.1.2 **PROPOSED CLASSIFICATION AND LABELLING**

Proposed Classification and Labelling according to Regulation (EC) No 1272/2008							
Classification		Labelling					
Hazard Class and Category	Hazard statements	Pictograms	Signal word	Hazard statements	Suppl. Hazard statements	Precautionary statements	SCLs and M-factors
Corrosive to metal	H290 - May be corrosive to metals	GHS05	warning	May be corrosive to metals	-	P234 P390 P406	-
Flammable liquid – category 3	H226 - Flammable liquid and vapour	GHS02	Warning	Flammable liquid and vapour		P210 P233 P240 P242	

						P243 P280 P303+P361+P353 P403+P235 P501	
Acute tox. 4 (oral)	H302	GHS07	warning	H302	-	Prevention P264, P270 Disposal P501	
Acute tox. 3 (Inhalation - vapour)	H331	GHS06	danger	H331	EUH071	Prevention P261, P271 Response P304+P340, P311 Storage P403+P233, P405 Disposal P501	
Skin Corr. 1A	H314	GHS05	danger	H314	-	Prevention P280, P260, P264 Response P310, P305+P351+P338, P304+P340, P303+P361+P353, P301+P330+P331, Storage P405 Disposal P501	<u>Skin Corr. 1B; H314:</u> 10% ≤ C < 90% <u>Skin Corr. 1A; H314:</u> C ≥ 90% <u>Skin Irrit. 2; H315:</u> 2% ≤ C < 10%
Eye dam./Irrit. 1	H318	-	-	-	-	Prevention H280 Response P310, P305+P351+P338	Eye dam./Irrit. 1; H318: C ≥ 10% <u>Eye Irrit. 2; H319:</u> 2% ≤ C < 10%

1.3.2 Classification and labelling for the representative product(s)

1.3.2.1 PROPOSED CLASSIFICATION AND LABELLING

Proposed Classification and Labelling according to Regulation (EC) No 1272/2008							
Protectol® FM 85							
Classification		Labelling					
Hazard Class and Category	Hazard statements	Pictograms	Signal word	Hazard statements	Suppl. Hazard statements	Precautionary statements	SCLs and M-factors
Corrosive to metal	H290 - May be corrosive to metals	GHS05	warning	May be corrosive to metals	-	P234 P390 P406	-
Acute tox. 4 (oral)	H302	GHS07	warning	H302	-	Prevention P264, P270 Disposal P501	
Acute tox. 3 (Inhalation - vapour)	H331	GHS06	danger	H331	EUH071	Prevention P261, P271 Response P304+P340, P311 Storage P405 P403+P233 Disposal P501	
Skin Corr. 1B	H314	GHS05	danger	H314	-	Prevention P280, P260, P264 Response P310, P305+P351+P338, P304+P340, P303+P361+P353, P301+P330+P331, Storage P405	Skin Corr. 1B; H314 if 10% ≤ C < 90%

						Disposal P501	
Eye dam./Irrit. 1	H318	-	-	-	-	Prevention P280 Response P310, P305+P351+P338	Eye dam./Irrit. 1; H318: C ≥ 10%

1.3.2.2 **PACKAGING OF THE BIOCIDAL PRODUCT**

Protectol® FM 85					
Type of packaging	Size/volume of the packaging	Material of the packaging	Type and material of closure(s)	Intended user (e.g. Professional, non-Professional)	Compatibility of the product with the proposed packaging materials (Yes/No)
IBC; Drum, Sample bottles	1050 L(IBC), 220 L (Drums) 1L (bottles)	PE (outer container corrosion resistant steel) or brown glass (bottles)	PE	professional	yes

2 SUMMARY OF THE HUMAN HEALTH RISK ASSESSMENT

2.1 SUMMARY OF THE ASSESMENT OF EFFECTS ON HUMAN HEALTH

Introductory note:	
<p>The repeated dose toxicity via the oral route of formic acid is assessed with its non-corrosive salts, sodium formate and potassium diformate, in order to achieve sufficiently high dose levels. Neurotoxicity is assessed with methanol. This read across approach is in accordance with Article 6(3) of the EU No. 528/2012 (BPR) following point 1.5(2) under Annex IV: “<i>common precursors and/or the likelihood of common breakdown products via physical and biological processes, which result in structurally similar chemicals and indicates the presence of dangerous properties</i>”. The full read-across justification, which was performed following the Read-Across Assessment Framework developed by ECHA, can be found in Appendix VII.</p>	
Endpoint	Brief description
Toxicokinetics	<p>Absorption: rapid, but no quantitative data available</p> <p>Distribution: seemingly a significant proportion of formate distributes in the tissue, but more likely undergoes rapid metabolism and excretion</p> <p>Metabolism: rapid: hepatic first pass effect; oxidation to CO₂; no indication of accumulation</p> <p>Excretion: Rapid elimination via exhalation of CO₂; low urinary excretion of formic acid</p>
Acute toxicity	<p>predominantly determined by formic acid’s inherent irritating/corrosive properties. The toxicity values after oral uptake and inhalation in rats suggest formic acid to be acutely harmful. The clinical signs give no evidence of specific systemic adverse effects.</p> <p>Proposed classification:</p> <p>Acute tox 4 (oral) H302 LD₅₀ 730 mg/kg bw¹</p> <p>Acute tox 3 (inhal) H331 LC₅₀ 7.4 mg/l</p>
Corrosion and irritation	<p>Formic acid is corrosive to skin and eye. Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive in the EU (12th ATP). Respiratory irritation: we propose to classify formic acid as EUH071, ‘corrosive to the respiratory tract’, as its corrosive properties determine its toxicity.</p>
Sensitisation	<p>Formic acid is not a skin sensitizer. There is no indication that formic acid would be a respiratory sensitizer.</p>
Repeated dose toxicity	<p>The short-term toxicity of formic acid has not been investigated.</p> <p>The medium-term oral toxicity was studied in the rat and the pig. Oral administration of potassium diformate led to largely reversible local irritation effects in the stomach and histological changes of the stomach and gastrointestinal tract. High doses may produce adverse effects, such as decrease in body weight gain, possibly due to the inherent irritating potential.</p>

¹ Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

	<p>Rat: NOAEL_{local} < 420 mg formate/kg bw/d NOAEL_{sys} 840 mg/kg bw/d</p> <p>Pig: < 149 mg formate/kg bw/d</p> <p>Medium-term inhalation toxicity was studied in rats and mice exposed to formic acid vapours for 13 weeks. Histological changes were observed in the upper respiratory tract. In addition, a decrease in body weight gain was observed at the highest dose level in mice.</p> <p>Medium-term inhalation toxicity:</p> <p>overall NOAEC_{local} = 60 mg formic acid/m³ NOAEC_{systemic} = 244 mg formic acid/m³</p> <p>The long-term oral toxicity was studied in the rat and the pig. Oral administration of potassium diformate led to local irritation effects in the stomach, which were confirmed histo-pathologically. In the high dose animals, body weight (gain) was decreased and there was a lower incidence of pelvic mineralization in the kidney.</p> <p>NOAEL_{systemic} = 280 mg formate/kg bw/d</p>
Genotoxicity	Available data indicate that formic acid has no genotoxic potential.
Carcinogenicity	No data are available on formic acid. A carcinogenicity study on potassium diformate indicates that potassium diformate has no carcinogenic potential.
Reproductive toxicity	<p>No data are available on formic acid.</p> <p>No developmental toxicity and teratogenicity was observed for formate in rats and rabbits.</p> <p>No adverse effects on fertility were observed for formate in rats.</p> <p>No adverse effects on or via lactation are expected for formic acid.</p> <p>Two-generation study, rat:</p> <p>NOAEL_{parental} = 200 mg formate/kg bw/d NOAEL_{offspring} = 670 mg formate/kg bw/d NOAEL_{reproduction parameters} = 670 mg formate/kg bw/d</p> <p>Teratogenicity studies, rat, rabbit:</p> <p>NOAEL_{maternal} = 640 mg formate/kg bw/d NOAEL_{developmental} = 640 mg formate/kg bw/d</p>
Neurotoxicity	<p>At moderate doses, no neurotoxic effects are expected for formic acid.</p> <p>When the metabolic capacity to dispose of formate is exceeded, formate accumulation and adverse effects on the optical nerve and photoreceptors can occur. However, these symptoms are considered to be an exclusive sequel of acute methanol intoxication in primates.</p>
Immunotoxicity	There are no indications that Formic Acid has the potential to induce adverse effects involving the immune system.
Disruption of the endocrine system	ED criteria are not met for Human Health
Other effects	Workplace measurements, health records from industry and case reports show that local corrosive effects prevail but systemic effects may result after contact of concentrated formic acid to extended areas of the body

	<p>surface. Occupational and accidental dermal exposure records report skin corrosion and metabolic acidosis. After oral exposure observations range from moderate burns around the mouth to severe corrosion of the gastro-intestinal tract with destruction of the oesophagus, perforation of the stomach, and corrosion of the small intestine together with massive bleeding and systemic toxicity, potentially leading to the death of the patient. For inhalation exposure at the threshold limit of 5 ppm or 9.5 mg/m³ an effect on the blood pH is unlikely.</p>
--	---

2.2 REFERENCE VALUES

	Study	NOAEL/ LOAEL	Overall assessment factor	Value
AEL _{short-term}	90-day feeding study, potassium diformate, rat	840 mg formate/kg bw/d (2100 mg formate/kg bw/d)	100	8.4 mg formate/kg bw/d
AEL _{medium-term}	90-day feeding study, potassium diformate, rat	840 mg formate/kg bw/d (2100 mg formate/kg bw/d)	100	8.4 mg formate/kg bw/d
AEL _{long-term}	2-year feeding study, potassium diformate, rat	280 mg formate/ kg bw/d (1400 mg formate/kg bw/d)	100	2.8 mg formate/kg bw/d Rounded to 3 mg formate/kg bw/d ²
ARfD	Not required	/	/	/
ADI	EU SANCO D3/AS D, 2005; JECFA, 2003	/	/	3 mg/kg bw/d
Occupational exposure limit	EU WEL, MAK/TLV (8-hour TWA) IOELV Commission Directive 2006/15/EC	/	/	5 ppm or 9.5 mg/m ³ 5 ppm or 9 mg/m ³
AEC _{resp tract irrit}	inhalation, 13 weeks, formic acid, rat/mice	60 mg/m ³	10	6 mg/m ³

² We refer to TAB entry TOX-4 as the impact of rounding is less than 10%. Please note that for this CAR, the risk characterization has been performed with the non-rounded 2.8 mg formate/kg bw/d value. The decision for rounding the AEL long-term was taken at HH WG I-2022; however it was decided that there was no need to alter the risk characterization of the CAR. For product approval, the rounded 3 mg formate/kg bw/d value should be used.

2.3 RISK CHARACTERISATION

Summary table: scenarios			
Scenario number	Scenario (e.g. mixing/loading)	Primary or secondary exposure Description of scenario	Exposed group (e.g. Professional, non-Professional, bystanders)
1.	Animal house disinfection by fogging	1a.primary exposure during mixing and loading by professional contractors	Professionals: professional contractors
		1b.application of the in use solution by fogging, manipulated from outside	
		1c. cleaning of equipment and disposal of containers	
2.	Disinfection of footwear in dipping troughs	2a.primary exposure during mixing and loading by experienced farm workers in dipping troughs	Professionals: experienced farm workers
		2b.passive use of footbath/trough by experienced farm workers	
		2c.draining of disinfection solution and disposal of containers by experienced farm workers	
3.	Animal feet disinfection and disinfection of transport vehicles	3a.primary exposure during mixing and loading by experienced farm workers in animal feet baths/vehicle troughs	Professionals: experienced farm workers
		3b.passive animal use of animal feet bath, passive use of vehicle trough by experienced farm workers	
		3c.draining of disinfection solution and disposal of containers by experienced farm workers	
4.	Reuse/restocking of disinfected animal housing	Secondary exposure while opening animal housings sealed for treatment (fogging)	Professionals: experienced farm workers

The risk assessment performed for formic acid, PT3, covers professional animal house disinfection by fogging, and disinfection of footwear, animal feet and transport vehicles in dipping troughs. A deciding factor in identifying safe uses is the high vapour pressure of formic acid. Inhalation exposure to formic acid is relevant in all scenarios.

Exposure for professional animal house disinfection by fogging was assessed. The assessment includes exposure at re-entry by professional bystanders. Systemic exposure was determined for the dermal and inhalation route. A quantitative assessment was done for inhalation of vapour. Where relevant, a qualitative assessment was included for local dermal and inhalation exposure.

There is no concern for professionals using the biocidal product during fogging of animal housing PT3.1, when appropriate PPE are applied. During mixing and loading, additionally

appropriate RPE are required when ventilation is insufficient. Sufficient ventilation for safe re-entry should be ensured.

Exposures for disinfection of footwear, animal feet and transport vehicles in dipping troughs were assessed for agricultural workers. Systemic exposure was determined for the dermal and inhalation route. A quantitative assessment was done for inhalation of vapour. Where relevant, a qualitative assessment was included for local dermal and inhalation exposure.

There is no concern for professionals using the biocidal product preparing, using and emptying dipping troughs, when appropriate PPE are applied during mixing and loading. During mixing and loading, additionally appropriate RPE are required when ventilation is insufficient. For exposure to in-use dilutions of FA (skin/eye irritant), appropriate PPE should be used during post-application as well.

Possible scenario combinations were calculated for professionals (experienced farmers).

There is no concern for professionals during combined use of the biocidal product for restocking of treated animal housings, and for preparing and using dipping troughs, when PPE as discussed for single scenarios are applied.

Inhalation exposure at re-entry after fogging is acceptable only when ventilation measures are in place to reduce the FA concentration in air to below 6 mg/m³.

For the representative products in this CAR, indirect exposure of the general public should be avoided by implementation of appropriate RMM, considering the volatility and corrosive properties of the a.s., and the fact that PPE and RPE are not applicable for the general public.

The following RMM is proposed:

Use is not authorized in areas where public can be received/present.

Animal health:

Animal exposure assessment and measures for animal health are required at product authorisation stage.

Conclusion of risk characterisation for professional user				
Scenario, Tier	Relevant reference value²	Estimated uptake (syst: mg/kg bw/d; local: mg/m³)	Estimated uptake/reference value (%)	Acceptable (yes/no)
Scenario 1, fogging, T1 no PPE	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	16.431	587	no
	Local inhalation vapour AEC 6 mg/m ³	3.8	63.3	yes
Scenario 1, fogging, T2 impermeable coveralls, boots, gloves and face protection	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.0377	1.3	yes
	Local inhalation vapour AEC 6 mg/m ³	3.8	63.3	yes

Scenario 2, footwear disinfection, 5% FA, T1 no PPE	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.212	7.6	yes
	Local inhalation vapour AEC 6 mg/m ³	3.8	63.3	yes
Scenario 2, footwear disinfection, 5% FA, T2 coated coveralls, boots, gloves and face protection;	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.0329	1.2	yes
	Local inhalation vapour AEC 6 mg/m ³	3.8	63.3	yes
Scenario 3, animal feet/transport vehicle disinfection, 5% FA, T1 no PPE	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.0475	1.7	yes
	Local inhalation vapour AEC 6 mg/m ³	3.5 (M&L), 1.0 (appl), 0.38 (post-appl)	58.3 (M&L), 16.7 (appl), 6.3 (post- appl)	yes
Scenario 3, animal feet/transport vehicle disinfection, 5% FA, T2 coveralls, boots, gloves and face protection	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.0228	0.8	yes
	Local inhalation vapour AEC 6 mg/m ³	3.5 (M&L), 1.0 (appl), 0.38 (post-appl)	58.3 (M&L), 16.7 (appl), 6.3 (post- appl)	yes
Scenario 4, restocking of animal housing after ambient temperature fogging, T1, all a.s. evaporated	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	11.5	411	no
	Local inhalation vapour AEC 6 mg/m ³	550	9167	no
Scenario 4, restocking of animal housing after ambient temperature fogging, T2/3, ventilation until <AEC	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.125	4.5	yes
	Local inhalation vapour AEC 6 mg/m ³	<6	<100	yes
Scenario 4, restocking of animal housing after thermal fogging, T1, all a.s. evaporated	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	9	321	no
	Local inhalation vapour AEC 6 mg/m ³	430	7167	no

Scenario 4, restocking of animal housing after thermal fogging, T2/3, ventilation until <AEC	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.125	4.5	yes
	Local inhalation vapour AEC 6 mg/m ³	<6	<100	yes
Scenarios 2,3,4 restocking after ambient temperature fogging, footwear, animal feet disinfection, T1	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	11.76	420	no
	Local inhalation vapour AEC 6 mg/m ³	550	9167	no
Scenarios 2,3,4 restocking after ambient temperature fogging, footwear, animal feet disinfection, T2	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.181	6.5	yes
	Local inhalation vapour AEC 6 mg/m ³	<6	<100	yes
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection, T1	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	9.26	331	no
	Local inhalation vapour AEC 6 mg/m ³	430	7167	no
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection, T2	Systemic effects AEL _{long-term} 2.8 mg/kg bw/d	0.181	6.5	yes
	Local inhalation vapour AEC 6 mg/m ³	<6	<100	yes
Local exposure				
conc	task	classification	Hazard category	Potential exposure route
55%	Mixing & loading For fogging, footwear disinfection, animal feet/vehicle transport disinfection	Skin corr 1B EUH071	High	Skin, eye, RT
	Conclusion on risk: ACCEPTABLE +engineering controls +low frequency +short duration			

	+professionals using PPE; RPE @insufficient ventilation +professionals following instructions for use +good standard of personal hygiene			
5%	Dipping troughs Draining & disposal	Skin irrit 2 Eye irrit 2	low	Skin, eye, RT
Conclusion on risk: ACCEPTABLE +engineering controls +reversible effect +professionals following instructions for use +experience expected				

3 SUMMARY OF THE ENVIRONMENTAL RISK ASSESSMENT

3.1 FATE AND BEHAVIOUR IN THE ENVIRONMENT

Summary table on compartments exposed and assessed		
Compartment	Exposed (Y/N)	Assessed (Y/N)
Freshwater	Y	Y
Sediment	N	N
Seawater	N	N
Seawater sediment	N	N
STP	Y	Y
Air	Y	N
Soil	Y	Y
Groundwater	Y	Y
Biota	N	N

Summary table on relevant physico-chemical and fate and behaviour parameter of the active substance			
	Value	Unit	Remarks
Molecular weight	46.03	g/mol	
Melting point	8	°C	
Boiling point	100.23	°C	
Vapour pressure (at 12 °C)	2400	Pa	
Water solubility (at 12 °C)	1.09x10 ⁶	mg/l	
Log10 Octanol/water partition coefficient	-2.10	---	(pH 7)
Organic carbon/water partition coefficient (Koc)	30	l/kg	(pH 7)
Henry's Law Constant (at 12 °C)	0.101	Pa/m ³ /mol	
Acid dissociation constant	3.7	---	Predominant species at a pH of 7 is formate, which is reflected in the pH dependent Koc.
Biodegradability	Ready biodegradable		
DT50 for degradation in soil (12 °C)	1	day	
Half-life for biodegradation in manure	19.9	d	(12 °C)

3.2 EFFECTS ASSESSMENT

Summary table on calculated PNEC values	
Compartment	PNEC
Freshwater	≥ 2 mg/L
STP	> 50 mg/L
Soil	≥ 1.29 mg/kg _{wwt} (≥ 1.47 mg/kg _{dwt})
Groundwater	Not applicable

For groundwater, calculated PECs are compared to the reference value of 0.1 µg/L.

3.3 EXPOSURE ASSESSMENT

Whenever applicable, PEC values were calculated for all stable types. Only the maximum values are presented in the summary table below.

Summary table on calculated PEC values				
	PEC _{STP}	PEC _{water}	PEC _{soil,twa}	PEC _{GW} ¹
	[mg/L]	[mg/L]	[mg/kg _{wwt}]	[µg/L]
Scenario 1 (Footwear)	2.00E-02	7.285E-03	2.828E-01	7.285E+01
Scenario 2 (Animal's feet)	2.43E+00	2.43E-01	2.527E-01	6.508E+01
Scenario 3 (Animal housing, fogging)	5.49E-02	5.49E-03	2.238E-03	2.757
1 TIER 1: porewater concentration				

3.4 RISK CHARACTERIZATION

Whenever applicable, PEC/PNEC values were calculated for all stable types. Only the maximum values are presented in the summary table below.

Summary table on calculated PEC/PNEC values			
	PEC/PNEC_{STP}	PEC/PNEC_{water}	PEC/PNEC_{soil}
Scenario 1 (Footwear)	< 4.00x10 ⁻⁴	≤ 3.64x10 ⁻³	≤ 0.219
Scenario 2 (Animal's feet)	< 4.85x10 ⁻²	≤ 1.22x10 ⁻¹	≤ 0.196
Scenario 3 (Animal housing, fogging)	< 1.10x10 ⁻³	≤ 2.75x10 ⁻³	≤ 1.73x10 ⁻³

Groundwater concentrations for all scenarios were below the threshold of 0.1 µg/L after refinement of the exposure calculation.

Conclusion:

The risks for the environment from the intended uses of the representative product for PT3 are acceptable.

4 ASSESSMENT OF EXCLUSION, SUBSTITUTION CRITERIA AND POP

Conclusion on exclusion criteria	The exclusion criteria in BPR Article 5(1)a-c are not met.
Conclusion on CMR	Formic acid is not classified and does not meet the criteria to be classified as CMR
Conclusion on ED assessment	Formic acid does not have endocrine disrupting activities.
Conclusion on PBT and vP/vB criteria	Formic acid is not a PBT/vPvB substance.

Conclusion on substitution criteria	The substitution criteria in BPR Article 10(1)a-f are not met.
--	--


Conclusion on LRTAP/POP assessment	Formic acid does not meet the criteria for being a POP or LRTAP.
---	--

PART A : ASSESSMENT OF INTRINSIC PROPERTIES AND EFFECTS OF THE ACTIVE SUBSTANCE

1 GENERAL SUBSTANCE INFORMATION

1.1 IDENTIFICATION OF THE SUBSTANCE

Summary table on substance identity	
Common name (ISO name, synonyms)	Formic Acid
Chemical name (EC name, CA name, IUPAC name)	Methanoic Acid
EC number	200-579-1
CAS number	64-18-6
other CAS numbers (e.g. deleted, related, preferred, alternate)	/
Molecular formula	CH ₂ O ₂
SMILES notation	C(=O)O
Molar mass	46.025 g/mol
Information on optical activity and typical ratio of (stereo) isomers (if applicable and appropriate)	Not relevant
Description of the manufacturing process and identity of the source (for UVCB substances only)	Not relevant
Degree of purity (%)	Min. 99%

Structural formula


Origin of the natural active substance or precursor(s) of the active substance
Please refer to BASF PT3 Confidential Annex.

Method of manufacture
Please refer to BASF PT3 Confidential Annex.

1.2 COMPOSITION OF THE SUBSTANCE (REFERENCE SPECIFICATIONS)

Main constituent(s)			
Constituent (chemical name)	Typical concentration (%(w/w))	Concentration range (%(w/w))	Remarks / Discussion
Formic Acid	Please refer to BASF PT3 Confidential Annex.		

Impurities			
Constituent (chemical name)	Typical concentration (%(w/w))	Concentration range (%(w/w))	Remarks / Discussion
Please refer to BASF PT3 Confidential Annex.			

Additives				
Constituent (chemical name)	Function	Typical concentration (%(w/w))	Concentration range (%(w/w))	Remarks / Discussion
Please refer to BASF PT3 Confidential Annex.				

1.3 PHYSICAL AND CHEMICAL PROPERTIES OF THE ACTIVE SUBSTANCE

Property	Result	Test method applied or description in case of deviation	Remarks / Discussion / Justification for waiving	References
Aggregate state at 20°C and 101.3 kPa (99.4% (w/w))	The substance is a clear and colorless liquid which is homogeneous at 20 °C and 101.3 kPa.	Organoleptic	/	Study no. 07L00084, [REDACTED], (2007)
Physical state (appearance) at 20°C and 101.3 kPa (99.4% (w/w))	Liquid	Organoleptic	/	Study no. 07L00084, [REDACTED], (2007)
Colour at 20°C and 101.3 kPa (99.4% (w/w))	Colourless	Organoleptic	/	Study no. 07L00084, [REDACTED], (2007)
Odour at 20°C and 101.3 kPa 1. 99-100% 2. 85%	Pungent	Organoleptic	/	3. BASF AG (2007) BPD ID B3_04 4. [REDACTED], (2007a)
Melting / freezing point (99.4% (w/w))	8 °C	OECD 102	No decomposition observed	[REDACTED], (2018) 20181112_07L00084 Amendment01 Final Report BPD_ID_A3_01.pdf
Boiling point at (99.4% (w/w))	100.23 °C	OECD 103	Obtained by interpolation	Study no. 07L00084, [REDACTED], (2007)
Relative density (99.4% (w/w))	D ₄ ²⁰ = 1.2195	OECD 109	/	Study no. 07L00084, [REDACTED], (2007)

Acidity/alkalinity	pH _{85% formic acid} = -1.6 At 1%: pH = 2.2	German Industrial Standard DIN 19268	Potentiometric measurement	Study no. 07L00172, [REDACTED], (2007)
	90.9530 ± 0.0663 % acidity	CIPAC MT 191	On 85% formic acid in water sample. Since test item is an acid, only acidity was tested.	Study no 16011907G975, [REDACTED], (2016a)
	pH = 2.18	CIPAC MT 75	At 24.8 °C On 1% aqueous solution of 85% formic acid sample	Study no 16011907G907, [REDACTED], (2016c)
	pH = 2.13	CIPAC MT 75.3	At 19.1 °C On 1% aqueous solution of 99% formic acid sample	Study no. S16-06389 [REDACTED], (2017)
	108.03% (m/m) mean acidity	CIPAC MT 191	On 99% formic acid	Study no. S16-06390 [REDACTED], (2017)
Vapour pressure (99.4% (w/w))	At 20 °C: 42.71 hPa At 25 °C: 54.96 hPa At 50 °C: 170.7 hPa	OECD 104	Extrapolated from regression-derived equation	Study no. 07L00084, [REDACTED], (2007)
Henry's law constant	At 20 °C: 0.16 Pa.m ³ /mol		Calculation based on measured relative density as a surrogate for water solubility and measured vapour pressure	ECT Oekotoxikologie GmbH (2015)
Surface tension (99.4% (w/w))	At 20 °C: 71.5 mN/m (at 1g/L)	OECD 115	The test item is not surface active	Study no. 07L00084, [REDACTED], (2007)

Water solubility at 20 °C	Completely miscible Corresponding to 1220 g/L (= D ₄ ²⁰)	SOP PCE/006/04 (BASF AG, GKA Analytik, chapter 4: visual method) Based on OECD 105 Deviation: Preparation of saturated solution was not possible and results are not expected to be different since missing part of the test solution for a pure solution is water.	At pH 5 / 7 / 9 At 20.1 ± 0.1 °C Temperature dependence was not investigated due to complete miscibility.	Study no. 02L00109, ██████████, (2002)
Partition coefficient (n-octanol/water) and its pH dependency Surface tension at 20 °C	At pH 5: Log K _{ow} = -1.9 At pH 7: Log K _{ow} = -2.1 At pH 9: Log K _{ow} = -2.3	EC method A.8	At 23 ± 1 °C The purity of the test solution (performed on a 85.3 w/w solution including water as "impurity") is seen as not relevant, and is not expected to influence the outcome	Study no. 02L00109, ██████████, (2002)
Thermal stability and identity of breakdown products (99.4% (w/w))	Decomposition onset temperature: 350 °C Energy release: >150 J/g Auto-ignition temperature: 528 °C (corrected according to EN 14522)	OECD 113 EC method A.15	Combustion products are H ₂ O and CO ₂ At room temperature and during incomplete combustion CO and H ₂ may be formed	Study no. SIK-Nr.07/1018, ██████████, (2007)
Reactivity towards container material (99.4% (w/w))	Compatible: - stainless steel, types 1.4306, 1.4307, 1.4311, 1.4404, 1.4541, 1.4571	Based on experience	Formic acid and solutions of formic acid are acidic. Therefore, materials which are not sufficiently resistant towards acids should not be	██████████, (2007a)

	- plastics: different types of PE like HD-PE; PP (for plugs and caps) Not compatible: - carbon steel, paper, board		used to avoid equipment damage and spoilage of products	
Dissociation constant (99.4% (w/w))	At 20 °C: pK _a = 3.70	OECD 112	/	Study no. 07L00084, ██████████, (2007)
Granulometry	Waived	-	Not applicable, substance is not a powder or granule	-
Viscosity (capillary viscometer) (99.4% (w/w))	Dynamic viscosity At 20 °C: 1.80 mPa.s At 40 °C: 1.22 mPa.s Kinematic viscosity At 20 °C: 1.47 mm ² /s At 40 °C: 1.02 mm ² /s	OECD 114	/	Study no. 07L00084, ██████████, (2007)
Solubility in organic solvents, including effect of temperature on solubility (99.4% (w/w))	Miscible at ratios: 1:9, 1:1 and 9:1 Miscible at 20 and 30 °C Corresponding to: > 850 g/L N,N-dimethylformamide > 92.9 g/L 1,4-dioxane > 1190 g/L Dichloromethane	SOP PCE/006/04 (BASF AG, GKA Analytik, chapter 4: visual method) Based on OECD 105 Deviation: Preparation of saturated solution was not possible and results are not expected to be different since missing part of the test solution for a pure solution is water.	3 solvents: - N,N-dimethylformamide (Density 0.9445 g/cm ³ at 25°C) - 1,4-dioxane (Density 1.0329 g/cm ³ at 20°C) - Dichloromethane (Density 1.3255 g/cm ³ at 20°C)	Study no. 07L00084, ██████████, (2007)

Stability in organic solvents used in biocidal products and identity of relevant degradation products	Waived	-	Organic solvents not used in the biocidal products	-
---	--------	---	--	---

1.4 PHYSICAL HAZARDS AND RESPECTIVE CHARACTERISTICS

Property	Result	Test method applied or description in case of deviation	Remarks / Discussion / Justification for waiving	References
Explosives	The substance is not explosive	UN Manual of Tests and Criteria (2010)	The substance has no chemical groups indicating explosive properties	██████████, (2006)
Flammable gases	Waived	-	Not applicable	-
Flammable aerosols	Waived	-	Not applicable	-
Oxidising gases	Waived	-	Not applicable	-
Gases under pressure	Waived	-	Not applicable	-
Flammable liquids	Flash point = 49.5 °C Is a flammable liquid category 3, as its flash point is ≥ 23 °C and ≤ 60 °C (H226)	EC method A.9	Closed cup; corrected for atmospheric pressure and rounded to units of 0.5 °C	Study no. SIK-Nr.07/1018, ██████████, (2007)
Flammable solids	Waived	-	Not applicable	-


Self-reactive substances and mixture	The substance is not self-reactive	UN Manual of Tests and Criteria (2010)	The substance has no chemical groups indicating explosive or self-reactive properties	-
Pyrophoric liquids	Waived	-	Not a pyrophoric liquid, based on auto-ignition temperature (528 °C) and experience in manufacture and handling	Study no. SIK-Nr.07/1018, [REDACTED], (2007)
Pyrophoric solids	Waived	-	Not applicable	-
Self-heating substances and mixtures	Waived	-	Not applicable, substance has a melting point of 8 °C	-
Substances and mixtures which in contact with water emit flammable gases	Waived	-	The active substance is a weak acid that, in presence of water, will partially dissociate and provide ions (ion hydronium and formate). This dissociation do not liberate any flammable gas. This is a well known process.	-
Oxidising liquids	The substance is not an oxidising liquid	UN Manual of Tests and Criteria (2010)	The compound contains oxygen but this element is chemically bonded only to carbon and hydrogen The compound does not contain any halogen atoms	[REDACTED] (2006)
Oxidising solids	Waived	-	Not applicable	-
Organic peroxide	Waived	-	Not applicable	-

Corrosive to metals	Corrosive to steel Not corrosive to aluminium	UN Test C.1 (37.4)	On 99% formic acid	Study no. 16092902G979 ██████████, (2017) Study no 16011907G979 ██████████, (2016b)
	Corrosive to steel Not corrosive to aluminium	UN Test C.1 (37.4)	On 85% formic acid in water sample	Study no 16011907G979 ██████████, (2016b)
	Compatible materials: - stainless steel, types 1.4306, 1.4307, 1.4311, 1.4404, 1.4541, 1.4571 Not compatible: - carbon steel As a conclusion, a classification as Corrosive to metals (H290) is justified.	Based on experience	On 99% formic acid	██████████, (2007a)
Auto-ignition temperature (liquids and gases)	Auto-ignition temperature: 528 °C (corrected according to EN 14522)	EC method A.15	/	Study no. SIK-Nr.07/1018, ██████████, (2007)
Relative self-ignition temperature for solids	Waived	-	Not applicable	-
Dust explosion hazard	Waived	-	Not applicable	-

1.5 HAZARD IDENTIFICATION FOR PHYSICO-CHEMICAL PROPERTIES

Formic acid is a flammable liquid category 3. Further it does not present any other hazard from a physico-chemical point of view with regard to the available information. It presents a high self-ignition temperature, and has no explosive or oxidising properties.

1.6 ANALYTICAL METHODS FOR DETECTION AND IDENTIFICATION



Analytical methods								
Analyte (type of analyte e.g. active substance, metabolite etc.)	Compartment	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ), Maximum Residue Limits or other limits	Reference
				Fortification range / Number of measurements	Mean	RSD		
<i>Titration with sodium hydroxide solution 1mol/L Formic Acid</i>	<i>Pure 100%</i>	5 concentrations >0.9999 0.2-1 g test item	No interfering substances	5 repl recovery excellent	-	-	No LoD	 (2017)
	<i>Diluted with water 85%</i>	5 concentrations >0.9999 0.2-1 g test item	No interfering substances	5 repl recovery excellent	-	-	No LoD	Idem

GCMS (column: DB FFAP, 30 m x 0.32 mm (inner diameter), film thickness 0.25 µm, batch: USN526534H , AGILENTInsert) Electron impact (EI) positive	-	-	GC-MS analysis of the test item was performed and showed the absence of any other acid or impurity that could interfere with the titration.	-	-	-	-	Idem
CIPAC Method MT 30.5, "Water, Karl Fischer method using pyridine-free reagents", Hydranal- Composite 1, titer 0.8 - 1.2 mg/mL	Pure 100% Diluted with water 85%	12 conc R=1.0000 0.118- 6.347% w/w water		12 replicates	103%	3.37%	LoQ = 0.122% w/w	Idem

<p>Remark: The enzymatic method of determination of formic acid in aqueous solutions is acknowledged to represent a specific, sensitive, and reliable method, and in Germany it is contained in the official list of methods which are suited to examine foodstuffs.</p> <p><i>Photometer (wavelength 334, 340, or 360 nm) to detect formation of NADH</i></p>	<p><i>Soil</i></p>	<p><i>Linearity is given in the range 0.2 mg formic acid /l sample solution to 200 mg formic acid/l sample (cf. full test description in Section A4.1_01).</i></p>	<p><i>Enzyme is highly specific for formic acid. Test may be disturbed by: Low or high pH outside approx. 7-8 Reducing agents Colour, turbidity, or protein</i></p>	<p>25</p>	<table border="1"> <thead> <tr> <th>Fortification [mg/kg]</th> <th>Concentration, mean [mg/kg soil]</th> <th>Recovery [%]</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>1.59</td> <td>n.d.</td> </tr> <tr> <td>5</td> <td>1.61</td> <td>31</td> </tr> <tr> <td>10</td> <td>9.05</td> <td>85</td> </tr> <tr> <td>50</td> <td>47.8</td> <td>93</td> </tr> </tbody> </table>	Fortification [mg/kg]	Concentration, mean [mg/kg soil]	Recovery [%]	0	1.59	n.d.	5	1.61	31	10	9.05	85	50	47.8	93	<p>LoQ was set at 10 mg/L for soil extracts</p>	<p>██████████ (2007)</p>
					Fortification [mg/kg]	Concentration, mean [mg/kg soil]	Recovery [%]															
					0	1.59	n.d.															
					5	1.61	31															
					10	9.05	85															
					50	47.8	93															
					<table border="1"> <thead> <tr> <th>Fortification [mg/kg]</th> <th>Concentration, mean [mg/kg soil]</th> <th>Rel SD [%]</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>1.59</td> <td>70</td> </tr> <tr> <td>5</td> <td>1.61</td> <td>53</td> </tr> <tr> <td>10</td> <td>9.05</td> <td>3.8</td> </tr> <tr> <td>50</td> <td>47.8</td> <td>4.7</td> </tr> </tbody> </table>	Fortification [mg/kg]	Concentration, mean [mg/kg soil]	Rel SD [%]	0	1.59	70	5	1.61	53	10	9.05	3.8	50	47.8	4.7		
					Fortification [mg/kg]	Concentration, mean [mg/kg soil]	Rel SD [%]															
					0	1.59	70															
					5	1.61	53															
10	9.05	3.8																				
50	47.8	4.7																				

Photometer (wavelength 334, 340, or 360 nm) to detect formation of NADH	Water surface	Surface water: given in the range 0.2 to 5 mg/L. $R^2=0.99998$ for the regression curve for all measurements Linearity confirmed in the range 0.2-100 mg/L (Keller and Hartmann, 2013: cf. Section A4.1_03).	None (enzyme specific for formic acid)	(5 measurements at each of the four fortification levels) and blanks	Fortification level [mg/L]	Recovery [%] Drinking water	Recovery [%] Surface water	LoQ of 0.2 mg/L	[REDACTED] (2007)
					0.2	103	116		
					0.5	91	n.d.		
					2	103	81		
					5	101	78		
					Fortification level [mg/L]	Rel SD[%] Drinking water	Rel SD [%] Surface water		
					0.2	17	7.7		
					0.5	2.4	n.d.		
					2	6.6	1.6		
					5	3.7	1.7		

Photometer (wavelength 334, 340, or 360 nm) to detect formation of NADH	Drinking water	Drinking water: given in the range 0.2 to 5 mg/L. $R^2 = 0.9997$ for the regression curve for all measurements Linearity confirmed in the range 0.2-100 mg/L (Keller and Hartmann, 2013: cf. Section A4.1_03)	None (enzyme specific for formic acid) Drinking water: precipitation of magnesium phosphate caused turbidity that was removed by filtering the solution.	(5 measurements at each of the four fortification levels)	Fortification level [mg/L]	Recovery [%] Drinking water	Recovery [%] Surface water	LoQ of 0.2 mg/L	[REDACTED] (2007)
					0.2	103	116		
					0.5	91	n.d.		
					2	103	81		
					5	101	78		
					Fortification level [mg/L]	Rel SD[%] Drinking water	Rel SD [%] Surface water		
					0.2	17	7.7		
					0.5	2.4	n.d.		
					2	6.6	1.6		
					5	3.7	1.7		

<p>Ion chromatography</p> <p><i>Material and conditions:</i></p> <p><i>Ion chromatographer DIONEX DX 120 with conductivity detector and autosampler.</i></p>	Air	Formic acid, 1.2 to 47.8 mg/L	Methoxyacetic acid cannot be completely separated from formic acid	Measures were performed at three different concentration (6 replicates by concentration):	<table border="1"> <thead> <tr> <th>Formic acid [mg/m³]</th> <th>Recovery [%]</th> </tr> </thead> <tbody> <tr> <td>0.9</td> <td>95</td> </tr> <tr> <td>9.0</td> <td>95</td> </tr> <tr> <td>18.0</td> <td>94</td> </tr> </tbody> </table>	Formic acid [mg/m ³]	Recovery [%]	0.9	95	9.0	95	18.0	94	<table border="1"> <thead> <tr> <th>Formic acid [mg/m³]</th> <th>Relative standard deviation [%]</th> </tr> </thead> <tbody> <tr> <td>0.9</td> <td>9,7</td> </tr> <tr> <td>9.0</td> <td>6,4</td> </tr> <tr> <td>18.0</td> <td>3,8</td> </tr> </tbody> </table>	Formic acid [mg/m ³]	Relative standard deviation [%]	0.9	9,7	9.0	6,4	18.0	3,8	<p><i>Absolute limit of quantification: 0.1 µg formic acid. This corresponds to a relative limit of quantification of 0.12 mg/m³ for an air sample volume of 140 L, an absorption volume of 10 mL, and an injection volume of 50 µL</i></p>	 (2007)
Formic acid [mg/m ³]	Recovery [%]																							
0.9	95																							
9.0	95																							
18.0	94																							
Formic acid [mg/m ³]	Relative standard deviation [%]																							
0.9	9,7																							
9.0	6,4																							
18.0	3,8																							
<p>Photometer (wavelength 334, 340, or 360 nm) to detect formation of NADH</p>	<i>animal and human body fluids and tissues</i>	<i>Linearity is given in the range 0.2 mg formic acid/l sample solution to 200 mg formic acid/l sample</i>	None (enzyme specific for formic acid)	n.a.	<i>100% because formic acid is water soluble and the volatility is low. The enzyme reaction is complete under the specified test conditions.</i>	<i>Coefficient of variance: 0.48 – 2.40 %</i>	<i>Detection limit 0.2 mg/l sample</i>	 (2007)																

Photometer (wavelength 334, 340, or 360 nm) to detect formation of NADH	<i>food and feedstuffs</i>	<i>Linearity is given in the range 0.2 mg formic acid/l sample solution to 200 mg formic acid/l sample</i>	<i>None (enzyme specific for formic acid)</i>	16	Fortification level [mg/L]	Number of measurements	Mean concentration [mg/L]	Rel SD [%]	<i>Detection limit 0.2 mg/l sample</i>	[REDACTED] (2007)
					0	6	9.96	2.5		
					10	4	18.77	11		
					50	5	62.88	0.9		

No data submitted for sediments:

Based on the physico-chemical properties as well as the environmental fate of formic acid, the compartment sediment is of no concern for this substance.

Formic acid

- is readily biodegradable,
- is completely miscible with water,
- has a low potential for adsorption (log Kow -1.9 to -2.3; log Koc < 1.25)
- will predominantly distribute into the compartment water (93.5%), while a negligible fraction will be associated with the sediment (5.9E-05%) according to the Mackay Level I model (BPD ID IIA4.1.1.3_01).

It can be concluded that formic acid will be rapidly removed from the environment due to biodegradability. As it is completely miscible with water and has a low adsorption potential formic acid will not distribute into the compartment sediment. This is supported by the Mackay level I model result, which shows that formic acid will predominantly distribute into the compartment water. Therefore, no analytical method for the detection of formic acid is provided.

2 EFFECTS AGAINST TARGET ORGANISMS

2.1 FUNCTION AND FIELD OF USE ENVISAGED

FUNCTION

Main Group 1 : DISINFECTANTS

- *PT2* "Disinfectants and algacides not intended for direct application to humans or animals"
- *PT3* "Veterinary hygiene"
- *PT4* "Food and feed area"
- *PT5* "Drinking water"

Main Group 2 : PRESERVATIVES

- *PT6* "Preservatives for products during storage"

With Bactericidal, yeasticidal & fungicidal activity.

To control the spread of microorganisms which may be harmful to human health.

FIELD OF USE ENVISAGED

The Formic Acid-based Biocidal products are wide-spread and have the following aims :

- *PT2* : Disinfection of industrial and institutional premises and machinery, for Cleaning-In-Place procedures, bathroom surfaces, toilets and sanitary ware in the domestic and institutional environment i.e. walls, toilets and other hard surfaces.
Products are applied by non-professionals by pouring and wiping; professionals apply the diluted concentrate as cleaning-in-place. Products to be used by professionals are concentrated formulations and by general public RTU formulations.
- *PT3* : Disinfection of areas in which animals are housed, kept and transported.
Products to be used for animal house disinfection (by fogging), for disinfection of footwear (by dipping) and for animal's feet and animal transport vehicles disinfection
Products to be used by professionals (i.e. professional contractors or experienced farm workers)
- *PT4* : Disinfection of working areas and production surfaces including food preparation and consumption areas.
Products to be used for hard surface disinfection (by trigger spraying) and for Cleaning-In-Place procedures. Products to be used by professionals.
- *PT5* : Disinfection of drinking water for animals
Products to be used by professionals
- *PT6* : Preservation of industrial, consumer, household and institutional products, washing and cleaning fluids and other detergents, and formulation of detergent end product.

2.2 INTENDED USES

Summary table of intended use(s)	
Product Type	PT3 Veterinary hygiene
Product description	Formic Acid-based Biocidal products are recommended for the disinfection of hard surfaces in which animals are housed and kept and vehicles (including tyres) used in the transport of animals, footwear and machinery connected with livestock farming.
Target organisms (including development stage)	Bacteria Yeasts
Description of use(s)	Formic Acid-based Biocidal products are applied directly to the surface to be treated. Disinfection of animal housings : Formic Acid-based Biocidal products are applied by the use of a fogging system. Means of transport, footwear and machinery : Formic Acid-based Biocidal products may be used by dipping, e. g. by the use of troughs containing disinfectant through which vehicles, the footwear of personnel or animals may pass.
Mode of action	The biocidal activity of Formic Acid, i.e. acidulant action and corrosion which causes enzyme denaturation and inhibition, cellular structure disruption, and impairment of cellular metabolic pathways. This mode of action is considered to depend on the low pH-value. Secondly, formic acid does inhibit cytochrome C oxidase and thus impairs cellular energy supply. Organisms and tissues with a high energy demand are specifically susceptible : Acidulant: acidification of cytoplasm; Inhibitor for decarboxylases and haemin enzymes such as catalase; Organic acids in general may disrupt the proton-motive force, as well as inhibit substrate transport, energy-yielding processes and macromolecular synthesis. Acidulant action is responsible for formic acid being most effective at lower pH values (below 3.5), but enzyme inhibition and other modes also provide some antimicrobial action at higher pH values. Enzyme inhibition is less significant in the control of fungi; therefore, higher concentrations of formic acid are needed to control fungi. The activity of formic acid against some viruses is presumably explained by the action of acid in denaturing polypeptide chains.
Objects to be protected	Humans and animals The aim of the treatments is to control infectious diseases.
Concentration of product in the in-use formulation/product	Representative product used in efficacy tests : Protectol® FM 85 with 85% Formic Acid
Application rate(s)	The product Protectol® FM 85 (based on 85% FORMIC ACID) is bactericidal and yeasticidal at 5.88 % (⇔ 5% FORMIC ACID) in 30 min at +10°C on hard/non-porous surfaces with prior cleaning.
Frequency of application	Disinfection of animal housings : One application is made prior to introduction of animals.

Summary table of intended use(s)	
	Means of transport, footwear and machinery : Vehicles and relevant machinery are sprayed down with disinfectant after each period of work (daily occurrence). Vehicles will use a drive through each time that they visit critical areas. Footwear will be disinfected in a footbath each time that a critical area is visited.
Season/period for use (where relevant)	Not relevant
Field of use (indoors/outdoors)	Indoor
Category(ies) of user(s)	Industrial and professional users, depending on the respective product.
Instruction for use	<p>Fogging of animal houses Clean the animal house and let the surfaces dry. The fogging equipment is loaded manually with the disinfectant outdoors and diluted with the appropriate amount of clean water. Put the equipment inside, close the house well and start the fogging process automatically from the outside. No people or animals should be in the animal house during fogging. Fog the house for at least 2 h up to overnight. Start ventilation and air the house well for ca. 2 hours before restocking.</p> <p>Hoof disinfection Dilute the concentrated product to the desired in-use concentration with clean water in a suitable container or in the bath itself. Mix well. Add the diluted concentrated product to an appropriate bath or mat, where the animals can walk through. Renew the solution if the use solution is getting too dirty.</p> <p>Wheel/boot disinfection Clean and rinse surfaces prior to disinfection Decant or dispense the concentrated product and dilute with water. Fill the bath with the diluted concentrate and place at the entrances/ exits. Make sure the wheels / boots are kept at the required contact time before moving on. Footbaths must be kept indoors/under roof in order to prevent dilution by rain.</p>

2.3 SUMMARY ON EFFICACY

2.3.1 Efficacy

General overview

Formic acid-based products exert toxic effects on the target organisms.

Since the representative product ██████████, no efficacy studies have been performed and submitted using only the active substance. Therefore, to review efficacy data available for formic acid, please see information in **Part B** of this CAR.

2.3.2 Mode of action

Different modes of action are reasonably considered to contribute to the biocidal activity of Formic Acid, i.e. acidulant action and corrosion which causes enzyme denaturation and inhibition, cellular structure disruption, and impairment of cellular metabolic pathways.

This mode of action is considered to depend on the low pH-value. Secondly, formic acid does inhibit cytochrome C oxidase and thus impairs cellular energy supply. Organisms and tissues with a high energy demand are specifically susceptible.

1. Acidulant: acidification of cytoplasm;
2. Inhibitor for decarboxylases and haemin enzymes such as catalase;
3. Organic acids in general may disrupt the proton-motive force, as well as inhibit substrate transport, energy-yielding processes and macromolecular synthesis.

Acidulant action is responsible for formic acid being most effective at lower pH values (below 3.5), but enzyme inhibition and other modes also provide some antimicrobial action at higher pH values. Enzyme inhibition is less significant in the control of fungi; therefore, higher concentrations of formic acid are needed to control fungi. The activity of formic acid against some viruses is presumably explained by the action of acid in denaturing polypeptide chains.

2.3.3 Resistance

There is no adaptation to cope with acidic pH values or denaturated proteins, nor is there a mechanism known to exist that a sub-lethal energy supply, due to an incomplete cytochrome C oxidase inhibition, would lead to undesired side-effects or resistance against this inhibitor.

No incidence of resistance to formic acid has been recorded until now.

2.4 CONCLUSION ON EFFICACY


Since the representative product is [REDACTED], no efficacy studies have been performed and submitted using only the active substance. Therefore, to review efficacy data available for formic acid and to read a conclusion on efficacy, please see information in **Part B** of this CAR.

3 ASSESSMENT OF EFFECTS ON HUMAN HEALTH


3.1 TOXICOKINETICS

In aqueous solution and at neutral pH, formic acid and water-soluble formate salts dissociate and are present as the formate anion in solution. The behaviour of chemical dissociation in water has particularly been investigated with potassium diformate [CAS No. 20642-05-1], which served as test compound in several toxicity studies (DocIIIA6.2_01; FA_BPR_Ann_II_8_8_01: ██████████, 1997). Based on the physico-chemical properties, it is justified to include kinetic and metabolism studies conducted with water-soluble formate salts.

Potassium formate is expected to form the following equilibriums in aqueous solutions:



Mapping the pH as function of dilution and titer curve allowed to estimate the buffer effect of the diformate system



and to calculate the concentration profile of diformate, formic acid and formate as function of concentration in aqueous solutions. The same procedure was applied to formic acid.

The calculations indicate that in aqueous solutions

- i) at pH <4 and at concentrations >0.1% the equilibrium in equation 1 is in favor of potassium diformate.
- ii) at pH of 4 to 5, and at dilution down to 0.001%, most of the formic acid content is released from potassium formate.
- iii) further dilution and increase of pH above 5, the concentrations of formic acid and diformate decrease rapidly, leaving only formate left at pH 7 and above. No diformate exists above pH 7.




Fig. 3.1 Formic acid – Formate ratio vs. pH by dilution

Formic acid occurs naturally in animals and most plants. Formic acid is an inherent ingredient in human food. The content reported for some common foods and beverages: fruits 20 to 40 mg/kg; honey 20 to 2000 mg/kg; wines 1 to 340 mg/kg; roasted coffee 30 to 40 mg/kg; cheese 20 to 200 mg/kg. Formic acid was added intentionally to some foods such as ice cream, soft drinks and fruit drinks as a flavor adjunct. The dietary consumption in adults was estimated to range between 0.4 and 1.2 mg/kg per day (DocIIIA6.2_09; FA_BPR_Ann_II_8_8_08; Boeniger, 1987). JECFA/IPCS (2003, originally published by WHO, 1997; BPD ID A6.15.4_01b, FA_BPR_Ann_II_8_16_1_01) stated that endogenous formate is generally present in human blood at levels of 0.07 – 0.4 mM (3.2 – 18.4 mg/l). Further, formic acid is required for the biosynthesis of purines and pyrimidines in the intermediary metabolism.

Formic acid is considered to be available by all potential exposure routes.

The toxicokinetic properties and the metabolism of formic acid have been investigated after oral, inhalation, intravenous, or intraperitoneal administration, in different species: rat, mouse, dog, monkey, pig, and humans. None of the studies were performed according to regulatory guidelines (some are pre-guideline). Nevertheless, the studies were conducted in accordance with generally accepted scientific principles, techniques and methods, and hence are acceptable for assessment. In addition, PBPK models were developed based on data collected after intravenous and inhalation exposure.

Justification for read-across:

The repeated dose toxicity via the oral route of formic acid is assessed with its non-corrosive salts, sodium formate and potassium diformate, in order to achieve sufficiently high dose levels. Neurotoxicity is assessed with methanol. A read across approach is provided in accordance with Article 6(3) of the EU No. 528/2012 (BPR) following point 1.5(2) under Annex IV: "common precursors and/or the likelihood of common breakdown products via physical and biological processes, which result in structurally similar chemicals and indicates the

presence of dangerous properties". The full read-across justification, which was performed following the Read-Across Assessment Framework developed by ECHA, can be found in Appendix VII.

The read-across justification concludes that the hypothesis that systemic toxicity of formic acid can be established by its salts, sodium formate and potassium diformate, and a closely related substance methanol, as these chemicals have a common breakdown product *in vivo*, is supported by the available information on physicochemical properties and its toxicokinetics.

When making use of this read-across, reference values will be derived for formate and expressed as mg formate/kg bw/d. At physiological pH 7, formic acid and potassium diformate are both exclusively present as formate anion. Therefore, inside the body, the major form present after exposure to either formic acid or potassium diformate is formate. (pKa of FA is 3.70 at 20°C). In water, there is an equilibrium between formic acid and the dissociated acid ($\text{HCOOH} \leftrightarrow \text{H}^+ + \text{COOH}^-$). Once its corrosive properties have been exerted, only formate is released/available. Therefore, for those HH endpoints where read-across is relevant, the endpoint will be expressed as formate. A conversion is not needed as the difference between formic acid and formate is limited to 1 H⁺ (MW of formate is 1 less than formic acid).

Toxicokinetics

The toxicokinetic properties of formic acid and sodium formate were studied in **human volunteers** following **oral** ingestion (DocIIIA6.2_07; FA_BPR_Ann_II_8_8_06: Malorny, 1969b). Formate and formic acid were both rapidly absorbed and reached peak plasma levels within 10 to 30 min after ingestion. Resorption of the unprotonated acid started already in the stomach; sodium formate was converted to the unprotonated acid under the pH conditions of the stomach. After ingestion of a single dose of 1000 mg formic acid (12.5 mg/kg bw), the increase in the plasma level of formate was barely distinguishable against a baseline (about 4 mg/l), while a transient 3- to 4-fold increase in formate (20 mg/l plasma) was noted after ingestion of 2000 mg formic acid (26.7 mg/kg bw). Formate was eliminated from the plasma with a half-life time $t_{1/2} = 45$ min. The background urinary formate excretion in humans was approx. 13 mg/24 hours. The average urinary excretion accounted for approx. 2 - 4 % of the administered dose, but was very variable among the individuals. The major part of ~65 - >80 % was excreted within the first 6 hours after ingestion and returned to normal levels at 12 hours after dosing. The blood pH remained unchanged following single formate or formic acid doses that were equivalent to 3000 mg formic acid. Urine volume and pH were increased as long as formate was excreted via urine.

In a human pharmacokinetic study (Hanzlik et al. (2005); FA_BPR_Ann_II_8_8_10) females (n=14) ingested 3900 mg calcium formate (equivalent to 2700 mg formate). The endogenous formate level was approx. 0.024 ± 0.008 mM in this study. Absorption was fast and the mean maximal serum level of 0.50 mM was seen at 60 minutes after dosing.

The toxicokinetic properties from plasma formate concentrations were studied in the **pig** following **oral** ingestion of potassium diformate by ██████████, 1998 (DocIIIA6.2-10; FA_BPR_Ann_II_8_8_09). Potassium diformate dissociated to formate *in vivo* as expected when it was fed with the diet to pigs. Absorption was rapid; the mean half maximal plasma level of approx. 200 mg formate/l plasma was reached in less than 2 hours, and the mean maximal plasma level $C_{\text{max}} = 386.4$ mg/l was seen 4 to 5 hours after feeding had been started. The values were derived from those 4 pigs which consumed at least 80% of the feed within 40 minutes after it had been offered. Formate was rapidly and completely eliminated. The mean biological half-life was calculated to be $t_{1/2} = 2.73$ hours, i.e. about 25 % of the amount in blood will be removed per hour (first-order elimination, $k_{\text{el}} = 0.25 \text{ h}^{-1}$). In 3 pigs, control plasma levels (mean: 1.9 mg/l) were reached within 12 hours; after 24 hours all pigs had

normal plasma levels. There was no indication of an accumulation of formate. Only 13.5 % of the high oral dose was found systemically bioavailable. This seemed to be due mainly to the metabolic activity of the liver (hepatic "first-pass effect") and secondly to the urinary elimination. Furthermore, it was assumed that not 100 % of the dose was resorbed, while part of it might also have been subjected to degradation by the gut microflora. A quantitative gastro-intestinal absorption rate could not be derived from this study.

A PBPK model (multicompartment dynamic system) developed by Bouchard et al., 2001 (DocIIIA6.2-03; FA_BPR_Ann_II_8_8_02), described the toxicokinetics of methanol, formaldehyde and formic acid in rats, monkeys, and humans for up to 48 h following **inhalation** exposure to methanol. The volume of distribution of formate of 6.4 to 4.2 l/kg bw suggested that a significant proportion of formate distributes in the tissue, but more likely undergoes rapid metabolism and excretion, thus leading to an apparently high distribution volume. The metabolism rate constant ratio $k_{\text{form}}/k_{\text{fald}}$ was twice as high in rats as in monkeys (0.53 vs. 0.26). Thus, in monkeys and plausibly in humans, a much larger fraction of formaldehyde is rapidly converted to unobserved forms rather than metabolized to formic acid and further to CO₂. For humans, the simulations showed that after continuous inhalation of 260 mg methanol/m³ (200 ppm) for 5 days, methanol-related blood and urinary formate levels (0.16 mg/L and 1.5 mg/L, respectively) remained far below reported baseline levels in unexposed subjects (4.9-10.3 and 6.3-13 mg/L, respectively). Furthermore, the model predicted that an 8-hour inhalation of 650 - 2600 mg/m³ (500 to 2000 ppm) methanol would be required to reach endogenous baseline values of formate.

Additional information on distribution is provided in section 3.6.1 on sub-chronic oral toxicity: systemic bioavailability data were provided in the study by [REDACTED] (1998; BPD ID A6.4.1_01, FA_BPR_Ann_II_8_9_2_01) notably it was reported that formate plasma levels of approx. 90 to 160 mg /L were regularly found in rats after oral exposure to potassium diformate. In section 3.14 on Further human data, for a case report on suicidal ingestion of Formic Acid, data on post-mortem formate concentrations are available.

Crossing of barriers as blood/brain, blood/testes, blood/placenta, and exposure via the breastmilk: It may be deduced from the physico-chemical properties of formic acid that the possibility of formate to cross the mentioned barriers is low. The substance is highly soluble in water and the logKow is around -2.0. The pKa is 3.70 at 20°C, and therefore formic acid (and the related salt potassium diformate) is almost exclusively present in the ionised form at physiological pH (DocIIIA6.2-01; FA_BPR_Ann_II_8_8_01). It is known that only the unionised form is likely to cross biological membranes, and that substances with a logP of 2-4 would likely cross membranes. The physico-chemical properties of formic acid differ largely, hence it is unlikely that formate would cross biological membranes. This does not preclude the uptake by means of active transport systems. Penetration into (and through) membranes may occur in minor quantities because the small size of the formate molecule. Transfer into breast milk may be given due to the high solubility in water. In this context it should also be mentioned that endogenous formic acid is produced in the intermediary metabolism in humans, and that the C1-fragment is required in the biosynthesis of amino acids and nucleic acids (DocIIIA6.2-09; FA_BPR_Ann_II_8_8_08), i.e. there is a need in the developing fetus. Excess blood formate is rapidly metabolised to background levels in humans, i.e. formate does not accumulate. Finally, there were no adverse effects noted in the testes, the brain, or the development of offspring, in any of the numerous studies requiring repeated dosing. This includes all subchronic and chronic repeated dose studies, carcinogenicity studies, multigeneration reproduction and teratogenicity studies, conducted in several species (rat, mouse, rabbit, pig) with either sodium formate or potassium diformate. Neurotoxicity is known to occur in humans only in the optical nerve following severe methanol intoxication leading to very high blood formate levels over an extended period of time (DocIIIA6.9; FA_BPR_Ann_II_8_13_2_0). Thus, though formate crossing of the blood/brain, blood/testes, blood/placenta barriers, and the exposure via the breast milk cannot be fully excluded, no adverse effects were seen in the parental animals and their progeny of several species following high-level long-term dosing, or dosing during reproduction and development, of either sodium formate or potassium diformate.

Metabolism

The metabolism of formic acid in animals has been extensively documented. Formic acid is an intermediate in normal metabolism. It takes part in the metabolism of one-carbon compounds and its carbon may appear in methyl groups undergoing transmethylation. The metabolic oxidation of formate to CO₂ involves tetrahydrofolate (THF). Formyl-THF synthetase catalyzes the binding of formate to THF to yield 10-formyl-THF. The latter liberates CO₂, and the folate moiety is reduced to THF by Formyl THF dehydrogenase.




Fig. 3.2 Oxidation of formate to CO₂

The oxidation rate of formate to CO₂ depends on the hepatic folate pathway, i.e. the levels of folate coenzymes and folate-dependent enzymes. These levels are higher in rodents than in primates, and consequently the rate of formate oxidation to CO₂ is also higher in rodents (DocIIIA6.2_04; FA_BPR_Ann_II_8_8_03: NTP, 2004). In monkeys, the maximum elimination rate of formate is reported to be about 34 mg/kg bw/h, whereas in rats it was about 73 mg/kg bw/h (BPD ID A6.2_12; ; FA_BPR_Ann_II_8_8_13: Kavet & Nauss, 1990). The formate plasma elimination half-life in various species following intravenous infusion (see table 3.1-1) was discussed in a review by Malorny, 1969a (DocIIIA6.2_06; FA_BPR_Ann_II_8_8_05). There is a clear species difference in the extent of formic acid metabolism and elimination rate which is consequently dose-dependent. As humans and primates have reduced capacity for formate oxidation compared with rodents and dogs, humans and primates are more susceptible to formate intoxication.

Formic acid was rapidly oxidised to CO₂ and water by the liver in human volunteers, while a minor part of 2 to 4 % was excreted unchanged into the urine within 24 hours (DocIIIA6.2_07; FA_BPR_Ann_II_8_8_06: Malorny, 1969b). Based on the first-order elimination kinetics (see Table 3.1-1), it is evident that after exposure to one single dose of formic acid or formate salt that was systemically bioavailable, normal blood levels will be reached within 4 to 5 hours post-application in humans.

In the recent single dose human study (FA_BPR_Ann_II_8_8_10: Hanzlik et al, 2005), a mono-exponential decline of serum concentrations with an average half-life of 59 +/- 7 minutes was seen, and baseline levels were reached within 240 minutes after dosing (see figure and legend below).




Fig. 3.3 Plasma formate concentration versus time for 14 adult female human subjects following administration of placebo (A) or calcium diformate (B).

This finding is in good correlation with the earlier reported human half-life of 45 minutes (Malorny (1969b); BPD ID A6.2_07; FA_BPR_Ann_II_8_8_06).

The disappearance of formate from blood is shown in Table 3.1-1.

Table 3.1-1: First-order elimination half-lives of formate in blood plasma in various species

Species	t _{1/2} (min)	Source
Rat	12	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Guinea pig	22	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Rabbit	32	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Monkey	30 - 50	BPD ID A6.2_11; FA_BPR_Ann_II_8_8_12: Clay et al., 1975
Human	45	BPD ID A6.2_07; FA_BPR_Ann_II_8_8_06: Malorny, 1969b
Human	59	FA_BPR_Ann_II_8_8_10: Hanzlik et al., 2005
Cat	67	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Dog	77	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Pig	87	BPD ID A6.2_08; FA_BPR_Ann_II_8_8_07: Makar et al., 1990
Pig	164	BPD ID A6.2_10; FA_BPR_Ann_II_8_8_09: ██████████, 1998

The pig shows the most limited metabolic capacities of reported test species (mouse >rat >monkey >human >pig). Formate metabolism in the pig in comparison to the rat was studied by Makar et al. (1990) (DocIIIA6.2_08; FA_BPR_Ann_II_8_8_07). 14C-radiolabeled formate was applied i.p. and determined in blood only, not including urine levels and exhaled CO₂. No complete mass balance was provided. The species-specific metabolic capacities of the liver to convert formate were also analysed. The results indicated that the pig has very low levels of folates and low levels of key enzyme in the folate pathway as compared to rodents, monkey and humans. The pig's ability to dispose of formate was found more limited and much slower than that observed in rats or monkeys. It was suggested that the pig may be a suitable model for studying formate metabolism, because accumulation of formate and susceptibility to its toxic effects must be considered.

In humans, formate bioaccumulation is less likely to occur, based on the results of the early and the more recent single dose human studies (e.g. FA_BPR_Ann_II_8_8_10: Hanzlik et al, 2005), and based on the results of a recent repeat dose human study (Altaweel et al., 2009: FA_BPR_Ann_II_8_8_11). No formate accumulation was noted in a 14-day human study (12 females) who ingested 3900 mg calcium formate/day. The baseline serum formate level was 0.539 ± 0.06 mM in this study, maximal serum levels were approx. 0.8 mM (see below; figure and legend from Altaweel et al. (2009); FA_BPR_Ann_II_8_8_11).




Fig. 3.4 Serum formate concentrations determined prior to and throughout the 14-day study. The concentrations observed at screening (S) do not vary significantly, either for time groups or for individual subjects, over the course of the study. Observations 1a, 2a, 3a, 8a, and 15 were made prior to the first dose of the day indicated, while observations 1b, 2b, 3b, and 8b were made 40–60 min after ingesting the second 1,300 mg dose of calcium formate

on the day indicated. The sample times "a" and "b" correspond, respectively, to the times of trough and peak levels of serum formate following single doses (see Ref. 36). The data show no accumulation of serum formate with repeated administration of 1,300 mg calcium formate three times per day over 14 days.

Data on maximum blood levels of formate reported after single or repeated dosing of formic acid or formate salt are summarised in Table 3.1-2.

Massive serum formate levels are seen in primates (humans, monkeys) following methanol intoxication (see below). Higher formate serum levels are achievable following oral ingestion, compared to inhalation or dermal absorption. Massive serum formate levels are seen in primates (humans, monkeys) following methanol intoxication, whereas levels remain low in rats (not listed) unless the formic acid oxidase is inhibited by N2O-treatment. Under these conditions, formate levels are comparable to those seen in primates, i.e. the metabolic capacity of the rat was lowered to that of the primates, and under these conditions, the rat is also susceptible to toxic optical neurotoxicity.

Table 3.1-2: Maximum formate blood levels either after dosage of formic acid or formate salt or following methanol poisoning (see also Table 3.1-3)

Species	Substance	Route	Dose [mg/kg bw]	Peak blood level [mg/l]	Reference
Dog	Formic acid or Na formate	i.v. (~10 min)	~54 (as formic acid)	~200	BPD ID A6.2_06; FA_BPR_Ann_II_8_8_05: Malorny, 1969a
Pig	Na formate [CAS No. 141-53-7]	i.p.	~350 (as formic acid)	~470	BPD ID A6.2_08; FA_BPR_Ann_II_8_8_07: Makar et al., 1990
Pig	Potassium diformate [CAS No. 20642-05-1]	oral feed	~700 (as formic acid)	~400	BPD ID A6.2_10; FA_BPR_Ann_II_8_8_09: ██████████, 1998
Human (single cases)	Formic acid or Na formate	oral	~13	4 - 5 (baseline)	BPD ID A6.2_07; FA_BPR_Ann_II_8_8_06: Malorny, 1969b
		oral	~27	20	
		oral	~40	85	
Human (n=14) Single dose	Calcium formate	oral	2700 mg (as formate)	0.50 mM (mean)	FA_BPR_Ann_II_8_8_10: Hanzlik et al., 2005
Human (n=12) 14-day repeated dose	Calcium formate	oral	2700 mg/day (as formate)	0.572 mM (mean)	FA_BPR_Ann_II_8_8_11: Altaweel et al., 2009
Rat, N ₂ O-pre-treated	Methanol intoxication	oral	4000 mg/kg (methanol)	16 mM	Cited in

					BPD ID A6.10_01; FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000
Monkeys	Methanol intoxication	oral	Dose not stated (methanol)	11.4 mM	Cited in BPD ID A6.10_01; FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000
Humans	Methanol intoxication	oral	Dose not stated (methanol)	19.3 mM	Cited in BPD ID A6.10_01; FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000

A great deal of knowledge about the metabolism of formic acid has been extensively documented within the investigations into the mechanism of methanol intoxication. Formic acid is one of the main metabolites of methanol. The absorption, distribution and elimination of methanol and formate have successfully been modeled after inhalation exposure to methanol in various species including humans. The model predictions were in good agreement with experimental data in various species, i.e. rat, monkey, and human data, suggesting that the values of the pharmacokinetic constants used in the model are close to real values (DOCIIIA6.2_03; FA_BPR_Ann_II_8_8_02: Bouchard et al., 2001).

Formate has to be considered as the causative agent for optical neural damage in methanol-intoxicated humans and animals (DOCIIIA6.2_05; FA_BPR_Ann_II_8_8_04: Martin-Amat et al., 1978; DocIIIA6.10_01; FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000). The blood levels of formate that correlated with the emergence of pathological changes were very high: In a review by Eells et al. (2000) the following values after accidental and experimental methanol intoxication were summarised (see fig 3.5 representing Table 2 from Eells et al. (2000)):

TABLE 2. Blood formate, pH and bicarbonate concentrations in methanol-intoxicated rats, monkeys and humans.

Species	Blood Formate (mM)	Blood Bicarbonate (mEq/L)	Blood pH
N ₂ O - Treated Rats ^a	16.1 ± 0.7	7.7 ± 1.2	6.91 ± 0.06
Monkeys ^b	11.4 ± 1.2	6.5 ± 0.5	7.19 ± 0.02
Humans ^{c,d}	19.3 ± 4.4	3.2 ± 0.4	6.93 ± 0.02

Note: Methanol-intoxicated rats were exposed to a mixture of N₂O/O₂ (1:1) for 4 hours prior to methanol administration (4 g/kg at zero time followed by 2g/kg at 12-hour intervals) and exposure to the gas mixture was continued throughout the experiment. Blood formate concentrations and blood gas measurements were determined 60 hours after the initial dose of methanol. Each value represents the mean ± SE for 6 rats. Rodent data was compiled from studies by Eells *et al.*, (1996)^a. The monkey data was compiled from studies by Martin-Amat *et al.*, (1977)^b and the human data was compiled from studies conducted by McMartin *et al.*, (1980)^c and Eells *et al.*, (1991)^d.

Fig. 3.5 Eells et al. (2000) Table 2

In four monkeys (Rhesus, *Maccaca mulatta*) receiving ~142 mg/kg bw/h of Na formate by i.v. infusion, the (steady-state) blood levels of formate amounted to 540, 950, 1350, and 1530 mg/l after 12, 20, 30 and 34 hours, respectively (DocIIIA6.2_05; FA_BPR_Ann_II_8_8_04: Martin-Amat et al., 1978). After 10 hours, all animals accumulated maximum formate in blood between 10 and 30 mEq/L (460 – 1380 mg/l). Under this extreme dosing regimen, the elimination half-lives had increased considerably up to about 5 hours, evidently due to metabolic overload and saturation [compare the dose of 142 mg/kg bw/h with the maximum metabolic capacity of 34 mg/kg bw/h, see above].

Critical blood concentrations of 8 – 15 mM formate (= 360 – 680 mg/l) maintained over 30 – 40 hours were considered potentially detrimental, producing experimental ocular toxicity in monkeys (DocIIIA6.2_05; FA_BPR_Ann_II_8_8_04: Martin-Amat et al., 1978) and were associated with visual toxicity in acute cases of human methanol intoxication (DocIIIA6.10_01; FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000).

Inhalation exposure to formic acid is supposed to be limited due to the warning of its pungent smell and its respiratory irritation unless through accidental events. 60 mg/m³ is considered to be the 13-weeks NOAEC for histological changes in the nasal region of rats and mice [see 13-week studies on rats and mice, DOCIIIA6.4.3_01/ FA_BPR_Ann_II_8_9_2_03 and DOCIIIA6.4.3_02/ FA_BPR_Ann_II_8_9_2_04, and section 3.6.4]. As for solid formate, inhalable quantities of solid formate salts are limited.

Assuming 100% absorption, a human body weight of 60 kg, and a high respiration volume of 1.25 m³/h under working conditions (BPD ID A6.12.8_01; FA_BPR_Ann_II_8_12_8_01: NIOSH, 1990), this concentration would correspond to a systemic dose of 610 mg/8 h or ~10.2 mg/kg bw/d or ~1.3 mg/kg bw/h.

Compared with the maximum conversion rate of formate to CO₂ in primates (BPD ID A6.2_12; FA_BPR_Ann_II_8_8_13: Kavet & Nauss, 1990), such an exposure level would not result in accumulation of formate.

At the maximum occupational exposure level of 5 ppm (9.5 mg/m³), the systemic dose would be only 1.6 mg/kg bw/d or 0.2 mg/kg bw/h under these assumptions (BPD ID BPD ID A6.12.8_01; FA_BPR_Ann_II_8_12_8_01: NIOSH, 1990).

Table 3.1-3 Main results from key and supporting study summaries

Summary table of toxicokinetic studies

Method Guideline, GLP status, Reliability	Species, Strain, Sex, No/Group	Test substance, Dose levels, Duration of exposure	Results	Remarks (e.g. major deviations)	Reference
In vitro / Physico-chemical studies on the behaviour of the TS in	n.a.	Formic acid [CAS No. 64-18-6] Potassium diformate [CAS No. 20642-05-1]	At physiological pH 7, formic acid and potassium diformate are both exclusively present as formate anion		██████████, 1997 BPD ID A6.2_01 FA_BPR_Ann_II_8_8_01

aqueous solutions. No guideline available		Route: not applicable Test procedure: Titration, calculations			
In vivo / No data, pharmacological standards	Dog, not specified, 6/group	Formic acid [CAS No. 64-18-6] and Na formate [CAS No. 141-53-7] Route: i.v. (~10 min) Dose: ~54 mg/kg bw Sampling intervals : 0, 1, 2, 4 hours after dosing: blood pH, formate blood levels	<u>Elimination:</u> from blood $t_{1/2} = 77$ min $k_{el} = 0.54$ h ⁻¹ <u>Blood levels:</u> Max. ~200 mg/l, Return to normal after 4 h <u>Baseline blood level:</u> ~7 – 12 mg/l (but high variance) <u>Blood pH:</u> transient acidosis, severe after formic acid and slight after Na formate, Return to normal after 3 to 4 h		Malorny, 1969a BPD ID A6.2_06 FA_BPR_Ann_II_8_8_05
In vivo / No data, pharmacological standards	Humans m + f 12, 7 and 2-3 per group	Formic acid [CAS No. 64-18-6] and Na formate [CAS No. 141-53-7] Route: oral Formic acid: 0.4% aqueous solution Na formate: in food Single dose Formic acid: 2000mg Na formate: 1.48, 2.96, 4.44 g (equivalent to 1000, 2000, and 3000 mg formic acid = ~13, 27, and	<u>Absorption:</u> rapid, maximum in blood after 10 – 30min <u>Bioavailability:</u> at 13 mg/kg bw in blood barely measureable, at ≥27 mg/kg max. 3-4fold increase in blood. Baseline blood level: ~3 – 4 mg/l (2 subjects) and 18 mg/l (1 subject) Max. blood level: 20 - 85 mg/l at 2000 mg <u>Elimination:</u> from blood $t_{1/2} = 45$ min =>		Malorny, 1969b BPD ID A6.2_07 FA_BPR_Ann_II_8_8_06

		40 mg/kg bw) Sampling: kinetics plasma levels: blood after 5, 120 min; urine after 15 min to 6 h urinary excretion: before, 0-6, 6-12, 12-24 hrs after ingestion blood pH: before, at 15, 30, 45, 60, 75, 90 min after ingestion	$k_{el} = 0.92 \text{ h}^{-1}$ Clinical signs: transient gastric irritation immediately after the ingestion of 2 g formic acid as 0.4% aqueous solution.		
In vivo / no data	Pig (crossbred) n=6 sex not reported <i>Control animal:</i> Rat Sprague-Dawley male, n=5	^{14}C -Na formate [CAS No. 141-53-7] Route: i.p. Dose: 500 mg/kg bw. (~350 mg formic acid/kg) Blood kinetics and liver folate metabolism (comparison among various species) Sampling intervals: 90, 180, 240, 300 min after dosing	<u>Elimination:</u> from blood $t_{1/2} = 87 \text{ min}$ $k_{el} = 0.48 \text{ h}^{-1}$ Max. blood level: ~470 mg/L The pig shows the most limited metabolic capacities of reported test species (mouse >>rat >monkey >human >pig).		Makar et al., 1990 BPD ID A6.2_08 FA_BPR_Ann_II_8_8_07
In vivo / No data, pharmacol. standards	Pig Crossbred (50% Duroc, 25% Yorkshire, 25% Danish Landrace)	Potassium diformate [CAS No. 20642-05-1] Route: 6% in oral feed	<u>Absorption:</u> rapid with maximum in blood after ~4 h <u>Bioavailability:</u>		██████████, 1998 BPD ID A6.2_10 FA_BPR_Ann_II_8_8_09

	n=4, female	<p>High single dose: 1000 mg/ kg bw (= ~700 mg formic acid/kg bw)</p> <p>Blood sampling: before, 0.5, 1, 1.5, 2, 3, 4, 5, 7, 12, 24 hours after at least 80% of the feed was eaten.</p> <p>Plasma formate concentrations used for calculation of the biological half-life ($t_{1/2}$), AUC, and C_{max}.</p> <p>Calculations according to a two compartment pharmacokinetic model, absorption and elimination processes considered to follow first-order kinetics.</p>	<p>Mean dose systemically bioavailable (AUC) = 2834.6 mg x h/l = ~13.5 % of the mean dose applied</p> <p>Baseline blood level: ~1.9 mg/l</p> <p>Max. blood level, C_{max} = 386 mg/l</p> <p><u>Elimination:</u> from blood $t_{1/2}$ = 2.73 h k_{el} = 0.25 h⁻¹</p> <p>Plasma formate concentrations returned to baseline after ~12 h p.a.</p>		
In vivo / No data, pharmacol. standards	Human subjects , females, n=14	<p>Calcium formate [CAS No. 544-17-2]</p> <p>Route: oral</p> <p>Single oral dose, 3900 mg (i.e. 2700 mg formate),</p>	<p>Endogenous formate level 0.024 ± 0.008 mM</p> <p><u>Absorption:</u> maximal serum level (mean: 0.50 mM) @ 60 min after dosing.</p> <p><u>Elimination:</u> mono-exponential</p>		Hanzlik et al., 2005 FA_BPR_Ann_II_8_8_10

		split into 6 doses of 650 mg each	decline of serum concentrations, average half-life 59 +/- 7 min. Baseline levels within 240 minutes post dosing		
In vivo / No data, pharmacol. standards	Human subjects , females, n=12	Calcium formate [CAS No. 544-17-2] Route: oral 14-days study, 3900 mg/day (i.e. 2700 mg formate/day), split into 3 daily doses of 1300 mg each	Mean basal serum formate level before study initiation: 0.539 ± 0.06 mM Formate levels only slightly increased at 40-60 min after dosing: up to 0.8 mM: No formate accumulation: serum formate level on day 15: 0.582 ± 0.091 mM; no significant difference between this value and the basal level before treatment (p=0.268).		Altaweel et al., 2009 FA_BPR_Ann_II_8_8_1_1

3.1.1 Short summary of the toxicokinetic information

Conclusions:

Formic acid is considered to be available by all potential routes of exposure. Inhalation may be the most relevant route during production and application.

For risk characterisation a value of 100% is used for oral absorption (rapid, but no quantitative data available) and for absorption via inhalation (no data available).

Dietary consumption of formic acid and its salts (estimated 0.4 and 1.2 mg/kg bw/day), inhalation as air contaminant as well as the endogenous turn-over maintain a baseline blood level of about 3 to 18 mg/l in humans; in a more recent study, it was found to be 0.539 mM.

Biotransformation of formate to CO₂ in primates is rapid: the first-order elimination half-life in human blood is approx. 45 min, corresponding to an elimination constant of about 0.9 h⁻¹, and the metabolic oxidation rate of formate is reported to be 34 mg/kg bw/h (0.75 mmol/kg bw/h). In human volunteers, a minor part was excreted unchanged into the urine within 24 hours. No accumulation is expected to occur, except at prolonged exposures above the critical capacity limit.

The steady-state blood concentration from a continuous dosage of 10 mg formic acid/kg bw/h that is systemically bioavailable will be of the order of 11 mg/l in humans, while a continuous

dose of 30 mg/ kg bw/h, at the borderline of metabolic saturation, is supposed to level off at 33 mg/l (see assumptions and estimation below).

Following inhalation, the experimental NOAEC of 61 mg/m³ in mice, corresponding to a 8-h dose of ~1.3 mg/kg bw/h would remain well below the metabolic capacity limit and result in a transient steady-state of approx. 17 mg/l in blood (in addition to the baseline level).

At the maximum occupational exposure level of 5 ppm (9.5 mg/m³), the systemic dose would be only 0.2 mg/kg bw/h, and the increment expected in blood would be indistinguishable from the endogenous fraction in blood.

Toxic effects are only expected, if the maximum metabolic oxidation rate becomes exhausted [>34 mg/kg bw/d], and thus critical formate blood concentrations are reached. These are reportedly in the range of 8 to 15 mM (= 360 – 680 mg/l).

The estimation below demonstrates that a bioavailable body burden of 1 mg formate/kg bw/h still fails to produce blood increases that are distinguishable from the baseline level, remaining at a factor of 300 to 600 below toxicologically relevant blood levels.

Estimation of a steady-state blood level:

The single-dose data can be used to estimate a blood concentration in equilibrium (steady state):

Assumption: Continuous oral uptake

Exemplary dose [D]: 1, 10, and 30 mg/(kg bw*h)

Gastro-intestinal bioavailability: 100 %

Elimination constant [k_{el}]: 0.9 h⁻¹ (from BPD ID A6.2_07/ FA_BPR_Ann_II_8_8_06)

Distribution volume [V_d]: 1 litre/kg bw*)

*) Note: Reportedly, the distribution volumes for formate range from about 4 to 6 litre/kg bw. But these values appear to be governed mainly by the rapid metabolism and excretion from the circulatory system. However, these processes are already comprised in the elimination constant k_{el} . Hence, a *high* distribution volume in the algorithm would be a bias resulting in underestimating the blood level. Therefore, adopting the conservative assumption of 1 litre/kg bw for the distribution volume of formate appears to be the more appropriate approach.

The steady-state concentration [C_{eq}] is described by the following equation:

$$C_{eq} = \frac{\text{Dose [mg/h]}}{\text{bw} * V_d * k_{el}} \text{ mg/l}$$

Table 3.1-4 Predicted steady-state concentration in blood C_{eq} during continuous dosage of Na-formate above baseline level

	Elimination constant k_{el} [h ⁻¹] / distribution volume V_d [L/kg]	Predicted steady-state concentration in blood C_{eq} during continuous dosage of Na-formate above baseline level (excluding baseline) (assumed absorption rate 100 %)	Baseline level in blood [mg/l]	Toxicologically relevant level [mg/l]
		Dose [mg/(kg*h)]		

		1	10	30		
Pig	0.25 / 1	4	40	120	~2 ¹⁾	No data
Human	0.90 / 1	1.1	11	33	3 – 18 ²⁾	>360 ³⁾

¹⁾ from BPD ID A6.2_10/ FA_BPR_Ann_II_8_8_09

²⁾ from BPD ID A6.2_07/ FA_BPR_Ann_II_8_8_06

³⁾ BPD ID A6.10_01/ FA_BPR_Ann_II_8_13_5_01

Dermal absorption

Dermal absorption of formic acid has not been investigated. Due to the corrosive properties of formic acid, no dermal absorption study is requested. In a first tier of risk assessment, a worst case value for dermal absorption of 100% is used for external dermal exposure. Severe metabolic acidosis resulting from dermal contact with formic acid as described in several case reports (see section 3.3 and 3.14), demonstrated rapid dermal absorption through the acid-burned skin.

3.1.2 Values and conclusions used for the risk assessment

Value(s) used in the Risk Assessment – Oral absorption	
Value(s)	100%
Justification for the selected value(s)	Formic acid is rapidly absorbed after oral ingestion by humans (DocIIIA6.2_07; FA_BPR_Ann_II_8_8_06: Malorny, 1969b) Rapid absorption, but no quantitative data available ³

Value(s) used in the Risk Assessment – Dermal absorption	
Value(s)**	100%
Justification for the selected value(s)	Dermal absorption of formic acid has not been investigated. A dermal absorption of 100% is used for external dermal exposure because rapid dermal absorption was demonstrated following acid skin burns in several case reports.

** the dermal absorption value is applicable for the active substance and might not be usable in product authorization

³ Due to animal welfare reasons an oral absorption study was not provided for formic acid as corrosive substance. However, the available toxicokinetic data and data on absorption after accidental or suicidal oral ingestion of the substance by humans indicate rapid and almost quantitative absorption.

Generally, the smaller the molecule the more easily it may be taken up. With a molecular weight of 46.03 g/mol formic acid is very favorable for oral absorption.

Furthermore, formic acid is miscible with water at any ratio which also favors oral absorption since water-soluble substances will readily dissolve into gastrointestinal fluids. Additionally, molecules with a molecular weight lower than 200 may pass through aqueous pores or may be carried through the epithelial barrier by the bulk passage of water.

Together with the observed clinical signs after oral ingestion, it is highly probable that formic acid is orally absorbed to a high extent.

As worst case 100% absorption is assumed.

Value(s) used in the Risk Assessment – Inhalatory absorption	
Value(s)	100%
Justification for the selected value(s)	no data available (assumed 100% resorption)

Conclusion(s) used in the Risk Assessment – Distribution	
Conclusion	No data
Justification for the conclusion	no data available; assumed distribution in the aqueous compartment: seemingly a significant proportion of formate distributes in the tissue, but more likely undergoes rapid metabolism and excretion Assumptions presented are based on a PBPK model. The physico-chemical properties of formic acid suggest the likelihood of it crossing blood/brain, blood/testes, and blood/placenta barriers is low. Transfer into breast milk may occur due to high water solubility.

Conclusion(s) used in the Risk Assessment – Metabolism	
Conclusion	Rapid oxidation to CO ₂ and H ₂ O No toxicologically significant metabolites
Justification for the conclusion	maximum elimination rate of formate: Monkey: 34 mg/(kg bw*h) Rat: 73 mg/(kg bw*h)

Conclusion(s) used in Risk Assessment – Elimination									
Conclusion	Rapid elimination from blood plasma No potential for accumulation Rate and extent of excretion: human: 2 to 4%/24h unchanged into the urine, ~65 - >80% thereof excreted within the first 6h.								
Justification for the conclusion	Humans: elimination half-life (t _{1/2}) = 45 min corresponding to an elimination constant of about 0.9 h ⁻¹ Rapid biotransformation of formate to CO ₂ in primates Metabolic oxidation rate of formate 34 mg/kg bw/h (monkey). Human volunteers: minor part was excreted unchanged into the urine within 24 hours (see above). No accumulation is expected to occur, except at prolonged exposures above the critical capacity limit.								
	<table border="1"> <thead> <tr> <th>Species</th> <th>t_{1/2} (min)</th> </tr> </thead> <tbody> <tr> <td>Rat</td> <td>12</td> </tr> <tr> <td>Guinea pig</td> <td>22</td> </tr> <tr> <td>Rabbit</td> <td>32</td> </tr> </tbody> </table>	Species	t_{1/2} (min)	Rat	12	Guinea pig	22	Rabbit	32
Species	t_{1/2} (min)								
Rat	12								
Guinea pig	22								
Rabbit	32								

		Monkey	30 - 50	
		Human	45	
		Cat	67	
		Dog	77	
		Pig	87	
		Minipig	164	

3.2 ACUTE TOXICITY

The acute toxic action profile of formic acid is predominantly determined by its inherent irritating/corrosive properties. The toxicity values after oral uptake and inhalation in rats suggest formic acid to be acutely harmful. The clinical signs give no evidence of specific systemic adverse effects.

3.2.1 Acute oral toxicity

Summary table of animal studies on acute oral toxicity						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance Dose levels, Type of administration (gavage, in diet, other)	Signs of toxicity (nature, onset, duration, severity, reversibility)	Value LD50	Remarks (e.g. major deviations)	Reference
OECD 401 GLP: no Rel: 1	Rat Wistar m + f 5/sex/group	Formic acid purity 99% Lot/batch: no data 501, 631, 794, 1000 mg/kg bw gavage	Clinical signs: - observed 30 min after dosing: unkept fur, hunched posture, stagger, aggressiveness, dyspnoea, sedation and ataxia, lateral and abdominal position, convulsions, bloody noses, blood in urine. - later: hypothermia, pale limbs, body weight loss. - Symptoms subsided and were absent in all animals but one which showed symptoms until d14.	730 mg/kg bw (m +f) Males: 863 mg/kg bw Females: 618 mg/kg bw		BPD ID A6.1.1_01, FA_BPR_Ann_II_8_7_1_01: [REDACTED], 1985

Formic acid is of moderate toxicity via the oral route when tested in the rat. Oral LD₅₀ = 730 mg/kg bw.

After a single *oral* administration of formic acid in the rat (DocIIIA6.1.1-01, FA_BPR_Ann_II_8_7_1_01: ██████████, 1985), there was a clear dose-response relationship with respect to mortality and decrease in body-weight gain of survivors. The clinical symptoms and pathological organ lesions (hyperemia of the stomach and intestines, congestion in spleens, mottled livers and kidneys, discoloration of kidneys and pancreas) are largely nonspecific and can be explained primarily by the local corrosive character of formic acid, associated secondary systemic effects.

For human data: see section 3.14.

Several case reports report on fatal suicidal ingestion of formic acid (see section 3.14 for a detailed discussion). Due to the corrosivity of formic acid, local effects occur at all dose levels. The amount ingested and the concentration determine the grade and the location of the effects. Therefore, the observations range from moderate burns around the mouth to severe corrosion of the gastro-intestinal tract with destruction of the oesophagus, perforation of the stomach, and corrosion of the small intestine together with massive bleeding and systemic toxicity. Systemic toxicity was seen after ingestion **of 30 g formic acid** or more. Prognosis is poor after massive oral ingestion (>45 to 200 g formic acid); prognosis is moderate after moderate oral ingestion (approx. 30 to 45 g); lesions, but low mortality, are expected in most cases with low amounts ingested (<30g); persistent lesions due to tissue corrosion must be expected in cases with >10 g formic acid ingested. Tissue destruction of the gastrointestinal tract may result in fatal bleeding, septic shock, or stricture which may require surgical treatment. Reversibility of effects was often seen in cases with low amounts ingested (<10 g formic acid).

Important note:

Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

Value used in the Risk Assessment – Acute oral toxicity	
Value	LD ₅₀ 730 mg/kg bw ⁴
Justification for the selected value	BPD ID A6.1.1_01, FA_BPR_Ann_II_8_7_1_01: ██████████, 1985 Acute oral toxicity of formic acid has been assessed in a study according to OECD 401.

⁴ Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

3.2.2 Acute dermal toxicity

Summary table of animal studies on acute dermal toxicity					
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance, Vehicle, Dose levels, Surface area,	Value LD ₅₀	Remarks (e.g. major deviations)	Reference
OECD 402 GLP: yes Rel: 1	Rat Wistar m + f 5/sex/group	Sodium formate [CAS No. 141-53-7] purity 100% Lot/batch: 1292066 2000 mg/kg bw limit test Vehicle: 0.5% CMC 24 hours, semi-occlusive Surface area 40 cm ² (10% of body surface)	>2000 mg/kg bw No clinical signs, or local, or systemic effects observed. No mortality.	Other test substance: sodium formate	BPD ID A6.1.2_01, FA_BPR_Ann_II_8_7_3_01 ██████████ 2007

No acute *dermal* study has been conducted with formic acid itself because of its corrosive properties. After single dermal exposure of the sodium salt in the rat (DocIIIA6.2.1-01, FA_BPR_Ann_II_8_7_3_01: ██████████ 2007), no local irritation and systemic effects were observed. Dermal LD₅₀ of sodium formate >2000 mg/kg bw.

Human case reports on acute 'accidental' dermal (and inhalation) exposure are rather rare. Besides local effects, severe acid skin burns and respiratory tract irritation, patients suffered and recovered rapidly from metabolic acidosis (described in section 3.3 and 3.14).

Value used in the Risk Assessment – Acute dermal toxicity	
Value	No data available on formic acid Supportive data: Sodium formate: LD ₅₀ >2000 mg/kg bw

Justification for the selected value	<p>According to regulation (EU) 528/2012 Annex II 8.7 acute toxicity studies does not generally need to be conducted if the substance is classified as corrosive to the skin due to animal welfare reasons.</p> <p>Hence, the information on the acute dermal toxicity of the corresponding salt, sodium formate, is only supportive information, as no information on acute dermal toxicity is needed for this substance.</p> <p>BPD ID A6.1.2_01, FA_BPR_Ann_II_8_7_3_01: ██████████ 2007</p> <p>Acute dermal toxicity of sodium formate has been assessed in a study according to OECD 402.</p>
---	--

Data waiving	
Information requirement	Acute dermal toxicity of formic acid
Justification	Corrosive substance

3.2.3 Acute inhalation toxicity

Summary table of animal studies on acute inhalation toxicity					
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance, form (gas, vapour, dust, mist) and particle size (MMAD) Actual and nominal concentration, Type of administration (nose only / whole body/ head only)	Value LC50	Remarks (e.g. major deviations)	Reference

Comparable to OECD 403 GLP: no Rel. 2	Rat Sprague-Dawley m+f 10/sex/group	Formic acid purity 98% Lot/batch: no data 2.82, 6.60, 8.08, 10.6, 14.7 mg/l (analytical); 4.03, 8.50, 10.58, 13.40, 17.90 mg/l (nominal) 4 hours whole body vapour	7.4 mg/l (m+f) Males: 7.3 mg/l Females : 7.5 mg/l Clinical signs (in all treated groups): Closed lids, snout swiping, discharge from the nose and eye, corrosion of nose and eyes, salivation, corneal opacity, loss of pain reflex, dyspnea, respiration sounds, flatulence, apathy, hunched posture, unsteady gait Symptoms persisted until d14 after treatment (except for the 2.82 mg/l group: symptom free at d11) Mortality: within 7 days post exposure (inflated lungs, dilated hearts). BW at d7: dose-dependent decrease	BPD ID A6.1.3_01; FA_BPR_Ann_II_8_7_2_01 ██████████ 1980
---	--	--	---	--

Formic acid is of moderate toxicity via inhalation when tested in the rat. LC_{50} (4hrs) = 7.4 mg/l = 7400 mg/m³.

Following a 4-hour *inhalation* of formic acid vapours in rats (DocIIIA6.1.3-01; FA_BPR_Ann_II_8_7_2_01: ██████████, 1980), clinical signs indicated corrosive properties of the test substance, evidenced by the occurrence of corneal opacity and corrosion of the dorsal nose in some cases. Symptoms persisted until termination 14 days after the rats were exposed to 6.6 mg/l or above. Deaths occurred within 7 days. Inflated lungs and dilated hearts were seen in animals that died; gross pathology revealed no changes in animals sacrificed at termination.

Human case reports on acute 'accidental' inhalation (and dermal) exposure are rather rare. Besides local effects, severe acid skin burns and respiratory tract irritation, patients suffered and recovered rapidly from metabolic acidosis (described in section 3.3 and 3.14).

Note: the applicant has submitted a re-interpretation of the 1980 ██████████ study (FA_BPR_Ann_II_8_7_2_01-new) and concludes to a higher LC_{50} value of 7.85 mg/l. BE cannot accept this re-interpretation. The applicant's justification for this re-interpretation can be found in the PT3 specific BASF confidential Annex to the PT3 CAR, along with BE's clarification for refusal.

Value used in the Risk Assessment – Acute inhalation toxicity	
Value	LC ₅₀ 7.4 mg/l
Justification for the selected value	DocIIIA6.1.3-01; FA_BPR_Ann_II_8_7_2_01: ██████████ 1980 Acute inhalation toxicity of formic acid has been assessed in a study comparable to OECD 403.

3.2.4 Overall conclusion on acute toxicity

Value used in the Risk Assessment – Acute systemic toxicity	
Value	See below
Justification for the selected value	Appropriate studies are available for determining the LD ₅₀ oral and LC ₅₀ inhalation of formic acid. The acute toxic action profile of formic acid is predominantly determined by its inherent irritating/corrosive properties. The toxicity values after oral uptake and inhalation in rats suggest formic acid to be acutely harmful. The clinical signs give no evidence of specific systemic adverse effects.
Classification according to CLP and DSD	Acute toxicity, oral, cat. 4, H302 Acute toxicity, inhalation, cat. 3, H331 Corrosive properties determine the toxicity of formic acid; additional labelling EUH071

Value/conclusion used in the Risk Assessment – Acute local effects	
Value/conclusion	LD ₅₀ oral 730 mg/kg bw (formic acid) ⁵ LC ₅₀ inhalation 7.4 mg/l (formic acid) LD ₅₀ dermal >2000 mg/kg bw (Na formate)

⁵ Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

Justification for the selected value/conclusion	<p>Appropriate studies are available for determining the LD₅₀ oral and LC₅₀ inhalation of formic acid.</p> <p>Due to the corrosivity of formic acid, local effects occur at all dose levels. Pathological organ lesions recorded after oral administration included hyperemia of the stomach and intestines, congestion in spleens, mottled livers and kidneys, discoloration of kidneys and pancreas. Clinical signs after inhalation, closed lids, snout swiping, discharge from the nose and eye, corrosion of nose and eyes, salivation, corneal opacity, loss of pain reflex, dyspnea, respiration sounds, flatulence, apathy, hunched posture, unsteady gait, indicated the corrosive properties of formic acid, evidenced by the occurrence of corneal opacity and corrosion of the dorsal nose. The acute dermal toxicity of formic acid was not tested and not requested because of its corrosive properties.</p>
--	--

3.3 IRRITATION AND CORROSION

3.3.1 Skin corrosion and irritation

Summary table of animal studies on skin corrosion/irritation					
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance, Vehicle, Dose levels, Duration of exposure	Results <i>Average score (24, 48, 72 h), observations and time point of onset, reversibility, other adverse local/systemic effects, histopathological findings</i>	Remarks (e.g. major deviations)	Reference
n.a. corrosive substance; no <i>in vivo</i> testing acc to OECD 404 performed					
OECD 406 Buehler Test GLP: yes Rel. 1	Guinea pig Female 20/group 10 naïve controls	Formic acid purity 85.3% Induction: 7.5% formic acid in water challenge: 2% formic acid in water	Result: not sensitizing Pre-test: Min. irritant conc.: 5%; Max. non-irritant conc. 2%		BPD ID A6.1.5_01; FA_BPR_Ann_II_8_3_01 ██████████ 2002.

Summary table of human data on skin corrosion/irritation				
Type of data/ report, Reliability	Test substance	Relevant information about the study	Observations	Reference
Case report	Formic acid conc. not known	Route of exposure: dermal 1 male, 35-year-old	Accidental splash from a container on the maxilla, chin, around mouth, thorax Clinical signs: burning pain, sialorrhoea, nausea, vomiting Skin: blisters, necrotic areas Systemic: blood pressure 110/60, pulse and breathing regular, blood gases and acido-balance normal, no formic acid detected in blood and urine Result: Skin corrosion Reversible within 8 days	BPD ID A6.12.2_07a ; FA_BPR_Ann_II_8_12_2_07 Malizia et al., 1977
Case report	Formic acid undiluted, conc. not known	Route of exposure: dermal 1 female, 15-year-old	Accidental splash on lower extremities (20% of total body surface) Clinical signs: burns, nausea, vomiting (4 hrs after exposure = start treatment) Skin: depth of burns not determined, became full-thickness. Gross oedema on d2 and d3 without fever, ocular damage or pulmonary complications. Burns surgically revived on d16, grafted several times. Major scarring of burned areas persisted. Urine: brownish, haemoglobinuria Blood: pH 7.23, HCO ₃ 16.7 mmol/l, base deficit 9.5, haemolysis Patient recovered rapidly from metabolic acidosis	BPD ID A6.12.2_08 ; FA_BPR_Ann_II_8_12_2_08 Sigurdsson et al., 1983

			<p>Result: Skin corrosion Mild metabolic acidosis Reversibility: No: severe burns required several grafts, major scarring</p>	
Case report	Formic acid 90%	Route of exposure: dermal 1 female, 3-year-old	<p>Accidental splash on right torso and extremities (35% of total body surface) Clinical signs: severe distress (10 min after exposure = start treatment) Skin: full-thickness second- and third-degree burns. Required several skin grafts during several months Urine: initially dark red, haemoglobinuria resolved within few days without kidney failure Blood: pH 6.85, HCO₃ 16.7 mmol/l, base deficit -29.7 on 100% oxygen, bicarbonate 6mEq/l; initial serum formate level 400 µg/ml, haemolysis Patient recovered rapidly from metabolic acidosis. Result: Skin corrosion Metabolic acidosis Reversibility: No: severe burns required several grafts</p>	<p>BPD ID A6.12.2_09 ; FA_BPR_Ann_II_8_12_2_09 Chan et al., 1995</p>

No skin and eye irritation study reports are available on formic acid itself. Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive (according to DSD: C, R 35) in the EU (12th ATP) (see DOC-III A6.4.1_e / FA_BPR_Ann_II_8_2_0: Justification and A6.4.1_s/ FA_BPR_Ann_II_8_1_0: Justification).

A Buehler test was made available for assessment of skin sensitization (██████████ 2002; BPD ID A6.1.5_01; FA_BPR_Ann_II_8_3_01). There was no evidence of a sensitising potential in guinea pigs using the method of Buehler. During the irritation screen performed for this study with formic acid diluted in water, the minimum irritant concentration was found to be 5% formic acid in water; the maximum non-irritant concentration was found to be 2% formic acid in water.

Sodium formate [CAS No. 141-53-7] produced no skin irritation in an acute dermal toxicity test (see section 3.2.2).

Human data: see 3.14 for a detailed discussion.

The corrosive potential of formic acid has been reported on several occasions after accidental dermal exposure in humans and documented in case reports. Malizia et al., 1977 (DocIIIA6.12.2-07; FA_BPR_Ann_II_8_12_2_07) reported blisters and necrotic areas on the skin of a man after an accidental exposure from a formic acid splash on the face and thorax. The skin around the acid-burned region was hyperaemic and oedematous. The local skin corrosion was without signs of systemic toxicity. The patient recovered after 8 days.

Sigurdsson et al., 1983 (DocIIIA6.12.2-08; FA_BPR_Ann_II_8_12_2_08) reported an agricultural accident with a girl who's legs were hit by a splash of formic acid. The patient complained of nausea and vomited on arrival at the hospital. The burns turned out to be full-thickness. Gross oedema formed on d2 and d3. The burn was surgically revised and grafted. However, major scarring of the burned area persisted. Apart from the local skin corrosion and scarring, there was absorption of formic acid, which caused metabolic acidosis with hemolysis and hemoglobinuria.

Another accidental splash exposure on the right torso and extremities of a 3-year-old girl was reported by Chan et al., 1995 (DocIIIA6.12.2-08; FA_BPR_Ann_II_8_12_2_09). The patient was in severe distress. The dermal exposure to formic acid caused severe systemic toxicity: severe metabolic acidosis with haemolysis and haemoglobinuria. Only 10 minutes after the accident medical treatment started and further dermal absorption prevented. Nevertheless, the initial serum formate level was 400 µg/ml. Full-thickness second- and third-degree burns affected 35% of the total body surface, and required several grafts and long-term treatment.

Conclusion used in the Risk Assessment – Skin irritation and corrosivity	
Value/conclusion	Formic acid is corrosive to skin
Justification for the value/conclusion	No skin and eye irritation study reports are available on formic acid itself. Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive in the EU (12 th ATP) Harmonized classification and SCLs: Skin Corr 1A; H314 Skin Corr. 1B; H314: 10% ≤ C < 90% Skin Corr. 1A; H314: C ≥ 90%

	Skin Irrit. 2; H315: 2% ≤ C < 10%
--	-----------------------------------

Data waiving	
Information requirement	Skin irritation study on formic acid
Justification	Formic acid is a corrosive substance

3.3.2 Eye irritation

Conclusion used in Risk Assessment – Eye irritation and corrosivity	
Value/conclusion	Formic acid is corrosive to the eye
Justification for the value/conclusion	<p>No skin and eye irritation study reports are available on formic acid itself. Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive in the EU (12th ATP)</p> <p>Harmonized classification and SCLs: Skin corr 1A, H314 Eye Irrit. 2; H319: 2% ≤ C < 10%</p> <p>Additional proposed classification and SCLs: Eye dam/irrit 1, H318 Eye dam. 1; H318: C ≥ 10%</p>

Data waiving	
Information requirement	Eye irritation study on formic acid
Justification	Formic acid is a corrosive substance

3.3.3 Respiratory tract irritation

Summary table of animal studies on respiratory tract irritation					
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance Dose levels, Duration of exposure	Results <i>clinical signs, histopathology, reversibility</i>	Remarks <i>(e.g. major deviations)</i>	Reference
Alarie (1973), ASTM (1984). GLP: yes Rel. 1	Mouse, Swiss Webster male, 5/group	Formi®LHS Potassium formate (1:2) Purity 99.8% Lot/batch: B0298 Inhalation nebulation, nose- only 267, 568, 622, and 802 mg/m ³ (analytical) MMAD: 1.2µm 10 min acclimatisation, 30 min exposure, 20 min recovery	RD50 = 615 mg/m ³ [RD50 = 608 mg/m ³ (Alarie, 1973) RD50 = 623 mg/m ³ (Bos et al., 1992)] ~ weak sensory irritant of the upper respiratory tract Asymptomatic decrease of the breathing rate at exposure time. Max. decrease towards the end. RD50 calculated from the mean of the 7 last measurements (minutes 18 to 30 of exposure). No other changes in behaviour. Breathing rate returned to normal within the recovery period. The tidal volume was not affected by treatment. Necropsy: 1 petechia noted in 1 lung lobe of 1 animal. Similar occasional findings in unexposed animals. No mortality. Reversibility: Yes, within the 20-min recovery period		BPD ID A6.1.6_01; FA_BPR_Ann_II_8_13_2 _01 ██████████ (1999)
In accordance with OECD 413	Rat, Fischer 344/N, m + f	Formic acid purity 95%	NOAEL _{local} : 30 mg/m ³		BPD ID A6.4.3_01; FA_BPR_Ann_II_8_9_2_

GLP: yes Rel. 1	10/sex	0, 15, 30, 61, 122, 244 mg/m ³ (nominal) 6h/d, 5d/wk, 13 weeks Vapour, whole body	<p>LOAEL_{local}: 61 mg/m³</p> <p>No clinical signs</p> <p>Local effects: nasal irritation, squamous metaplasia of the respiratory epithelium, olfactory degeneration, severity minimal to mild.</p> <p>Respiratory epithelium squamous metaplasia:</p> <table border="1"> <tr> <td>mg/m³</td> <td>0</td> <td>15</td> <td>30</td> <td>61</td> <td>122</td> <td>244</td> </tr> <tr> <td>male</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>9</td> </tr> <tr> <td>female</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>6</td> </tr> </table> <p>Olfactory epithelium degeneration: minimal to mild</p> <table border="1"> <tr> <td>mg/m³</td> <td>0</td> <td>15</td> <td>30</td> <td>61</td> <td>122</td> <td>244</td> </tr> <tr> <td>male</td> <td>0</td> <td>0</td> <td>0</td> <td>1</td> <td>1</td> <td>9</td> </tr> <tr> <td>female</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>5</td> </tr> </table>	mg/m ³	0	15	30	61	122	244	male	0	0	0	0	0	9	female	0	0	0	0	0	6	mg/m ³	0	15	30	61	122	244	male	0	0	0	1	1	9	female	0	0	0	0	0	5		03 Thompson, 1992
mg/m ³	0	15	30	61	122	244																																									
male	0	0	0	0	0	9																																									
female	0	0	0	0	0	6																																									
mg/m ³	0	15	30	61	122	244																																									
male	0	0	0	1	1	9																																									
female	0	0	0	0	0	5																																									
In accordance with OECD 413 GLP: yes Rel. 1	Mice B6C3F ₁ m + f 10/sex	Formic acid purity 95% 0, 15, 30, 61, 122, 244 mg/m ³ (nominal) 6h/d, 5d/wk, 13 weeks Vapour, whole body	<p>NOAEL_{local}: 61 mg/m³</p> <p>LOAEL_{local}: 122 mg/m³</p> <p>No clinical signs</p> <p>Local effects: nasal irritation, olfactory degeneration, severity minimal but dose-related.</p> <p>Olfactory epithelium degeneration: minimal</p> <table border="1"> <tr> <td>mg/m³</td> <td>0</td> <td>15</td> <td>30</td> <td>61</td> <td>122</td> <td>244</td> </tr> <tr> <td>male</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> </tr> <tr> <td>female</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> <td>5</td> </tr> </table>	mg/m ³	0	15	30	61	122	244	male	0	0	0	0	0	2	female	0	0	0	0	2	5		BPD ID A6.4.3_01; FA_BPR_Ann_II_8_9_2_04 Thompson, 1992																					
mg/m ³	0	15	30	61	122	244																																									
male	0	0	0	0	0	2																																									
female	0	0	0	0	2	5																																									

Summary table of human data on respiratory tract irritation				
Type of data/report, Reliability	Test substance	Relevant information about the study	Observations	Reference
Case report	Formic acid 98%	Route of exposure: inhalation 1 male, 39-year-old	<p>Accidental spray (aerosol) onto the face with concomitant inhalation</p> <p>Clinical signs: facial burns (3% of total body surface), dyspnoea</p> <p>Nasopharyngoscopy: mild supraglottic erythema, normal vocal cords</p> <p>Skin: second-degree burns</p> <p>Pulmonary function tests: Vital capacity reduced on d1, recovered largely within 14 days. Complains of dyspnoea till d15</p> <p><u>Day 1</u></p> <p>FVC (L): 3.74 (79% predicted) FEV₁ (L): 2.86 (73% predicted) FEV₁/FVC: 76.38 (92% predicted) FEF_{25%-75%} (l/sec): 2.32 (56% predicted)</p> <p><u>Day 15</u></p> <p>FVC (L): 4.35 (92% predicted) FEV₁ (L): 3.62 (92% predicted) FEV₁/FVC: 83.09 (101% predicted) FEF_{25%-75%} (l/sec): 3.82 (92% predicted)</p> <p>Reversible Pulmonary dysfunction: Reactive Airway Dysfunction Syndrome Reversible within 15 days</p>	BPD ID A6.12.2_10 ; FA_BPR_Ann_II_8_12_2_10 Yelon et al., 1996

The airway irritating properties were studied by exposing mice to potassium diformate at concentrations of 267, 568, 622, 802 mg/m³ for a single period of 30 minutes (DocIIIA6.1.6-01; FA_BPR_Ann_II_8_13_2_01: ██████████ 1999). Animals were necropsied 7 days after exposure. Inhalation of nebulized potassium diformate solutions irritated the upper airways and caused a decrease of the respiratory rate and post-inspiratory apnoea in a concentration-dependent manner. Treatment-related changes in tidal volume were not observed. The RD₅₀ values were

obtained with two calculation methods, which were in good agreement. Since the RD_{50} values found at each concentration level did not increase or decrease with increasing concentrations, it was concluded that except sensory irritation other possible toxic actions were absent. No other effects were observed (behaviour, body weight, lung weight, macroscopic and histopathological findings: lungs, nasal cavity). The overall RD_{50} was 615 mg/m^3 . This study detected clearly the irritating effects caused by potassium diformate, however without any histopathological changes. As such, this data does not allow a conclusion on a relationship between the RD_{50} and the concentration inducing histopathological changes in the respiratory tract.

In addition, in the acute inhalation study in rats (see 3.2. Acute toxicity) clinical signs indicated the corrosive properties of formic acid, evidenced by the occurrence of corneal opacity and corrosion of the dorsal nose. Symptoms persisted until termination 14 days after the rats had been exposed to 6600 mg/m^3 and above.

Further evidence of respiratory tract irritation is found in the histopathological data of the nasal cavity of the repeated dose inhalation toxicity studies performed with formic acid vapours (13-week inhalation, rat, mouse). See section 3.6. for a more detailed discussion.

Subchronic 13-week inhalation studies with formic acid vapour at concentrations of 0, 15, 30, 61, 122, 244 mg/m^3 were conducted in rats and mice (DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_03 and DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_04: Thompson, 1992). Both in the rat and the mouse, the inhalation of formic acid did not result in clinical effects. In the rat, microscopic changes occurred in the respiratory and olfactory epithelium of the nose. Changes on the respiratory epithelium consisted of a minimal squamous metaplasia in which the pseudostratified, ciliated columnar cells were replaced by a flattened, non-ciliated epithelium with approximately 2 to 5 cells in thickness. Squamous metaplasia occurred most often in the respiratory epithelium that lines the most dorsal portion of the dorsal meatus in the nose's anterior section (Level I). In the olfactory epithelium, degenerative changes were minimal to mild and generally limited to the area of the dorsal meatus in the mid-nasal section (Level II). Degeneration was characterised by a loss of the usual orderly arrangement of the pseudostratified layer of nuclei and by a slight reduction on the normal thickness of the olfactory epithelium. There was no necrosis. No evidence was seen of metaplasia of the olfactory epithelium or atrophy of the nerve fibres in the olfactory mucosa. In the mouse, microscopic changes were limited to the degeneration of the olfactory epithelium of the nose. The minimal degeneration occurred in the dorsal portion of the dorsal meatus in the anterior or mid-nasal section (Levels I and II). Degeneration was characterised by a loss of the usual orderly arrangement of the pseudostratified layer of nuclei and by a slight reduction on the normal thickness of the olfactory epithelium. In conclusion, both in the rat and the mouse the upper respiratory tract was the major target for toxicity. The overall $LOAEC_{\text{local}} = 122 \text{ mg formic acid/m}^3$ and $NOAEC_{\text{local}} = 60 \text{ mg formic acid/m}^3$, based on histological changes in the nasal region in both the rat and the mouse.

Human data

Due to the warning effect of the pungent smell of formic acid, only few human data due to (accidental) inhalation exposure is available. Yelon et al., 1996 (DocIIIA6.12.2-10; FA_BPR_Ann_II_8_12_2_10) reported a case of an inhalation injury as a result of aerosolized formic acid from an accidental spray in the face. Apart from the skin burns, the man complained of dyspnea. Despite the oxygen therapy and nebulised metaproterenol therapy, the patient continued to complain of dyspnea, it even worsened. Pulmonary function tests within the first 12 hours were consistent with mild restrictive disease (FEV_1 of 2.86L, 73% of predicted; normal FEV_1/FVC of 76.38%); the $FEF_{25\%-75\%}$ of 2.32L/sec (56% predicted) was consistent with small airway dysfunction. On day 3, the patient had improvement in dyspnea, but developed a nonproductive

cough at the same time. The patient continued to complain of dyspnea on moderate-severe exertion. The patient recovered slowly. As all criteria were met, the patient could be diagnosed with Reactive Airway Dysfunction Syndrome (RADS).

Based on physico-chemical data, animal data and human findings, the corrosive nature of formic acid is found to affect the respiratory tract. We propose additional labelling with EUH071, 'corrosive to the respiratory tract', as the corrosive properties determine the toxicity of formic acid (CLP Regulation Annex II, point 1.2.6).

Conclusion used in the Risk Assessment – Respiratory tract irritation	
Conclusion	formic acid is to be classified as EUH071, corrosive to the respiratory tract.
Justification for the conclusion	The corrosive properties of formic acid have been observed to affect the respiratory tract in appropriate studies relating to inhalation toxicity and in a human case report.

3.3.4 Overall conclusion on corrosion and irritation

Conclusion used in the Risk Assessment – Corrosion and irritation	
Value	Formic acid is corrosive to skin and eye, and to the respiratory tract.
Justification for the selected value	No skin and eye irritation study reports are available on formic acid itself. Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive in the EU (12 th ATP) The corrosive properties of formic acid were evidenced by numerous human case reports. In addition, based on physico-chemical data, animal data (acute inhalation toxicity, respiratory irritation test, repeated inhalation toxicity) and human findings, formic acid is observed to affect the respiratory tract. For NOAEC _{local} see 13-week inhalation study, rat, mouse; section 3.6.3 below). RD50 = 615 mg potassium diformate/m ³ .
Classification according to CLP and DSD	Harmonized classification and SCLs: Skin corr 1A, H314 Skin Corr. 1B; H314: 10% ≤ C < 90% Skin Corr. 1A; H314: C ≥ 90% Skin Irrit. 2; H315: 2% ≤ C < 10% Eye Irrit. 2; H319: 2% ≤ C < 10%

	<p>Additional proposed classification and SCLs: Eye dam/irrit 1, H318 Eye dam. 1; H318: C ≥ 10% EUH071</p>
--	--

3.4 SENSITISATION

3.4.1 Skin sensitisation

Summary table of animal studies on skin sensitisation					
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance, Vehicle, Dose levels, Route of exposure (topical/intradermal, if relevant), Duration of exposure	Results (EC3-value or amount of sensitised animals at induction dose)	Remarks (e.g. major deviations)	Reference
OECD 406 Buehler Test GLP: yes Rel. 1 Inductions, topically on d0, d7, d14; Challenge, topically on d28 Scoring 1 on d29 Scoring 2 on d30 Evaluation according to Magnusson and Kligman:	Guinea pig Female 20/group 10 naïve controls	Formic acid purity 85.3% Induction: 7.5% formic acid in water challenge: 2% formic acid in water	Result: not sensitizing Scoring after 24h: naïve control: 0/10 formic acid: 0/20 pos. control*: 13/20 Scoring after 48h: naïve control: 0/10 formic acid: 0/20 pos. control*: 14/20 Pre-test: Min. irritant conc.: 5%; Max. non-irritant conc. 2%		BPD ID A6.1.5_01; FA_BPR_Ann_II_8_3_01 ██████████ 2002.

<p>0=no visible change 1=discrete or patchy erythema 2= moderate and confluent erythema 3= intense erythema and swelling</p>			<p>Observations after induction 1, 2, 3: Discrete to moderate erythema in 20/20 test animals. Mean score 1.65, 1.85, 2.45, respectively. Swelling in 10/20, 11/20, 9/20 test animals, respectively.</p> <p>*Pos control : α-HCA, historic data 24h: 13/20 48h: 14/20</p>																																																																	
<p>OECD 406 GPMT GLP: yes Rel. 1</p>	<p>Guinea pig Female 20/group 10 naïve controls</p>	<p>Formi®LHS Potassium formate (1:2)</p> <p>Intradermal induction: 0.5% m/v in purified water and/or adjuvant Topical induction: 15% m/m in Vaseline Challenge application: 10 and 5% m/m in vaseline</p>	<p>Result: not sensitizing</p> <p><u>Incidences naïve control group:</u></p> <table border="1" data-bbox="965 635 1413 1118"> <thead> <tr> <th rowspan="2">Rea- ding time</th> <th rowspan="2">Con- centr- a-tion</th> <th colspan="4">Incidence</th> <th rowspan="2">% anima ls with incide n-ces ≥ 1</th> </tr> <tr> <th>0</th> <th>1</th> <th>2</th> <th>≥ 3</th> </tr> </thead> <tbody> <tr> <td>24h</td> <td>10%</td> <td>8</td> <td>2</td> <td>0</td> <td>0</td> <td>20%</td> </tr> <tr> <td>48h</td> <td>10%</td> <td>9</td> <td>1</td> <td>0</td> <td>0</td> <td>10%</td> </tr> <tr> <td>24h</td> <td>5%</td> <td>10</td> <td>0</td> <td>0</td> <td>0</td> <td>0%</td> </tr> <tr> <td>48h</td> <td>5%</td> <td>10</td> <td>0</td> <td>0</td> <td>0</td> <td>0%</td> </tr> <tr> <td>24h</td> <td>0%</td> <td>10</td> <td>0</td> <td>0</td> <td>0</td> <td>0%</td> </tr> <tr> <td>48h</td> <td>0%</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0%</td> </tr> </tbody> </table> <p><u>Incidences test group:</u></p> <table border="1" data-bbox="965 1214 1525 1385"> <thead> <tr> <th>Rea- ding time</th> <th>Con- centra- tion</th> <th>Incidence</th> <th>% anima ls with incide n-ces ≥ 1</th> <th>% of anim als with</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	Rea- ding time	Con- centr- a-tion	Incidence				% anima ls with incide n-ces ≥ 1	0	1	2	≥ 3	24h	10%	8	2	0	0	20%	48h	10%	9	1	0	0	10%	24h	5%	10	0	0	0	0%	48h	5%	10	0	0	0	0%	24h	0%	10	0	0	0	0%	48h	0%	0	0	0	0	0%	Rea- ding time	Con- centra- tion	Incidence	% anima ls with incide n-ces ≥ 1	% of anim als with							<p>Report number: 1516/22-1032, ██████████ 1998 BPD ID A6.1.5_02; FA_BPR_Ann_II_8_3_0 3 SIAP (2008)</p>
Rea- ding time	Con- centr- a-tion	Incidence				% anima ls with incide n-ces ≥ 1																																																														
		0	1	2	≥ 3																																																															
24h	10%	8	2	0	0	20%																																																														
48h	10%	9	1	0	0	10%																																																														
24h	5%	10	0	0	0	0%																																																														
48h	5%	10	0	0	0	0%																																																														
24h	0%	10	0	0	0	0%																																																														
48h	0%	0	0	0	0	0%																																																														
Rea- ding time	Con- centra- tion	Incidence	% anima ls with incide n-ces ≥ 1	% of anim als with																																																																

							react ions ⁶
		0	1	2	≥ 3		
24h	10%	14	6	0	0	30%	0%
48h	10%	19	1	0	0	5%	0%
24h	5%	16	4	0	0	20%	10%
48h	5%	20	0	0	0	0%	0%
24h	0%	18	2	0	0	10%	-
48h	0%	20	0	0	0	0%	-

Pre-test:
Min. irritant conc.:
- Intradermal injection: 0.5%
- Topical application: 15%
Max. non-irritant conc: 10%

Observation after intradermal injection:
Well-defined erythema was noted at injection sites with Freund's Complete Adjuvant (FCA) for both test and control animals.

Observation after topical induction:
Slight erythema was apparent in test animals following application of 15% Formi®LHS in Vaseline.

⁶ Data interpretation: The incidences of the test animals were compared to the naïve control animals at the same concentration and reading time. If the challenge response of a test animal was less marked or the same as the maximum reaction apparent among naïve control animals at the same concentration and reading time, those animals were not counted as animals with reactions. Furthermore, the percentage of test animals with reactions was reduced by the percentage of test animals with reactions treated with vaseline alone.

			<p>No erythema was apparent at the topical application sites in the control animals.</p> <p><u>Positive control:</u> 2-Mercaptobenzo-thiazole (MBTZ), historic control data 03-04/1997: 6/9 positive, 2/9 inconclusive, 1/9 negative 08-09/1997: 6/10 positive, 2/10 inconclusive, 2/10 negative</p>		
--	--	--	--	--	--

No guinea-pig maximisation test on the active substance, formic acid, was made available by the applicant. Instead, a Buehler test was made available. Nevertheless, the conduct of an additional Maximisation test (GPMT) is scientifically not justified. A negative GPMT result was obtained with potassium diformate, that liberates formate and formic acid in equimolar quantities in aqueous solution. This substance was included in the "Formic acid and formates" category that was treated in the OECD/ICCA-HPV program, and the negative result can be read across to formic acid. The final SIAP (2008) is publicly available at: <http://webnet.oecd.org/Hpv/UI/handler.axd?id=81d8d2fe-5244-4699-93ab-c501433db94c>. In the concept of skin sensitisation it is generally assumed that protein-hapten conjugates need to be formed by covalent binding in order to be recognised by the immune system. Therefore, a compound which is able to cause contact allergy must have electrophilic properties, either by itself or after metabolic transformation. This concept is generally accepted and provides the mechanistic basis for Structure-activity-relations (SAR) for the skin sensitisation endpoint. Both formic acid and formate lack electrophilic properties, and are, therefore, considered to lack sensitising properties. In fact formic acid is not contained in publicly available structural alert lists, and acknowledged recently available QSAR models (CAESAR, OASIS) predict that formic acid is not a skin sensitizer. The negative result of the Buehler test with formic acid in Guinea pigs fits into the described concept. Additionally, no case reports of skin sensitisation following skin contact of workers or of the general public were retrieved. Case reports of accidental dermal exposure to formic acid also do not indicate that skin sensitisation was seen. The considerations on structure and electrophilicity do not suggest the conduct of a GPMT. Under REACH the conduct of a maximisation test is not allowed because formic acid is corrosive to the skin. The RMS BE accepted the justification given by the applicant.

There was no evidence of a sensitising potential in guinea pigs using the method of Buehler. During the irritation screen with formic acid diluted in water, the minimum irritant concentration was found to be 5% formic acid in water; the maximum non-irritant concentration was found to be 2% formic acid in water. The inductions performed with 7.5% formic acid caused discrete or patchy erythema to intense erythema, swelling and eczematoid skin changes. No sensitisation responses were elicited by formic acid: no visual changes (score=0) were observed in both the naïve control and test animals. In contrast, the positive control (not included, but routinely conducted twice a year in the laboratory) showed a clear sensitising effect, which confirmed the validity of the study.

A GMPT was performed with potassium diformate (██████████ 1998; BPD ID A6.1.5_02; FA_BPR_Ann_II_8_3_03/SIAP 2008). In the pre-test a topical minimal irritation concentration of 15% and a maximal non-irritant concentration of 10% were established. For the intradermal

injection 0.5% with and without Freund's Complete Adjuvant (FCA) were used. Well defined erythema was noted for both test and control animals after intradermal injections with FCA. No erythema was apparent in test animals receiving the test substance without FCA and in control animals receiving purified water alone. During the induction slight erythema was apparent in test animals following topical application of 15% potassium diformate in Vaseline. No erythema was apparent at the topical application sites in the control animals. During the challenge application light erythema was noted in some control and test animals treated with the higher challenge concentration (10%). In addition, four test animals showed slight erythema at the lower challenge application site although two of these animals also had a slight response to application of the vehicle Vaseline. Those two animals were therefore not considered in the assessment of animals with reactions. The reactions had generally resolved by the 48-hour assessment, and it was noted that the dermal reactions seen in the test group animals were no more persistent or marked than those seen among the controls. In conclusion, it can be stated that no evidence of skin sensitising properties of potassium diformate was observed.

In addition, there is no data available (human data e.g. market surveillance data, animal data, open literature) which may be indicative of the potential of formic acid to cause skin sensitisation and sensitisation by inhalation in humans.

Conclusion used in Risk Assessment – Skin sensitisation	
Value/conclusion	Formic acid does not fulfill the criteria of the CLP regulation to be classified as a skin sensitiser
Justification for the value/conclusion	Skin sensitization (Buehler test) by formic acid has been assessed in an OECD 406 study (Buehler test). The results do not trigger a classification as skin sensitizer.

Data waiving	
Information requirement	Local Lymph Node Assay (LLNA),
Justification	. LLNA not available as FA is corrosive to skin: Step 2, Point 8.3, Title 1, Annex II of EU 528/2012 indicates in vivo testing (preferably with the LLNA) does not need to be conducted if the substance is classified for corrosivity.

3.4.2 Respiratory sensitisation

Conclusion used in the Risk Assessment – Respiratory sensitisation	
Value/conclusion	There is no indication that formic acid would be a respiratory sensitizer.
Justification for the value/conclusion	No data are available (human data e.g. market surveillance data, animal data, open literature) which may be indicative of the potential of formic acid to cause sensitisation by inhalation in humans. No respiratory sensitisation was seen with formic acid in two subchronic rat and mouse inhalation studies (see 3.6.3, Thompson 1992). Hence, there is no indication that formic acid would be a respiratory sensitizer.

3.4.3 Overall conclusion on sensitisation

Conclusion used in the Risk Assessment – Sensitisation	
Value	Formic acid is not a skin sensitizer. There is no indication that formic acid would be a respiratory sensitizer.
Justification for the selected value	Classification as a sensitizer is not triggered by appropriate tests. Studies in guinea pigs (method of Buehler) showed that there is no evidence that formic acid has a potential to induce skin sensitisation. In addition, there are no data available (human data including market surveillance, animal studies, open literature) that may be indicative of the potential of formic acid to cause skin sensitisation and sensitisation by inhalation in humans.
Classification according to CLP and DSD	none

3.5 SHORT TERM REPEATED DOSE TOXICITY

3.5.1 Short-term oral toxicity

No data are available on short-term oral toxicity.

Value used in the Risk Assessment – Short-term oral toxicity	
Value/conclusion	The short-term toxicity of formic acid has not been investigated.
Justification for the value/conclusion	The additional conduct of a study with repeated administration via the oral, dermal, or inhalation route was not considered to be necessary.

Data waiving	
Information requirement	short-term oral toxicity of formic acid
Justification	According to the Guidance on the BPR VIII Human Health – Part A Information Requirements (ECHA, 2014), no studies are required because subchronic rodent toxicity studies are available for the oral route (rat, potassium diformate). The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01; FA_BPR_Ann_II_8_8_01: ██████████ 1997).

3.5.2 Short-term dermal toxicity

No data are available on short-term dermal toxicity.

Value used in the Risk Assessment – Short-term dermal toxicity	
Value/conclusion	The short-term toxicity of formic acid has not been investigated.

Justification for the value/conclusion	The additional conduct of a study with repeated administration via the oral, dermal, or inhalation route was not considered to be necessary.
---	--

Data waiving	
Information requirement	short-term dermal toxicity of formic acid
Justification	Dermal repeated dose studies were not conducted for reasons of animal welfare, because formic acid and potassium diformate are both corrosive to the skin. Moreover, only limited repeated exposure is expected because of the corrosivity to the skin.

3.5.3 Short-term inhalation toxicity

No data are available on short-term inhalation toxicity.

Value used in Risk Assessment – Short-term inhalation toxicity	
Value/conclusion	The short-term toxicity of formic acid has not been investigated.
Justification for the value/conclusion	The additional conduct of a study with repeated administration via the oral, dermal, or inhalation route was not considered to be necessary.

Data waiving	
Information requirement	short-term inhalation toxicity of formic acid
Justification	According to the Guidance on the BPR VIII Human Health – Part A Information Requirements (ECHA, 2014), no studies are required because subchronic rodent toxicity studies are available for the inhalation route of exposure (rat and mouse, formic acid).

3.5.4 Overall conclusion on short-term repeated dose toxicity

Value used in the Risk Assessment – Short-term repeated dose systemic toxicity	
Value	The short-term toxicity of formic acid has not been investigated.
Justification for the selected value	<p>The additional conduct of a study with repeated administration via the oral, dermal, or inhalation route was not considered to be necessary.</p> <p>According to the Guidance on the BPR VIII Human Health – Part A Information Requirements (ECHA, 2014), no studies are required because subchronic rodent toxicity studies are available for the oral route (rat, potassium diformate) and the inhalation route of exposure (rat and mouse, formic acid). The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01: ██████████ 1997). Dermal repeated dose studies were not conducted for reasons of animal welfare, because formic acid and potassium diformate are both corrosive to the skin. Moreover, only limited repeated exposure is expected because of the corrosivity to the skin.</p>
Classification according to CLP and DSD	n.a.

Value/conclusion used in the Risk Assessment – Short-term repeated dose local effects	
Value/conclusion	The short-term toxicity of formic acid has not been investigated.
Justification for the selected value/conclusion	<p>The additional conduct of a study with repeated administration via the oral, dermal, or inhalation route was not considered to be necessary.</p> <p>According to the Guidance on the BPR VIII Human Health – Part A Information Requirements (ECHA, 2014), no studies are required because subchronic rodent toxicity studies are available for the oral route (rat, potassium diformate) and the inhalation route of exposure (rat and mouse, formic acid). The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01: ██████████ 1997). Dermal repeated dose studies were not conducted for reasons of animal welfare, because formic acid and potassium diformate are both corrosive to the skin. Moreover, only limited repeated exposure is expected because of the corrosivity to the skin.</p>

Classification according to CLP and DSD	n.a.
--	------

3.6 SUB-CHRONIC REPEATED DOSE TOXICITY

3.6.1 Sub-chronic oral toxicity

Summary table of oral sub-chronic animal studies (usually 90-day studies)						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/ group	Test substance Dose levels, Route of exposure (gavage, in diet, other), Duration of exposure	NOAEL, LOAEL	Results	Remarks (e.g. major deviations)	Reference
OECD 408 GLP: yes Rel. 1	Rat, CrI:CDBR m + f 10/sex/group 10/sex/satellite group	KHCO ₂ •H ₂ CO ₂ [CAS No. 20642-05-1] purity 95% 0, 600, 1200, 3000 mg Formi/kg bw/d (nominal) = 0, 420, 840, 2100 mg formate/kg bw/d Oral, feed 13 wk, 4 wk recovery continuous, 7 d/week	NOAEL _{Local} : as formate: <420 mg/kg bw/d LOAEL _{Local} : as formate: 420 mg/kg bw/d NOAEL _{Systemic} : as formate: 840 mg/kg bw/d LOAEL _{Systemic} : as formate:	No clinical signs No active substance related mortality. Local effects: gastric irritation = thickening of the stomach, usually involving the limiting ridge, dose-related increase in severity and incidence of squamous cell hyperplasia in the stomach (m + f) partial reversibility during the treatment-free period Systemic or target organ toxicity: not overt Bw: dose-dependent decrease in bw (males), decrease in bw at high dose (females); in the recovery period, the bw gain was in parallel for the high dose and control group,		BPD ID A6.4.1_01 FA_BPR_Ann_II_8_9_2_01 [REDACTED] 1998

			2100 mg/kg bw/d	<p>but no increase in body weight gain compared to the control.</p> <p>Food intake: only slight but dose-dependent decrease in food consumption (not stat. sign.), in recovery period comparable food intake for all groups.</p> <p>Haematology at week 13:</p> <p>Haematology at week 13:</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">Males</th> <th colspan="3">Females</th> </tr> <tr> <th>Low</th> <th>Int.</th> <th>High</th> <th>Low</th> <th>Int.</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>RBC</td> <td></td> <td></td> <td>↑</td> <td></td> <td></td> <td></td> </tr> <tr> <td>MCV</td> <td></td> <td></td> <td>↓</td> <td></td> <td>↓</td> <td>↓</td> </tr> <tr> <td>MCH</td> <td></td> <td></td> <td>↓</td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>MCHC</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↑</td> </tr> <tr> <td>Platelet</td> <td></td> <td></td> <td>↑</td> <td></td> <td></td> <td></td> </tr> <tr> <td>WBC</td> <td></td> <td></td> <td>↑</td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <p>Clinical chemistry at week 13:</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">Males</th> <th colspan="3">Females</th> </tr> <tr> <th>Low</th> <th>Int.</th> <th>High</th> <th>Low</th> <th>Int.</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>AST</td> <td></td> <td></td> <td>↑ (trend)</td> <td></td> <td></td> <td></td> </tr> <tr> <td>AP</td> <td></td> <td></td> <td>↑ (trend)</td> <td></td> <td></td> <td></td> </tr> <tr> <td>Tot bilir</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>Tot prot</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>K</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>Ca</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>Creatinine</td> <td></td> <td>↓</td> <td>↑</td> <td>↓</td> <td></td> <td></td> </tr> <tr> <td>Urea</td> <td></td> <td></td> <td></td> <td></td> <td>↑</td> <td>↑</td> </tr> <tr> <td>Tot Chol</td> <td></td> <td></td> <td></td> <td></td> <td>↑</td> <td>↑</td> </tr> <tr> <td>Globulin</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> <tr> <td>A/G ratio</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>↓</td> </tr> </tbody> </table>		Males			Females			Low	Int.	High	Low	Int.	High	RBC			↑				MCV			↓		↓	↓	MCH			↓			↓	MCHC						↑	Platelet			↑				WBC			↑					Males			Females			Low	Int.	High	Low	Int.	High	AST			↑ (trend)				AP			↑ (trend)				Tot bilir						↓	Tot prot						↓	K						↓	Ca						↓	Creatinine		↓	↑	↓			Urea					↑	↑	Tot Chol					↑	↑	Globulin						↓	A/G ratio						↓	
	Males			Females																																																																																																																																																		
	Low	Int.	High	Low	Int.	High																																																																																																																																																
RBC			↑																																																																																																																																																			
MCV			↓		↓	↓																																																																																																																																																
MCH			↓			↓																																																																																																																																																
MCHC						↑																																																																																																																																																
Platelet			↑																																																																																																																																																			
WBC			↑																																																																																																																																																			
	Males			Females																																																																																																																																																		
	Low	Int.	High	Low	Int.	High																																																																																																																																																
AST			↑ (trend)																																																																																																																																																			
AP			↑ (trend)																																																																																																																																																			
Tot bilir						↓																																																																																																																																																
Tot prot						↓																																																																																																																																																
K						↓																																																																																																																																																
Ca						↓																																																																																																																																																
Creatinine		↓	↑	↓																																																																																																																																																		
Urea					↑	↑																																																																																																																																																
Tot Chol					↑	↑																																																																																																																																																
Globulin						↓																																																																																																																																																
A/G ratio						↓																																																																																																																																																

				<p>Clinical chemistry at week 17:</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Males</th> <th colspan="2">Females</th> </tr> <tr> <th>Low</th> <th>High</th> <th>Low</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>AST</td> <td></td> <td>↑</td> <td></td> <td></td> </tr> <tr> <td>AP</td> <td></td> <td>↑</td> <td></td> <td></td> </tr> <tr> <td>Glucose</td> <td></td> <td>↓</td> <td></td> <td>↓</td> </tr> </tbody> </table> <p>all changes considered of no biological relevance, no dose-response and no microscopic changes observed.</p> <p>Absorption study (high-dose): formate plasma levels morning: 90 µg (f)-160 µg (m) formate/ml afternoon: < LOD rapid absorption and metabolism, no accumulation</p>		Males		Females		Low	High	Low	High	AST		↑			AP		↑			Glucose		↓		↓	
	Males		Females																										
	Low	High	Low	High																									
AST		↑																											
AP		↑																											
Glucose		↓		↓																									
<p>No guideline, but following scientific standards GLP: yes Rel. 2</p>	<p>Pig, Large White x Landrace hybrid breed f 6/group</p>	<p>$\text{KHCO}_2 \cdot \text{H}_2\text{CO}_2$ [CAS No. 20642-05-1] purity 98.7%</p> <p>0, 1.2%, 3.0%, 6.0% in the diet</p> <p><i>Mating to farrowing:</i> 0, 132, 324, 624 mg/(kg*d) = 0, 92, 226,</p>	<p>NOAEL_{Local}: as formate: < 149 mg/kg bw/d</p> <p>LOAEL_{Local}: as formate: 149 mg/kg bw/d</p> <p>NOAEL_{Systemic}: as formate:</p>	<p>No clinical signs</p> <p>No active substance related mortality.</p> <p>Local effects: gastric irritation = forestomach gastritis and erosion/ulcer in approx. 30 to 60% of the treated animals.</p> <p>No systemic or target organ toxicity:</p>	<p>BPD ID A6.4_02 FA_BPR_Ann_II_8_9_2_02 ██████████ 2004</p>																								

		<p>437 mg formate/(kg*d)</p> <p><i>Farrowing to weaning:</i> 0, 213, 513, 1086 mg/(kg*d) = 0, 149, 359, 760 mg formate/(kg*d)</p> <p>Oral, feed</p> <p>140 days: 114 from service to farrowing, and until day 26 post-partum</p> <p>continuous, 7 d/week</p>	<p>760 mg/kg bw/d</p> <p>LOAEL_{Systemic}: as formate: >760 mg/kg bw/d</p>	<p>No effect on bw (gain) and food intake.</p> <p>Haematology:</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">Week 15</th> <th colspan="3">Weaning</th> </tr> <tr> <th>Low</th> <th>Int.</th> <th>High</th> <th>Low</th> <th>Int.</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>RBC*</td> <td colspan="6">dose-dependent trend ↓</td> </tr> <tr> <td>Hb*</td> <td colspan="6">dose-dependent trend ↓</td> </tr> <tr> <td>WBC</td> <td colspan="6">dose-dependent trend ↓</td> </tr> <tr> <td>PCV</td> <td colspan="6">dose-dependent trend ↓</td> </tr> </tbody> </table> <p>* stat. sign. at week 15</p> <p>Clinical chemistry:</p> <table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="3">Week 15</th> <th colspan="3">Weaning</th> </tr> <tr> <th>Low</th> <th>Int.</th> <th>High</th> <th>Low</th> <th>Int.</th> <th>High</th> </tr> </thead> <tbody> <tr> <td>AP</td> <td colspan="6"></td> </tr> <tr> <td>K*</td> <td></td> <td>↑</td> <td></td> <td>↓</td> <td></td> <td>↑</td> </tr> </tbody> </table> <p>* dose-dependent.</p> <p>Reproduction parameters not affected.</p> <p>Development of piglets not affected at birth and until weaning.</p>		Week 15			Weaning			Low	Int.	High	Low	Int.	High	RBC*	dose-dependent trend ↓						Hb*	dose-dependent trend ↓						WBC	dose-dependent trend ↓						PCV	dose-dependent trend ↓							Week 15			Weaning			Low	Int.	High	Low	Int.	High	AP							K*		↑		↓		↑		
	Week 15			Weaning																																																																						
	Low	Int.	High	Low	Int.	High																																																																				
RBC*	dose-dependent trend ↓																																																																									
Hb*	dose-dependent trend ↓																																																																									
WBC	dose-dependent trend ↓																																																																									
PCV	dose-dependent trend ↓																																																																									
	Week 15			Weaning																																																																						
	Low	Int.	High	Low	Int.	High																																																																				
AP																																																																										
K*		↑		↓		↑																																																																				

No human data are available on subchronic oral toxicity.

The 90 day oral toxicity of potassium diformate was studied in rats (DocIIIA6.4.1-01; FA_BPR_Ann_II_8_9_2_01: ██████████ 1998). The formic acid salt, potassium diformate ("Formi"), was used as test material as it allowed to achieve high dose levels of the formate ion with the feed due to less irritating potency than formic acid itself. The systemic bioavailability of the test substance was considerable as reflected in the increased formate plasma levels of approx. 90 to 160 mg formate/l that were regularly found after the nocturnal feed intake of the rats in the high-dose group over the entire feeding period. The formate salt failed to produce any detectable target-organ toxicity. Local irritation effects in the stomach caused a dose-related thickening of the stomach at all dose levels, which was confirmed to be squamous cell hyperplasia. After the 4 week recovery period, the squamous cell hyperplasia in the forestomach subsided and was largely reversible. No overt systemic toxicity

was observed: There was a dose-dependent decrease in bw gain in males and a decrease in bw gain in the high dose females. However, the RMS is not convinced that the slight dose-related reduction in feed intake in males is entirely responsible for the significant decrease in bw gain. There was no reduction in feed intake in females. In the recovery period, body weight development in males and females was comparable between the high dose and control groups. In addition, is the observed systemic effect (dose-dependent bw gain decrease in males and bw gain decrease at the highest dose in females) secondary to the corrosive local GI tract effect? Using a precautionary approach the LOAEL_{systemic} according to the RMS is 2100 mg formate/kg bw/d, based on decreased bw gain in males and females. The NOAEL_{systemic} is 840 mg formate/kg bw/d. LOAEL_{local} = 420 mg formate/kg bw/d and NOAEL_{local} < 420 mg formate/kg bw/d, based on histological changes in the stomach.

The pig oral feed study was conducted to assess the safety of potassium diformate at dose levels of up to five times the recommended dose in the reproducing pig and its offspring (DocIIIA6.4.1-02; FA_BPR_Ann_II_8_9_2_02: ██████████ 2004). No guideline was followed, but the test design obeyed scientific standards. The study provided additional toxicity data on a species that has a more limited metabolism capability to dispose of formate than humans. Therefore, the pig appears to be a more appropriate test species than the rat: any symptomatology possibly related to formate in pig will have significance for the extrapolation to human beings. Potassium diformate ("Formi") was used as test material at nominal concentrations of 0%, 1.2%, 3%, and 6% in the feed. Dose levels of 0, 92, 226, and 437 mg formate/kg bw/d were achieved during 114 days of gestation, dose levels of 0, 149, 359, and 760 mg/kg bw/d during lactation until day 26 post-partum. There were no mortalities or clinical signs that were treatment-related. There was no indication of visual problems in any of the animals. Haematology, clinical chemistry, urinalysis, necropsy and histopathology did not indicate any systemic toxicity. At week 15 and weaning time points, there was a trend towards lowered red blood cell counts (RBC), hemoglobin concentration (Hb), white blood cell count, packed cell volume and haemoglobin. Plasma potassium levels were dose-dependently increased at week 15 and at weaning ($p < 0.05$). There was a clear trend in sodium concentration decrease with increasing dose at the Week 15 and weaning time points, and there was also a trend for potassium concentration to increase with dose at the same time points. Likewise, there was a clear trend towards a higher pH with increased dose levels. The increased potassium uptake was considered to be related with the observed effects, rather than with formic acid. Organ weights were not recorded, but the appearance was normal. Histopathology revealed local irritating effects, as evidenced by forestomach gastritis and erosion/ulcer in approx. 30 to 60 % of the treated animals. The reproduction parameters of the pig were not changed by the treatment. The development of the piglets was also unaffected at birth and up to day 26 post-partum. The NOAEL_{systemic} is 760 mg formate/kg bw/d, the highest dose tested, based on the lack of any systemic effects. LOAEL_{local} = 149 mg formate/kg bw/d and NOAEL_{local} < 149 mg formate/kg bw/d, based on histological changes in the stomach.

Value used in Risk Assessment – Sub-chronic oral toxicity

Value/conclusion	90 day oral toxicity, potassium diformate, rats: LOAEL _{systemic} 2100 mg formate/kg bw/d, NOAEL _{systemic} 840 mg formate/kg bw/d. LOAEL _{local} 420 mg formate/kg bw/d, NOAEL _{local} < 420 mg formate/kg bw/d
-------------------------	--

	140 day oral oxicity, potassium diformate, pig: LOAEL _{systemic} >760 mg formate/kg bw/d, NOAEL _{systemic} 760 mg formate/kg bw/d, LOAEL _{local} 149 mg formate/kg bw/d, NOAEL _{local} < 149 mg formate/kg bw/d
Justification for the value/conclusion	BPD ID A6.4.1_01, FA_BPR_Ann_II_8_9_2_01: ██████████ 1998 Subchronic oral toxicity of potassium diformate in the rat has been assessed in a study according to OECD 408. NOAEL _{systemic} = 840 mg formate/kg bw/d, based on decreased bw gain at 2100 mg formate/ kg bw/d; NOAEL _{local} < 420 mg formate/kg bw/d, based on histological changes in the stomach. BPD ID A6.4_02, FA_BPR_Ann_II_8_9_2_02: ██████████ 2004 Subchronic oral toxicity of potassium diformate in the pig has been assessed in a non-guideline study following scientific standards. NOAEL _{systemic} = 760 mg formate/kg bw/d, the highest dose tested, based on the lack of any systemic effects; NOAEL _{local} < 149 mg formate/kg bw/d, based on histological changes in the stomach.

Data waiving	
Information requirement	Subchronic oral toxicity study on formic acid
Justification	Subchronic toxicity studies are available for the oral route using potassium diformate. The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01; FA_BPR_Ann_II_8_8_01: ██████████ 1997).

3.6.2 Sub-chronic dermal toxicity

No data are available on subchronic dermal toxicity.

Value used in Risk Assessment – Sub-chronic dermal toxicity	
Value/conclusion	n.a.
Justification for the value/conclusion	n.a.

Data waiving	
Information requirement	Subchronic dermal toxicity study on formic acid
Justification	Subchronic dermal toxicity studies were not conducted for reasons of animal welfare, because formic acid and potassium diformate are both corrosive to the skin. In addition, formate salts differ in their local effects on skin and presumably also in the absorption characteristics compared to the acid, and therefore subchronic studies using formate were not considered to represent an adequate alternative to formic acid testing. Moreover, only limited repeated exposure is expected because of the corrosivity to the skin and because of the measures taken to prevent skin contact with the corrosive material.

3.6.3 Sub-chronic inhalation toxicity

Summary table of inhalatory sub-chronic animal studies (usually 90-day studies)						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/ group	Test substance, form (<i>gas, vapour, dust, mist</i>) and particle size (MMAD), Actual and nominal concentration, Type of administration (<i>nose only / whole body/ head only</i>),	NOAEL, LOAEL	Results	Remarks (<i>e.g. major deviations</i>)	Reference

		Duration of exposure																																														
In accordance with OECD 413 GLP: yes Rel. 1	Rat, Fischer 344/N, m + f 10/sex	Formic acid purity 95% 0, 15, 30, 61, 122, 244 mg/m ³ (nominal) Vapour, whole body 6h/d, 5d/wk 13 wk	NOAEL _{Local} : 30 mg/m ³ LOAEL _{Local} : 61 mg/m ³ NOAEL _{Systemic} : 244 mg/m ³ (highest dose tested) LOAEL _{Systemic} : Not achieved	No clinical signs No active substance related mortality. Local effects: nasal irritation, squamous metaplasia of the respiratory epithelium, olfactory degeneration, severity minimal to mild. Respiratory epithelium squamous metaplasia: <table border="1"> <tr><td>mg/m³</td><td>0</td><td>15</td><td>30</td><td>61</td><td>122</td><td>244</td></tr> <tr><td>male</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>9</td></tr> <tr><td>female</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>6</td></tr> </table> Olfactory epithelium degeneration: minimal to mild <table border="1"> <tr><td>mg/m³</td><td>0</td><td>15</td><td>30</td><td>61</td><td>122</td><td>244</td></tr> <tr><td>male</td><td>0</td><td>0</td><td>0</td><td>1</td><td>1</td><td>9</td></tr> <tr><td>female</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>5</td></tr> </table> No systemic toxic effects: No effect on body weight (gain) in males and females. Liver weight (abs. and rel.) increased for all treated groups in males, no effects in females. Relative lung weights decreased for all treated males, absolute lung weights only decreased for the 122 and 244 mg/m ³ males. Lung weights (abs. and rel.) decreased for all treated groups in females. Decrease in lung weight was without dose-response relationship and histopathological manifestations. Haematological and clinical chemistry changes were mild and generally	mg/m ³	0	15	30	61	122	244	male	0	0	0	0	0	9	female	0	0	0	0	0	6	mg/m ³	0	15	30	61	122	244	male	0	0	0	1	1	9	female	0	0	0	0	0	5		BPD ID A6.4.3_01 FA_BPR_Ann_II_8 _9_2_03 Thompson, 1992
mg/m ³	0	15	30	61	122	244																																										
male	0	0	0	0	0	9																																										
female	0	0	0	0	0	6																																										
mg/m ³	0	15	30	61	122	244																																										
male	0	0	0	1	1	9																																										
female	0	0	0	0	0	5																																										

				<p>unremarkable, and of no biological relevance:</p> <p>Slight neutropenia in males and females of all exposure levels after week 13: Statistically significant decreases in the number of segmented neutrophils ($p < 0.01$)</p> <p>Slight leukocytosis in males and females at 64 and 128 ppm, in females also at 8 ppm, after 3 days ($p < 0.01$)</p> <p>Slight decreases in urea nitrogen (UN), albumin, globulin, total protein, and creatinine in males and females at day 3 at 64 and 128 ppm, protein parameters only statistically significant for females ($p < 0.01$). A significant decrease in UN also in the female 16- and 32-ppm groups ($p < 0.01$).</p> <p>These changes were attributed to reduced feed intake during the first exposure period according to authors.</p> <p>Increase in sorbitol dehydrogenase in males of all groups exposed to ≥ 16 ppm after 3 days ($p < 0.01$). No changes for other liver-specific indicators.</p> <p>Increase in alkaline phosphatase in males at 128 ppm after 3 days, while decreases in females at 64 and 128 ppm at the same time point, and again increases in both top-dosed sexes after 13 weeks ($p < 0.01$).</p>		
--	--	--	--	--	--	--

				<p>Decrease in creatine kinase in males from 16 to 128 ppm after 3 days ($p < 0.01$).</p> <p>Decrease in amylase in females at 64 and 128 ppm after 3 and 23 days.</p> <p>Reproductive parameters: No effects on sperm motility, density or testicular or epididymal weights, no changes in the length of the oestrous cycle.</p>																						
<p>In accordance with OECD 413 GLP: yes Rel. 1</p>	<p>Mice B6C3F₁ m + f 10/sex</p>	<p>Formic acid purity 95% 0, 15, 30, 61, 122, 244 mg/m³ (nominal) Vapour, whole body 6h/d, 5d/wk 13 wk</p>	<p>NOAEL_{Local}: 61 mg/m³ LOAEL_{Local}: 122 mg/m³ NOAEL_{Systemic}: 122 mg/m³ LOAEL_{Systemic}: 244 mg/m³</p>	<p>No clinical signs No active substance related mortality. Local effects: nasal irritation, olfactory degeneration, severity minimal but dose-related. Olfactory epithelium degeneration: minimal</p> <table border="1"> <thead> <tr> <th>mg/m³</th> <th>0</th> <th>15</th> <th>30</th> <th>61</th> <th>122</th> <th>244</th> </tr> </thead> <tbody> <tr> <td>male</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> </tr> <tr> <td>female</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> <td>5</td> </tr> </tbody> </table> <p>Systemic toxic effects: Decrease in body weight gain in males and females at 244 mg/m³ (male terminal bw = 84% of the control, female terminal bw = 80% of the control). Relative liver weight increased for the 61, 122, 244 mg/m³ groups in males, and the 244 mg/m³ group in females. Relative kidney weights were increased in females in the 61, 122, and 244 mg/m³ groups. These changes were</p>	mg/m ³	0	15	30	61	122	244	male	0	0	0	0	0	2	female	0	0	0	0	2	5	<p>BPD ID A6.4.3_01 FA_BPR_Ann_II_8_9_2_04 Thompson, 1992</p>
mg/m ³	0	15	30	61	122	244																				
male	0	0	0	0	0	2																				
female	0	0	0	0	2	5																				

				without histopathological manifestations. Reproductive parameters: No effects on sperm motility, density or testicular or epididymal weights, no changes in the length of the oestrous cycle.		
--	--	--	--	--	--	--

No human data are available on subchronic inhalation toxicity.

Subchronic 13-week inhalation studies with formic acid vapour at concentrations of 0, 15, 30, 61, 122, 244 mg/m³ were conducted in rats and mice (DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_03 and DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_04: Thompson, 1992).

In the rat, the inhalation of formic acid did not result in clinical effects. All animals survived, and no effect on the body weight was observed. Changes in haematological and clinical chemistry changes measured at 3 time points (day 3, day 23, and at 13 weeks) were few and generally unremarkable. There were no gross lesions noted at necropsy. Absolute liver weights were increased in male rats in all exposure groups and relative liver weights were increased in males exposed to 61, 122, 244 mg/m³ formic acid. Absolute and relative lung weights were decreased in females in all treated groups. In males, relative lung weights were decreased for all treatment groups, absolute lung weights were decreased for the 122 and 244 mg/m³ groups. Microscopic changes occurred in the respiratory and olfactory epithelium of the nose. Changes on the respiratory epithelium consisted of minimal squamous metaplasia in which the pseudostratified, ciliated columnar cells were replaced by a flattened, non-ciliated epithelium of approximately 2 to 5 cells in thickness. Squamous metaplasia occurred most often in the respiratory epithelium that lines the most dorsal portion of the dorsal meatus in the nose's anterior section (Level I). In the olfactory epithelium, degenerative changes were minimal to mild and generally limited to the area of the dorsal meatus in the mid-nasal section (Level II). Degeneration was characterised by a loss of the usual orderly arrangement of the pseudostratified layer of nuclei and by a slight reduction on the normal thickness of the olfactory epithelium. There was no necrosis. No evidence was seen of metaplasia of the olfactory epithelium or atrophy of the nerve fibres in the olfactory mucosa. There were no effects on measures of sperm motility, density, or testicular or epididymal weights, and no changes in the length of the estrous cycle. In conclusion, the upper respiratory tract was the major target for toxicity in rats. There was no evidence of systemic toxicity. The NOAEC_{systemic} is 244 mg formic acid/m³, the highest dose tested, based on the lack of any systemic effects. LOAEC_{local} = 61 mg formic acid/m³ and NOAEC_{local} = 30 mg formic acid/m³, based on histological changes in the nasal region.

In the mouse, the inhalation of formic acid did not result in clinical effects. There was no mortality associated with the exposure to formic acid. Body weight gain was decreased for both males and females for the 244 mg/m³ group, and for the females for the 122 mg/m³ group. Relative liver weights were increased in males and females in the 122 and 244 mg/m³ groups and relative kidney weights were increased in females in the 61, 122, and 244 mg/m³ groups. There were no gross lesions noted at necropsy. Microscopic changes were limited to the degeneration of the olfactory epithelium of the nose in mice from the 122 mg/m³ and 244 mg/m³ formic acid groups. The minimal degeneration occurred in the dorsal portion of the dorsal meatus in the anterior or mid-nasal section (Levels I and II). Degeneration was characterised by a loss of the usual orderly arrangement of the pseudostratified layer of nuclei and by a slight reduction on the normal thickness of the olfactory epithelium.

Blood analysis (haematology, clinical chemistry, urinalysis) was not documented. There were no effects on the reproductive parameters evaluated. In conclusion, also in the mouse the upper respiratory tract was the major target for toxicity. LOAEC_{systemic} is 244 mg formic acid/m³, NOAEC_{systemic} is 122 mg formic acid/m³, based on the reduced body weight gain observed at 244 mg/m³. LOAEC_{local} = 122 mg formic acid/m³ and NOAEC_{local} = 61 mg formic acid/m³, based on histological changes in the nasal region.

Effects on the respiratory and olfactory epithelium at 13 weeks consisted of squamous metaplasia (minimal, rats) and degeneration (minimal, rats and mice), respectively. Based on the findings in the 13-week studies the overall NOAEC_{local} for microscopic lesions in the rats and mice is considered 60 mg/m³.

Note : The applicant does not agree with the estimated NOAEC for local effects in rats and proposes a local NOAEC in rats of 122 mg formic acid/m³ and a LOAEC of 244 mg formic acid/m³. However, eCA BE is adhering to the NOAEC_{local, rat} of 30 mg formic acid/m³. The applicant's justification for this re-interpretation can be found in the PT3 specific BASF confidential Annex to the PT3 CAR, along with BE's clarification for refusal.

Value used in Risk Assessment – Sub-chronic inhalation toxicity	
Value/conclusion	<p>13-week inhalation toxicity, formic acid, rat: LOAEC_{systemic} not achieved, NOAEC_{systemic} 244 mg formic acid/m³ LOAEC_{local} 61 mg formic acid/m³, NOAEC_{local} 30 mg formic acid/m³</p> <p>13-week inhalation toxicity, formic acid, mouse: LOAEC_{systemic} 244 mg formic acid/m³, NOAEC_{systemic} 122 mg formic acid/m³ LOAEC_{local} 122 mg formic acid/m³, NOAEC_{local} 61 mg formic acid/m³</p> <p>overall NOAEC_{local} for microscopic lesions in the rats and mice is considered 60 mg/m³</p>
Justification for the value/conclusion	<p>DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_03 ; DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_04: Thompson, 1992 Subchronic inhalation toxicity of formic acid in the rat and mouse has been assessed in a study in accordance with OECD 413.</p> <p>The upper respiratory tract was the major target organ: minimal to mild squamous metaplasia of the respiratory epithelium and minimal degeneration of the olfactory epithelium. In addition, a decrease in body weight gain was observed at the highest dose level in mice. NOAEC_{systemic} = 122 mg formic acid/m³, based on the reduced bodyweight gain observed at 244 mg/m³ in the mouse. The overall NOAEC_{local} = 60 mg formic acid/m³, based on histopathological changes in the nasal region of both rats and mice observed at 122 mg/m³.</p>

3.6.4 Overall conclusion on sub-chronic repeated dose toxicity

Value used in the Risk Assessment – Sub-chronic repeated dose systemic toxicity	
Value	medium-term oral toxicity : Rat: NOAEL _{systemic} = 840 mg formate/kg bw/d Pig: NOAEL _{systemic} = 760 mg formate/kg bw/d Medium-term inhalation toxicity: NOAEC _{systemic} = 122 mg formic acid/m ³
Justification for the selected value	<p>The medium-term oral toxicity of formic acid, administered as potassium diformate in the feed, was studied in the rat (90 days) and the pig (140 days). Local irritation effects in the stomach caused a dose-related thickening of the stomach at all dose levels, which was confirmed to be squamous cell hyperplasia of the stomach and gastrointestinal tract, and was largely reversible. High doses may produce adverse effects, such as decrease in body weight gain (rat), which might be due to the inherent irritating potential. In the rat, the NOAEL_{systemic} = 840 mg formate/kg bw/d, based on decreased bw gain at 2100 mg formate/ kg bw/d; in the pig, the NOAEL_{systemic} = 760 mg formate/kg bw/d, the highest dose tested, based on the lack of any systemic effects.</p> <p>Medium-term inhalation toxicity was studied in rats and mice exposed to formic acid vapours for 13 weeks. The upper respiratory tract was the major target organ: minimal to mild squamous metaplasia of the respiratory epithelium and minimal degeneration of the olfactory epithelium. In addition, a decrease in body weight gain was observed at the highest dose level in mice. NOAEC_{systemic} = 122 mg formic acid/m³, based on the reduced bodyweight gain observed at 244 mg/m³ in the mouse.</p>
Classification according to CLP and DSD	None

Value/conclusion used in the Risk Assessment – Sub-chronic repeated dose local effects	
Value/conclusion	medium-term oral toxicity :

	<p>Rat: NOAEL_{local} < 420 mg formate/kg bw/d</p> <p>Pig: < 149 mg formate/kg bw/d</p> <p>Medium-term inhalation toxicity:</p> <p>overall NOAEC_{local} = 60 mg formic acid/m³</p>
Justification for the selected value/conclusion	<p>The medium-term oral toxicity of formic acid, administered as potassium diformate in the feed, was studied in the rat (90 days) and the pig (140 days). Local irritation effects in the stomach caused a dose-related thickening of the stomach at all dose levels, which was confirmed to be squamous cell hyperplasia of the stomach and gastrointestinal tract, and was largely reversible. High doses may produce adverse effects, such as decrease in body weight gain (rat), which might be due to the inherent irritating potential. In the rat, the NOAEL_{local} < 420 mg formate/kg bw/d, based on histological changes in the stomach. In the pig, the NOAEL_{local} < 149 mg formate/kg bw/d, based on histological changes in the stomach.</p> <p>Medium-term inhalation toxicity was studied in rats and mice exposed to formic acid vapours for 13 weeks. The upper respiratory tract was the major target organ: minimal to mild squamous metaplasia of the respiratory epithelium and minimal degeneration of the olfactory epithelium. In addition, a decrease in body weight gain was observed at the highest dose level in mice. The overall NOAEC_{local} = 60 mg formic acid/m³, based on histopathological changes in the nasal region of both rats and mice observed at 122 mg/m³.</p>
Classification according to CLP and DSD	None

3.7 LONG-TERM REPEATED DOSE TOXICITY

3.7.1 Long-term oral toxicity

Summary table of oral long-term animal studies						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/ group	Test substance, Dose levels, Route of exposure (<i>gavage, in diet, other</i>), Duration of exposure	NOAEL, LOAEL	Results	Remarks (<i>e.g. major deviations</i>)	Reference
Comparable to 94/40/EEC GLP: yes Rel. 1	Rat, Wistar, m + f main: 50/sex/group interim: 20/sex/group	KHCO ₂ •H ₂ CO ₂ [CAS No. 20642-05-1] purity 98-99% 0, 50, 400, 2000 mg/(kg*d) = 0, 35, 280, 1400 mg formate/kg bw/d (nominal) Oral, feed continuous, 7 d/week 104 wk (interim kill at 52 wk)	NOAEL _{Local} : as formate: 35 mg/kg bw/d LOAEL _{Local} : as formate: 280 mg/kg bw/d NOAEL _{Systemic} : as formate: 280 mg/kg bw/d LOAEL _{Systemic} : as formate: 1400 mg/kg bw/d	<u>52 weeks</u> No clinical signs No active substance related mortality BW (gain): ↓ for high dose m+f Ophthalmoscopy: no effects on the eye Haematology, clinical chemistry, urinalysis: no consistent pattern of variation, no treatment effect Organ weight: no effect Necropsy: thick stomach (high dose) Histopathology: gastric irritation, stomach: foveolar epithelial (<i>males</i> grade 1: 11, grade 2: 3 /20 high dose animals vs 0/20 controls; <i>females</i> grade 1: 10, grade 2: 1 /20 high dose)		BPD ID A6.5_01/ BPD ID A6.7.-01 FA_BPR_Ann_II_8_9_3_01 FA_BPR_Ann_II_8_11_1_02 ██████████ 2002a/b

				<p>animals vs 0/20 controls) and basal cell hyperplasia (<i>males</i> grade 1: 10, grade 2: 2 /20 high dose animals vs 0/20 controls; <i>females</i> grade 1: 10, grade 2: 2 /20 high dose animals vs 0/20 controls), salivary gland: acinar cell hypertrophy (incidence high dose males 7/20, females 3/20 vs 0/20 controls), kidney: ↓ incidence of pelvic mineralisation (high dose males 0/20vs 6/20 control, high dose females 6/20 vs 14/20 controls)</p> <p><u>104 weeks</u></p> <p>No clinical signs</p> <p>No active substance related mortality</p> <p>BW (gain): ↓ for high dose, bw gain: 27% (m), 19% (f)</p> <p>Food intake: ↓ for high dose, 3% (m), 6% (f) over 104 weeks</p> <p>Ophthalmoscopy: no effects on the eye</p> <p>Haematology, clinical chemistry, urinalysis: no consistent pattern of variation, trend to ↓ RBC at pre-terminal investigation</p> <p>Organ weight: no effect</p> <p>Necropsy: nodules, raised focus, and thick stomach (high dose)</p> <p>Histopathology: gastric irritation,</p> <p>stomach: ↑ incidence and severity of basal cell/squamous</p>		
--	--	--	--	---	--	--

				<p>cell hyperplasia at the lining ridge (mid dose <i>males</i> grade 1: 13/39, grade 2: 6/39, high dose <i>males</i> grade 1: 9/43, grade 2: 19/43, grade 3: 14/43 vs 3/42 grade 1 and 1/42 grade 2 controls; mid dose <i>females</i> grade 1: 11/36, grade 2: 1/36, high dose <i>females</i> grade 1: 7/38, grade 2: 28/38, grade 3: 3/38 vs 4/39 grade 1 controls), foveolar epithelial hyperplasia (high dose <i>males</i> grade 1: 17/43, grade 2: 23/43 vs 1/41 grade 1 and 0/42 grade 2 controls; high dose <i>females</i> grade 1: 21/38 vs 0/39 grade 1 controls), acanthosis, hyperkeratosis (high dose) salivary gland: acinar cell hypertrophy ((high dose <i>males</i> 17/43 vs 0/42 controls, high dose <i>females</i> 10/38 vs 0/39 controls)</p> <p>duodenum: hypertrophy of the Brunner's glands (high dose <i>males</i> 16/43 vs 0/42 controls, high dose <i>females</i> 8/38 vs 0/39 controls)</p> <p>kidney: ↓ incidence of pelvic mineralisation (high dose <i>males</i> 4/43 vs 28/42 controls, high dose <i>females</i> 20/38 vs 37/39 controls) and papillary mineralisation (high dose <i>females</i> 2/38 vs 8/39 controls)</p>		
--	--	--	--	--	--	--

<p>94/40/EEC GLP: yes Rel. 1</p>	<p>Mouse, CD, m + f 51/sex</p>	<p>$\text{KHCO}_2 \bullet \text{H}_2\text{CO}_2$ [CAS No. 20642-05-1] purity 98- 99%: 0, 50, 400, 2000 mg/kg bw/d = 0, 35, 280, 1400 mg formate/kg bw/d (nominal), Oral, feed, continuous, 7 d/week, 80 wk</p>	<p>NOAEL_{Local/systemic}: as formate: 280 mg/kg bw/d LOAEL_{Local/systemic}: as formate: 1400 mg/kg bw/d</p>	<p><u>80 weeks</u> Clinical signs: none related to treatment No active substance related mortality BW (gain): slightly but significantly lower in high-dose males (p<0.05). No difference between control and low and mid dose animals and for all female groups. Food intake: comparable between all groups, although with a very slight increasing trend in the high-dose males. Macroscopic investigations: no effects Haematology: No adverse effects on RBC or WBC Ophthalmoscopy: not examined clinical chemistry, urinalysis: not examined Organ weight: no data Necropsy: some evidence of treatment-related thick stomach in high-dose females, the only macroscopic finding, but not noted in males Pathology: limited signs of chronic irritation in the stomach; otherwise</p>	<p>BPD ID A6.7_02. FA_BPR_Ann_II_8_11_2_01 ██████████ 2002b</p>
--	--	---	---	---	---

				<p>unremarkable</p> <p>Histopathology:</p> <p>No increase in any tumour type, slight local irritation of the forestomach with increased incidence of hyperplasia of the limiting ridge in high-dose males.</p> <p><u>Non-neoplastic observations:</u> gastric irritation, Thick stomach seen in some animals, no dose-response relationship in males, little correlation with microscopic findings</p> <p>Incidence of findings in the stomach:</p> <table border="1"> <thead> <tr> <th rowspan="2">(mg/kg bw/d)</th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> </tr> <tr> <td>thick</td> <td>6</td> <td>3</td> <td>7</td> <td>2</td> <td>1</td> <td>2</td> <td>3</td> <td>6</td> </tr> <tr> <td>raised focus</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>Incidence of limiting ridge hyperplasia in the stomach:</p> <table border="1"> <thead> <tr> <th rowspan="2">(mg/kg bw/d)</th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>36</td> <td>40</td> <td>36</td> <td>33</td> <td>37</td> <td>34</td> <td>35</td> <td>40</td> </tr> <tr> <td>grade 1</td> <td>4</td> <td>7</td> <td>6</td> <td>13</td> <td>7</td> <td>5</td> <td>7</td> <td>7</td> </tr> <tr> <td>grade 2</td> <td>0</td> <td>0</td> <td>0</td> <td>6</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>increased incidence of grade 1 (minimal) and grade 2 (slight) in high-dose males.</p> <p>NOAEL = 280 mg formate/kg bw/d</p>	(mg/kg bw/d)	males				females				0	35	280	1400	0	35	280	1400	n	51	51	51	51	51	51	51	51	thick	6	3	7	2	1	2	3	6	raised focus	0	0	0	0	0	2	0	0	(mg/kg bw/d)	males				females				0	35	280	1400	0	35	280	1400	n	36	40	36	33	37	34	35	40	grade 1	4	7	6	13	7	5	7	7	grade 2	0	0	0	6	0	0	0	0		
(mg/kg bw/d)	males					females																																																																																								
	0	35	280	1400	0	35	280	1400																																																																																						
n	51	51	51	51	51	51	51	51																																																																																						
thick	6	3	7	2	1	2	3	6																																																																																						
raised focus	0	0	0	0	0	2	0	0																																																																																						
(mg/kg bw/d)	males				females																																																																																									
	0	35	280	1400	0	35	280	1400																																																																																						
n	36	40	36	33	37	34	35	40																																																																																						
grade 1	4	7	6	13	7	5	7	7																																																																																						
grade 2	0	0	0	6	0	0	0	0																																																																																						

				<p><u>Neoplastic observations:</u> Increased incidence of primary lung tumours in high-dose males, but not in females: bronchiolo-alveolar adenomas and carcinomas</p> <p>Bronchiolo-alveolar tumour incidence:</p> <table border="1"> <thead> <tr> <th></th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>(mg/kg bw/d)</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>51</td> <td>19</td> <td>30</td> <td>51</td> <td>51</td> <td>25</td> <td>25</td> <td>51</td> </tr> <tr> <td>m. carcinoma</td> <td>0</td> <td>2</td> <td>5</td> <td>2</td> <td>0</td> <td>3</td> <td>0</td> <td>3</td> </tr> <tr> <td>b. adenoma</td> <td>4</td> <td>7</td> <td>11</td> <td>9</td> <td>5</td> <td>6</td> <td>4</td> <td>5</td> </tr> <tr> <td>all</td> <td>4</td> <td>9</td> <td>16</td> <td>11</td> <td>5</td> <td>9</td> <td>4</td> <td>8</td> </tr> </tbody> </table> <p>Alveolar epithelial tumour statistics: numbers of tumour bearing animals and results of test for dose response</p> <table border="1"> <thead> <tr> <th></th> <th colspan="2">males</th> <th colspan="2">females</th> <th>dose response</th> </tr> <tr> <th>(mg/kg bw/d)</th> <th>0</th> <th>1400</th> <th>0</th> <th>1400</th> <th></th> </tr> </thead> <tbody> <tr> <td>fatal</td> <td>0</td> <td>1</td> <td></td> <td></td> <td>ns (m, f)</td> </tr> <tr> <td>non-fatal</td> <td>4</td> <td>10</td> <td>5</td> <td>7</td> <td>ns</td> </tr> <tr> <td>all</td> <td>4</td> <td>11</td> <td></td> <td></td> <td>0.038*</td> </tr> </tbody> </table> <p>* increasing dose response; ns = not significant</p>		males				females				(mg/kg bw/d)	0	35	280	1400	0	35	280	1400	n	51	19	30	51	51	25	25	51	m. carcinoma	0	2	5	2	0	3	0	3	b. adenoma	4	7	11	9	5	6	4	5	all	4	9	16	11	5	9	4	8		males		females		dose response	(mg/kg bw/d)	0	1400	0	1400		fatal	0	1			ns (m, f)	non-fatal	4	10	5	7	ns	all	4	11			0.038*		
	males				females																																																																																					
(mg/kg bw/d)	0	35	280	1400	0	35	280	1400																																																																																		
n	51	19	30	51	51	25	25	51																																																																																		
m. carcinoma	0	2	5	2	0	3	0	3																																																																																		
b. adenoma	4	7	11	9	5	6	4	5																																																																																		
all	4	9	16	11	5	9	4	8																																																																																		
	males		females		dose response																																																																																					
(mg/kg bw/d)	0	1400	0	1400																																																																																						
fatal	0	1			ns (m, f)																																																																																					
non-fatal	4	10	5	7	ns																																																																																					
all	4	11			0.038*																																																																																					
No guideline, but following scientific standards GLP: no	Pig, crossbred f 7 control sows, 8 sows in treated groups	KHCO ₂ •H ₂ CO ₂ [CAS No. 20642-05-1] purity 95% 0, 1.2%, 3.6% in the diet	NOAEL as formate: 301 mg/(kg*d) (highest dose tested)	No signs of maternal toxicity (clinical signs, body weight development) or toxicity to reproduction or development at any dose level.		BPD ID A6.5_02 FA_BPR_Ann_II_8_9_4_0_JNS ██████████ 2003																																																																																				

Rel. 3		0, 140, and 430 mg/kg bw/d = 0, 98, 301 mg formate/kg bw/d nominal) Oral, feed continuous, 7 d/week >300 days				
--------	--	---	--	--	--	--

No human data are available on long-term oral toxicity.

The chronic oral toxicity of formate was investigated in the rat for up to 52 weeks and the effects on the incidence and morphology of tumours following oral administration of potassium diformate ("Formi") at 0, 50, 400, and 2000 mg/kg bw/d (0, 35, 280, 1400 mg formate/kg bw/d) for 104 weeks (DocIIIA6.5.-01/ FA_BPR_Ann_II_8_9_3_01 and DocIIIA6.7.-01/ FA_BPR_Ann_II_8_11_1_02: ██████████ 2002a/b). The formate salt failed to produce any target-organ toxicity. There were no treatment related clinical signs or mortality. Local irritation effects in the stomach caused thickening of the stomach, which was confirmed histopathologically. At 52 weeks, in the high dose animals foveolar epithelial hyperplasia in the stomach was characterized by an increase in the depth of intensely eosinophilic epithelium on the surface of the fundic mucosa. Basal cell hyperplasia was restricted to the squamous epithelium of the limiting ridge. In addition, there was minor acinar cell hypertrophy in the submaxillary salivary gland of some high dose animals. In the kidney there was a lower incidence of pelvic mineralisation in high dose animals. At 104 weeks, there was an increase in the incidence and severity of basal cell/squamous cell hyperplasia at the limiting ridge in high and intermediate dose animals. In addition to the basal proliferation, there was increased acanthosis and hyperkeratosis. Foveolar epithelial hyperplasia was similar as at 52 weeks. There was acinar cell hypertrophy in the submaxillary salivary gland of high dose animals. Brunner's gland hypertrophy characterized by large acinar cells was observed in the duodenum of high dose animals. In the kidney there was a lower incidence of pelvic mineralization in high dose animals. Body weight and body weight gain was decreased for high dose animals. Ophthalmoscopy showed no effect on the eye. Haematology, clinical chemistry, urinalysis, and organ weight showed no indications for treatment-related effects. In conclusion, there was no evidence of systemic target organ toxicity, including the eyes, due to formate administration. LOAEL_{systemic/local} (52 wk) = 1400 mg formate/kg bw/d, and NOAEL_{systemic/local} (52 wk) = 280 mg formate/kg bw/d, based on reduced bw gain and gastric hyperplasia. LOAEL_{systemic} (2 y) = 1400 mg formate/kg bw/d, and NOAEL_{systemic} (2 y) = 280 mg formate/kg bw/d, based on reduced bw gain. LOAEL_{local} (2 y) = 280 mg formate/kg bw/d, and NOAEL_{local} (2 y) = 35 mg formate/kg bw/d, based on hyperplastic changes in the stomach and gastrointestinal tract.

The effects on the incidence and morphology of tumours was investigated in the **mouse** following oral administration in the feed of potassium diformate ("Formi") at 0, 50, 400, and 2000 mg/kg bw/d (0, 35, 280, 1400 mg formate/kg bw/d) for 80 weeks (DocIIIA6.7.-02, FA_BPR_Ann_II_8_11_2_01: ██████████ 2002b; see also section 3.9). The animals were examined for mortality, clinical signs of toxicity and body weight. Haematological, but no clinical-chemical parameters were evaluated. The surviving animals were subjected to necropsy, and tissue slices prepared for histopathology. There were no treatment-related clinical signs, morbidity or mortality. Body weight gain was slightly but significantly lower in high-dose males, although with a very slight trend in increased food consumption in the high-dose males. There were no treatment-related effects on the red and white blood cell counts. Local irritation effects in the stomach caused thickening of the stomach but without dose-response relationship in the males, and with little correlation with microscopic findings. There was an increased incidence of limiting ridge hyperplasia in the forestomach of high-dose males. This was characterized by a minor increase of thickness and folding of the squamous epithelium at the limiting ridge, with a slightly more basophilic basal layer. This finding was considered to indicate an adaptive change to minor local irritation by the test substance. Minor limiting ridge hyperplasia was seen in all group including controls. Increased incidences of Grade 1 (minimal) and Grade 2 (slight) were seen in high-dose males. There was no evidence of progression to neoplasia. The spectrum of neoplasia was generally consistent with that expected in mice of this strain. However, there was a higher incidence of primary lung tumours (bronchiolo-alveolar adenomas and carcinomas) in high dose males than in controls. One primary tumour of the stomach was seen in one control female. According to the authors of this study, primary lung tumours are common background tumours in mice of this strain, and the incidence in the high dose males was within the background range of the laboratory. The incidence of the control males was slightly lower than expected, and the incidences across all treated groups showed no dose-related trend. Therefore, the slight background variation seen in high dose males was not considered to be of toxicological relevance, despite the statistical significance. In conclusion, the dietary administration of potassium formate to mice at dose levels up to 1400 mg formate/kg bw/d) for 80 weeks was well tolerated without treatment-related clinical effects or mortality. Treatment-related changes were limited to high-dose males and included decreased body weight, (not significant) increased food consumption, and an increased incidence of limiting ridge hyperplasia in the forestomach. The NOAEL for local/systemic toxicity was 280 mg formate/kg bw/d. There was no evidence of a tumorigenic effect in the stomach or any other tissue. The effects observed in this study and the NOAEL and LOAEL values derived from them are supportive of the effects and NOAEL and LOAEL values described in the ██████████ 2002a study on rats.

A chronic pig study on the effects of potassium diformate on ovulation and fertility in breeding sows was made available (DocIIIA6.5.-02, FA_BPR_Ann_II_8_9_4_0_JNS; ██████████ 2003). It focused on effects on fertility and, therefore, did not provide the full range of pathological and histopathological data which would be expected to be contained in guideline studies pertaining to chronic toxicity, reproduction toxicity, or developmental toxicity. However, the study provides additional data because the metabolic capability to dispose of formate is more limited in pigs. The study met generally accepted scientific standards, is well documented and, therefore, acceptable for assessment. In this study, pigs were fed 0, 140, 430 mg potassium diformate/kg bw/d (0, 98, 301 mg formate/kg bw/d) for over 300 days. No treatment-related effects were observed for maternal toxicity (clinical signs, mortality, body weight, feed consumption), nor on ovulation, fertility, gestation parameters, number of live born piglets, piglet viability and weight gain until weaning. NOAEL_{systemic} = 301 mg formate/kg bw/d, based on lack of systemic and local toxicity at the highest dose tested.

Value used in Risk Assessment – Long-term oral toxicity	
Value/conclusion	<p>104-w oral toxicity, potassium formate, rat: LOAEL_{systemic} (2 y) = 1400 mg formate/kg bw/d, NOAEL_{systemic} (2 y) = 280 mg formate/kg bw/d LOAEL_{local} (2 y) = 280 mg formate/kg bw/d, NOAEL_{local} (2 y) = 35 mg formate/kg bw/d</p> <p>>300d oral toxicity, potassium formate, pig: NOAEL_{systemic} = 301 mg formate/kg bw/d</p>
Justification for the value/conclusion	<p>BPD ID A6.5_01, FA_BPR_Ann_II_8_9_3_01: ██████████ 2002a</p> <p>Chronic oral toxicity of potassium formate in the rat has been assessed in a study in comparable to 94/40/EEC. There was no evidence of systemic target organ toxicity, including the eyes, due to formate administration. LOAEL_{systemic} (2 y) = 1400 mg formate/kg bw/d, and NOAEL_{systemic} (2 y) = 280 mg formate/kg bw/d, based on reduced bw gain. LOAEL_{local} (2 y) = 280 mg formate/kg bw/d, and NOAEL_{local} (2 y) = 35 mg formate/kg bw/d, based on hyperplastic changes in the stomach and gastrointestinal tract.</p> <p>BPD ID A6.5.-02, FA_BPR_Ann_II_8_9_4_0_JNS; ██████████ 2003:</p> <p>A chronic pig study on the effects of potassium diformate on ovulation and fertility in breeding sows was made available. No treatment-related effects were observed for maternal toxicity (clinical signs, mortality, body weight, feed consumption), nor on ovulation, fertility, gestation parameters, number of live born piglets, piglet viability and weight gain until weaning. NOAEL_{systemic} = 301 mg formate/kg bw/d, based on lack of systemic and local toxicity at the highest dose tested.</p>

Data waiving	
Information requirement	Chronic oral toxicity study on formic acid
Justification	A chronic toxicity study is available for the oral route using potassium diformate. The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01; FA_BPR_Ann_II_8_8_01: ██████████ 1997).

3.7.2 Long-term dermal toxicity

No data are available on long-term dermal toxicity.

Value used in Risk Assessment – Long-term dermal toxicity	
Value/conclusion	n.a.
Justification for the value/conclusion	n.a.

Data waiving	
Information requirement	Long-term dermal toxicity study on formic acid
Justification	Long-term oral toxicity test provides adequate information

3.7.3 Long-term inhalation toxicity

No data are available on long-term inhalation toxicity.

Value used in Risk Assessment – Long-term inhalation toxicity	
Value/conclusion	n.a.
Justification for the value/conclusion	n.a.

Data waiving	
Information requirement	Long-term inhalation toxicity study on formic acid
Justification	Long-term oral toxicity test provides adequate information

3.7.4 Overall conclusion on long-term repeated dose toxicity

Value used in the Risk Assessment – Long-term repeated dose systemic toxicity	
Value	long-term oral toxicity : Rat: NOAEL _{systemic} = 280 mg formate/kg bw/d
Justification for the selected value	The long-term oral toxicity of formic acid, administered as potassium diformate in the feed, was studied in the rat (2-year) and the pig (300 days). In the rat, local irritation effects in the stomach caused thickening of the stomach, which was confirmed histopathologically. There was an increase in the incidence and severity of basal cell/squamous cell hyperplasia, increased acanthosis, hyperkeratosis, foveolar epithelial hyperplasia, acinar cell hypertrophy in the submaxillary salivary gland, Brunner's gland hypertrophy in the duodenum. In the high dose animals, body weight (gain) was decreased and there was a lower incidence of pelvic mineralization in the kidney. NOAEL _{systemic} = 280 mg formate/kg bw/d, based on decreased bw gain at 1400 mg/kg bw/d in the 2-year rat study.
Classification according to CLP and DSD	None

Value/conclusion used in the Risk Assessment – Long-term repeated dose local effects	
Value/conclusion	long-term oral toxicity : Rat: NOAEL _{local} = 35 mg formate/kg bw/d
Justification for the selected value/conclusion	The long-term oral toxicity of formic acid, administered as potassium diformate in the feed, was studied in the rat (2-year) and the pig (300 days). In the rat, local irritation effects in the stomach caused thickening of the stomach, which was confirmed histopathologically. There was an increase in the incidence and severity of basal cell/squamous cell hyperplasia, increased acanthosis, hyperkeratosis, foveolar epithelial hyperplasia, acinar cell hypertrophy in the submaxillary salivary gland, Brunner's gland hypertrophy in the duodenum. In the high dose animals, body weight (gain) was decreased and there was a lower incidence of pelvic mineralization in the kidney. NOAEL _{local} = 35 mg formate/kg bw/d, based on hyperplastic changes in the stomach and gastrointestinal tract at 280 mg/kg bw/d in the 2-year rat study.

Classification according to CLP and DSD	none
--	------

3.8 GENOTOXICITY

3.8.1 In vitro

Summary table of in vitro genotoxicity studies					
Method, Guideline, GLP status, Reliability	Test substance, Doses	Relevant information about the study (e.g. cell type, strains)	Results	Remarks (e.g. major deviations)	Reference
Bacterial reverse mutation test Ames, pre-incubation variant, acc.to Haworth et al., Environ. Mutagen. 5(1): 3-142, 1983 GLP: no Rel. 4	Formic acid purity 98% dissolved in water 0, 10, 33, 100, 333, 1000, 3333 µg/plate	<i>Salmonella typhimurium</i> TA97, TA98, TA100, TA1535	+S9 : - -S9 : -	Not considered as key study for concluding on in vitro mutagenicity in bacterial cells Cytotoxicity at ≥1000 µg/plate (-/+S9) Test conducted with/without S9 from hamster and rat liver Positive controls confirmed the validity of the test Publication Deviations: -missing <i>E. coli</i> or TA102 strain; -2-Aminoanthracene as sole positive control (microsomal enzymes not tested) -No pH conditions stated	BPD ID A6.6.1_01 FA_BPR_Ann_II_8_5_1_01 Zeiger et al., 1992

				-individual plate counts are not presented	
Bacterial reverse mutation test Ames, OECD 471 GLP: yes Rel. 1 Standard plate test (SPT) Pre-incubation test (PIT)	Formic acid purity 85% dissolved in water SPT 0, 33, 100, 333, 1000, 2500, 5000 µg/plate PIT 0, 10, 33, 100, 333, 1000, 2500 µg/plate	<i>Salmonella typhimurium</i> TA1537, TA98, TA100, TA1535 <i>E. coli</i> WP2 uvrA	+S9 : - -S9 : -	Not mutagenic in bacterial cells SPT: Cytotoxicity at ≥1000 µg/plate PIT: Cytotoxicity at ≥100 µg/plate Depending on strain & test conditions Test conducted with/without S9 from rat liver Positive controls confirmed the validity of the test +S9: 2-Aminoanthracene as positive control; S9 batch characterized with benzo(a)pyrene (pur. ≥96%) in TA98 & TA100 No pH conditions stated	BPD ID A6.6.1_02 FA_BPR_Ann_II_8_5_1_02 ██████████ 2022
Mammalian chromosome aberration test, OECD 473 GLP: no data Rel. 2	Formic acid 2M stock solution dissolved in water 270-1380 µg/ml 270, 360, 450, 540, 630 µg/ml (6-14mM), at increased	CHO K1 cells	+S9 : ± -S9 : ±	Not clastogenic in mammalian cells Pos. results attributed to low pH (pH 6.1 - 6.4): dose-dependent increased aberration rate. <u>1st series</u> At initial pH 6.1, without buffering: 12 mM (-S9): 15.9%	BPD ID A6.6.2_01 FA_BPR_Ann_II_8_5_2_01 Morita et al., 1990

	buffer capacity up to 1380 µg/ml (30 mM)			<p>aberrations 10mM (+S9): 20.5% aberrations</p> <p>Toxic concentration 12 – 14 mM (pH ≤ 6.0).</p> <p><u>2nd series</u> Effect of neutralization of the medium</p> <table border="1"> <thead> <tr> <th>12-14 mM:</th> <th colspan="2">% aberrations</th> </tr> <tr> <th>initial pH</th> <th>-S9</th> <th>+S9</th> </tr> </thead> <tbody> <tr> <td>6.0</td> <td>12</td> <td>33</td> </tr> <tr> <td>6.4</td> <td>4</td> <td>2</td> </tr> <tr> <td>7.2</td> <td>0</td> <td>3</td> </tr> </tbody> </table> <p><u>3rd series</u> Effect of buffer capacity</p> <p>At enhanced buffer, toxic conc. increased to 30 mM.</p> <table border="1"> <thead> <tr> <th>Dose (mM)</th> <th>initial pH</th> <th>NaHCO3 34mM</th> <th>HEPES 30mM</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>7.4</td> <td>0.75</td> <td></td> </tr> <tr> <td>20</td> <td>6.1</td> <td>0.5</td> <td></td> </tr> <tr> <td>25</td> <td>5.8</td> <td>0.5</td> <td></td> </tr> <tr> <td>27.5</td> <td>5.7</td> <td>10.5</td> <td></td> </tr> <tr> <td>30</td> <td>5.4</td> <td>toxic</td> <td></td> </tr> <tr> <td>0</td> <td>8.5</td> <td></td> <td>0</td> </tr> <tr> <td>10</td> <td>7.6</td> <td></td> <td>0.5</td> </tr> <tr> <td>20</td> <td>7.1</td> <td></td> <td>0</td> </tr> <tr> <td>25</td> <td>6.7</td> <td></td> <td>12</td> </tr> <tr> <td>30</td> <td>5.9</td> <td></td> <td>toxic</td> </tr> </tbody> </table> <p>Chromosomal aberrations: chromatid specific: chromatid gaps, breaks, exchanges</p>	12-14 mM:	% aberrations		initial pH	-S9	+S9	6.0	12	33	6.4	4	2	7.2	0	3	Dose (mM)	initial pH	NaHCO3 34mM	HEPES 30mM	0	7.4	0.75		20	6.1	0.5		25	5.8	0.5		27.5	5.7	10.5		30	5.4	toxic		0	8.5		0	10	7.6		0.5	20	7.1		0	25	6.7		12	30	5.9		toxic	
12-14 mM:	% aberrations																																																															
initial pH	-S9	+S9																																																														
6.0	12	33																																																														
6.4	4	2																																																														
7.2	0	3																																																														
Dose (mM)	initial pH	NaHCO3 34mM	HEPES 30mM																																																													
0	7.4	0.75																																																														
20	6.1	0.5																																																														
25	5.8	0.5																																																														
27.5	5.7	10.5																																																														
30	5.4	toxic																																																														
0	8.5		0																																																													
10	7.6		0.5																																																													
20	7.1		0																																																													
25	6.7		12																																																													
30	5.9		toxic																																																													

				No positive control included, but positive results at acidic pH levels, demonstrated the sensitivity of the test system Testing program included acetic and lactic acid																					
In vitro mammalian cell gene mutation test (HPRT), OECD 476; EEC 2000/32, B.12 GLP: yes Rel. 1	Formic acid 85.3% Water 14.3% 31 – 500 µg/ml <i>1st experiment</i> <u>-S9</u> : 0, 31.25, 62.5, 125, 250, 500 µg/ml <u>-S9</u> : 0, 25, 50, 100, 200, 400 µg/ml <i>2nd experiment</i> <u>-S9/-S9</u> : 0, 100, 200, 300, 400, 500 µg/ml Vehicle control: culture medium Positive controls: EMS 300 µg/ml (- S9): MCA 10 µg/ml (+S9):	CHO K1 cells	+S9 : - -S9 : -	Not mutagenic in mammalian cells There was no increase in the number of mutant colonies with or without metabolic activation compared with the vehicle control. Cytotoxicity: -S9: # colonies and cell density not reduced +S9: # colonies ↓ at 200-300 µg/ml cell density ↓ at 300-400 µg/ml (2 nd exp.) 2 experiments, 6 replicates, pH and osmolality measured Mutant frequency (per 10 ⁶ cells), corrected: <table border="1"> <thead> <tr> <th></th> <th>Vehicle</th> <th>EMS</th> <th>MCA</th> </tr> </thead> <tbody> <tr> <td>1st exp -S9</td> <td>2.96</td> <td>295.88</td> <td></td> </tr> <tr> <td>1st exp +S9</td> <td>4.05</td> <td></td> <td>242.94</td> </tr> <tr> <td>2nd exp -S9</td> <td>2.88</td> <td>302.03</td> <td></td> </tr> <tr> <td>2nd exp + S9</td> <td>3.54</td> <td></td> <td>149.02</td> </tr> </tbody> </table>		Vehicle	EMS	MCA	1 st exp -S9	2.96	295.88		1 st exp +S9	4.05		242.94	2 nd exp -S9	2.88	302.03		2 nd exp + S9	3.54		149.02	BPD ID A6.6.3_01 FA_BPR_Ann_II_8_5_3_01 ██████████ 2002
	Vehicle	EMS	MCA																						
1 st exp -S9	2.96	295.88																							
1 st exp +S9	4.05		242.94																						
2 nd exp -S9	2.88	302.03																							
2 nd exp + S9	3.54		149.02																						

Formic acid was tested together with a high number of chemicals for its potential to induce reverse **mutations** in **bacterial** strain *Salmonella typhimurium* TA97, 98, 100, 1535 at concentrations between 100 and 3333 µg/plate in the presence and absence of metabolic activation (rat,

hamster derived), using the pre-incubation variant of the Ames test according to Haworth et al., 1983 (DocIIIA6.6.1-01, FA_BPR_Ann_II_8_5_1_01; Zeiger et al., 1992). Two series of tests were performed. In case the result had been negative or equivocal in the first run, the S9-mix concentration was enhanced from 10 (first test) to 30%. A negative solvent control (water) and appropriate positive controls were carried along. Formic acid did not induce reverse mutations in *S. typhimurium* at concentrations between 100 and 3333 µg/plate in the presence and absence of metabolic activation (rat and hamster source), where the positive controls led to a clear increase in revertant colonies. Slight cytotoxicity was reported at 3333 µg/plate, in isolated cases at 1000 µg/plate. The authors concluded that formic acid was to be considered not mutagenic in bacterial cells. The following methodological deficiencies were identified for this study: only four strains of bacteria were used; neither *E. coli* WP2 uvrA, *E. coli* WP2 uvrA (pKM101), or *S. typhimurium* TA102 were utilized; 2-Aminoanthracene was used as the sole positive control in the presence of S9-mix without further characterization of the S9 batch with a mutagen that requires metabolic activation by microsomal enzymes; pH conditions were not stated, and no individual plate counts were presented. Therefore this study could not be considered as key study for concluding on *in vitro* mutagenicity in bacterial cells.

A recent GLP-compliant study report in line with OECD 471 has been made available (DocIIIA6.6.1-02, FA_BPR_Ann_II_8_5_1_02; ██████████ 2022). Using both the standard plate (SPT) and pre-incubation (PIT) assay variant, formic acid was tested up to a dose of 5000 (SPT) and 2500 µg/plate (PIT), in the presence and absence of metabolic activation (rat derived). Formic acid did not lead to a relevant increase in the number of revertant colonies in the two assay variants, with or without S9 mix. Cytotoxicity was occasionally observed depending on the strain and test conditions at and above 1000 µg/plate (SPT) or at and above 100 µg/plate (PIT). All required bacterial tester strains were accounted for. The number of revertant colonies in the negative controls, with and without S9 mix, were within the range of the respective historical control data of each tester strain. Suitable positive controls were selected per strain which induced an appropriate mutagenic response, in line with historical control data. As positive control in the presence of metabolic activation, 2-aminoanthracene was used for all tester strains. The S9 batch was characterized with benzo(a)pyrene (pur. ≥96%) in TA98 and TA100 strains. pH conditions were not stated. However, as cytotoxicity was observed mainly at top dose levels, the impact of the pH value on the reliability of the study can be considered minor. Moreover, the selected top dose is in compliance with OECD TG 471. The study can be accepted as a key study. It can be concluded that formic acid is not mutagenic in bacterial cells.

Formic acid was tested for its potential to induce **chromosomal aberrations** in **mammalian** cells, CHO K1 cells (DocIIIA6.6.2-01, FA_BPR_Ann_II_8_5_2_01; Morita et al., 1990). A positive control was missing. The study was focused upon the influence of the pH of the medium, comprising various operations for shifting the pH as desired. Acetic and lactic acid were also tested in this study. In a first series, incubation was carried out in a standard medium without pH regulation. In a second series, the initial pH of the medium was adjusted to pH 6.0 with 14mM or 12 mM formic acid. These media were then neutralised to pH 6.4, and a second group to pH 7.2. In a third series, the effect of an increased buffer capacity was examined with 2 different buffer systems. All experiments were conducted with and without metabolic activation. There was a dose-related response in the chromosomal aberration rate. In the absence of additional buffer the effective doses of formic acid were 10-12 mM. Under the condition of enhanced buffer capacity, the effective doses increased. Depending on the buffer used, aberrant cells were seen at 25 or 27.5 mM and above. But there was no clastogenic activity at 20 or 25 mM formic acid. At 30 mM the formic acid was cytotoxic irrespective of the buffer system. Mainly chromatid-specific lesions (chromatid-type gaps and breaks with/without S9, chromatid exchanges with S9) were induced, also several-fold per cell at the high doses (= lower pH or buffer capacity). This also applied to

acetic and lactic acid, both included in the testing programme. It was concluded that formic acid is not itself clastogenic to these cells but that the acidic conditions of the medium were responsible for the chromosome aberrations observed (false –positive responses).

Formic acid was tested for its ability to induce gene **mutations** at the HPRT locus in **mammalian cells**, CHO K1 cells (DocIIIA6.6.3.-01, FA_BPR_Ann_II_8_5_3_01; ██████████ 2002). Two independent experiments were carried out with and without metabolic activation, including a vehicle and appropriate positive controls. The negative controls gave mutant frequencies within the range expected, and the positive controls led to the expected increase in the frequencies of forward mutations. Formic acid did not cause any increase in the mutant frequencies with or without S9-mix compared to the vehicle control. Cytotoxicity was observed in the presence of metabolic activation. Without S9, the number of colonies and cell density were not reduced at 500 µg/ml. Formic acid is not mutagenic in mammalian cells.

Conclusion used in Risk Assessment – Genotoxicity <i>in vitro</i>	
Conclusion	<i>In vitro</i> , formic acid was not mutagenic in bacterial and mammalian cells.
Justification for the conclusion	<p>BPD ID A6.6.1_02, FA_BPR_Ann_II_8_5_1_02: ██████████ 2022 BPD ID A6.6.2_01, FA_BPR_Ann_II_8_5_2_01 : Morita et al., 1990 BPD ID A6.6.3_01, FA_BPR_Ann_II_8_5_3_01: ██████████, 2002</p> <p><i>In vitro</i> genotoxicity of formic acid has been assessed in appropriate studies.</p> <p>There was no increase in the number of mutant colonies observed with or with metabolic activation. Cytotoxicity occurred at/above 1000 µg/plate (Ames-SPT) or at/above 100 µg/plate (Ames-PIT) or 300 µg/ml, respectively. In mammalian CHO cells, formic acid produced a dose-related response in the chromosomal aberration rate at an initial pH 6.1-6.4 without buffering, with an effective dose of 10-12 mM. Under the condition of enhanced buffer capacity, the effective doses increased. Depending on the buffer used, aberrant cells were seen at 25 or 27.5 mM and above. But there was no clastogenic activity at 20 or 25 mM formic acid. At 30 mM the formic acid was cytotoxic irrespective of the buffer system. Mainly chromatid-specific lesions were induced. It was concluded that formic acid is not itself clastogenic to these cells but that the acidic conditions of the medium were responsible for the chromosome aberrations.</p>

3.8.2 In vivo

No in vivo data on genotoxicity are available.

Conclusion used in Risk Assessment – Genotoxicity *in vivo*

Conclusion	n.a.
Justification for the conclusion	n.a.

Data waiving	
Information requirement	In vivo genotoxicity testing for formic acid
Justification	Formic acid gave negative results in the <i>in vitro</i> gene mutation study in bacteria, the <i>in vitro</i> cytogenicity study in mammalian cells, and <i>in vitro</i> gene mutation assay in mammalian cells. Therefore, no <i>in vivo</i> genotoxicity studies (bone marrow assay for chromosomal damage or a micronucleus test) are required.

3.8.3 Overall conclusion on genotoxicity

Conclusion used in the Risk Assessment – Genotoxicity	
Conclusion	Formic acid has no genotoxic potential.
Justification for the conclusion	<i>In vitro</i> , formic acid was not mutagenic in bacterial and mammalian cells. There was no increase in the number of mutant colonies observed with or with metabolic activation. In mammalian CHO cells, formic acid is not itself clastogenic but the acidic conditions of the medium were responsible for chromosome aberrations. <i>In vivo</i> data are not available and not required. The overall evaluation of the data leads to the conclusion that formic acid has no genotoxic potential itself.
Classification according to CLP and DSD	none

3.9 CARCINOGENICITY

Summary table of carcinogenicity studies in animals						
Method, Guideline, GLP status, Realibility	Species, Strain, Sex, No/ group	Test substance, Dose levels, Route of exposure, Duration of exposure	NOAEL, LOAEL	Results (Please indicate any results that might suggest carcinogenic effects, as well as other toxic effects)	Re-remarks (e.g. major deviations)	Reference
Comparable to 94/40/EEC GLP: yes Rel. 1	Rat, Wistar, m + f 50/sex	KHCO ₂ •H ₂ CO ₂ [CAS No. 20642-05-1] purity 98-99%: 0, 50, 400, 2000 mg/kg bw/d = 0, 35, 280, 1400 mg formate/kg bw/d (nominal), Oral, feed, continuous, 7 d/week, 104 wk	NOAEL _{Local} : as formate: 35 mg/kg bw/d LOAEL _{Local} : as formate: 280 mg/kg bw/d NOAEL _{Systemic} : as formate: 280 mg/kg bw/d LOAEL _{Systemic} : as formate: 1400 mg/kg bw/d	No increase in any tumour type, local irritation in the gastro-intestinal tract associated with hyperplasia. <u>Non-neoplastic observations:</u> gastric irritation, stomach: ↑ incidence and severity of basal cell/squamous cell hyperplasia at the lining ridge (mid dose <i>males</i> grade 1: 13/39, grade 2: 6/39, high dose males grade 1: 9/43, grade 2: 19/43, grade 3: 14/43 vs 3/42 grade 1 and 1/42 grade 2 controls; mid dose <i>females</i> grade 1: 11/36, grade 2: 1/36, high dose females grade 1: 7/38, grade 2: 28/38, grade 3: 3/38 vs 4/39 grade 1 controls), foveolar epithelial hyperplasia (high dose <i>males</i> grade 1: 17/43, grade 2: 23/43 vs 1/41 grade 1 and 0/42 grade 2 controls; high dose		BPD ID A6.5_01/ BPD ID A6.7.-01 FA_BPR_Ann_II_8_9_3_01 FA_BPR_Ann_II_8_11_1_02 ██████████ 2002a/b (see also 3.7.1)

				<p><i>females</i> grade 1: 21/38 vs 0/39 grade 1 controls), acanthosis, hyperkeratosis (high dose) salivary gland: acinar cell hypertrophy ((high dose males 17/43 vs 0/42 controls, high dose females 10/38 vs 0/39 controls) duodenum: hypertrophy of the Brunner's glands (high dose males 16/43 vs 0/42 controls, high dose females 8/38 vs 0/39 controls) kidney: ↓ incidence of pelvic mineralisation (high dose males 4/43 vs 28/42 controls, high dose females 20/38 vs 37/39 controls) and papillary mineralisation (high dose females 2/38 vs 8/39 controls)</p> <p><u>Neoplastic observations:</u> Reduced incidence of fibroadenoma in the mammary gland of high dose females</p>		
94/40/EEC GLP: yes Rel. 1	Mouse, CD, m + f 51/sex	KHCO ₂ •H ₂ CO ₂ [CAS No. 20642-05-1] purity 98-99%: 0, 50, 400, 2000 mg/kg bw/d = 0, 35, 280,	NOAEL _{Local/systemic} : as formate: 280 mg/kg bw/d LOAEL _{Local/systemic} : as formate:	No increase in any tumour type, slight local irritation of the forestomach with increased incidence of hyperplasia of the limiting ridge in high-dose males. <u>Non-neoplastic observations:</u> gastric irritation,		BPD ID A6.7_02. FA_BPR_Ann_II_8_11_2_01 [REDACTED] 2002b

		1400 mg formate/kg bw/d (nominal), Oral, feed, continuous, 7 d/week, 80 wk	1400 mg/kg bw/d	<p>Thick stomach seen in some animals, no dose-response relationship in males, little correlation with microscopic findings</p> <p>Incidence of findings in the stomach:</p> <table border="1"> <thead> <tr> <th rowspan="2">(mg/kg bw/d)</th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> <td>51</td> </tr> <tr> <td>thick</td> <td>6</td> <td>3</td> <td>7</td> <td>2</td> <td>1</td> <td>2</td> <td>3</td> <td>6</td> </tr> <tr> <td>raised focus</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> <td>2</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>Incidence of limiting ridge hyperplasia in the stomach:</p> <table border="1"> <thead> <tr> <th rowspan="2">(mg/kg bw/d)</th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>36</td> <td>40</td> <td>36</td> <td>33</td> <td>37</td> <td>34</td> <td>35</td> <td>40</td> </tr> <tr> <td>grade 1</td> <td>4</td> <td>7</td> <td>6</td> <td>13</td> <td>7</td> <td>5</td> <td>7</td> <td>7</td> </tr> <tr> <td>grade 2</td> <td>0</td> <td>0</td> <td>0</td> <td>6</td> <td>0</td> <td>0</td> <td>0</td> <td>0</td> </tr> </tbody> </table> <p>increased incidence of grade 1 (minimal) and grade 2 (slight) in high-dose males.</p> <p>NOAEL = 280 mg formate/kg bw/d</p> <p><u>Neoplastic observations:</u></p> <p>Increased incidence of primary lung tumours in high-dose males, but not in females: bronchiolo-alveolar adenomas and carcinomas</p> <p>Bronchiolo-alveolar tumour incidence:</p> <table border="1"> <thead> <tr> <th rowspan="2">(mg/kg bw/d)</th> <th colspan="4">males</th> <th colspan="4">females</th> </tr> <tr> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> <th>0</th> <th>35</th> <th>280</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>n</td> <td>51</td> <td>19</td> <td>30</td> <td>51</td> <td>51</td> <td>25</td> <td>25</td> <td>51</td> </tr> </tbody> </table>	(mg/kg bw/d)	males				females				0	35	280	1400	0	35	280	1400	n	51	51	51	51	51	51	51	51	thick	6	3	7	2	1	2	3	6	raised focus	0	0	0	0	0	2	0	0	(mg/kg bw/d)	males				females				0	35	280	1400	0	35	280	1400	n	36	40	36	33	37	34	35	40	grade 1	4	7	6	13	7	5	7	7	grade 2	0	0	0	6	0	0	0	0	(mg/kg bw/d)	males				females				0	35	280	1400	0	35	280	1400	n	51	19	30	51	51	25	25	51		
(mg/kg bw/d)	males					females																																																																																																																		
	0	35	280	1400	0	35	280	1400																																																																																																																
n	51	51	51	51	51	51	51	51																																																																																																																
thick	6	3	7	2	1	2	3	6																																																																																																																
raised focus	0	0	0	0	0	2	0	0																																																																																																																
(mg/kg bw/d)	males				females																																																																																																																			
	0	35	280	1400	0	35	280	1400																																																																																																																
n	36	40	36	33	37	34	35	40																																																																																																																
grade 1	4	7	6	13	7	5	7	7																																																																																																																
grade 2	0	0	0	6	0	0	0	0																																																																																																																
(mg/kg bw/d)	males				females																																																																																																																			
	0	35	280	1400	0	35	280	1400																																																																																																																
n	51	19	30	51	51	25	25	51																																																																																																																

				<table> <tr> <td>m. carcinoma</td> <td>0</td> <td>2</td> <td>5</td> <td>2</td> <td>0</td> <td>3</td> <td>0</td> <td>3</td> </tr> <tr> <td>b. adenoma</td> <td>4</td> <td>7</td> <td>11</td> <td>9</td> <td>5</td> <td>6</td> <td>4</td> <td>5</td> </tr> <tr> <td>all</td> <td>4</td> <td>9</td> <td>16</td> <td>11</td> <td>5</td> <td>9</td> <td>4</td> <td>8</td> </tr> </table> <p>Alveolar epithelial tumour statistics: numbers of tumour bearing animals and results of test for dose response</p> <table> <thead> <tr> <th rowspan="2"></th> <th colspan="2">males</th> <th colspan="2">females</th> <th rowspan="2">dose response</th> </tr> <tr> <th>(mg/kg bw/d)</th> <th>0</th> <th>1400</th> <th>0</th> <th>1400</th> </tr> </thead> <tbody> <tr> <td>fatal</td> <td></td> <td>0</td> <td>1</td> <td></td> <td>ns (m, f)</td> </tr> <tr> <td>non-fatal</td> <td></td> <td>4</td> <td>10</td> <td>5</td> <td>7</td> <td>ns</td> </tr> <tr> <td>all</td> <td></td> <td>4</td> <td>11</td> <td></td> <td></td> <td>0.038*</td> </tr> </tbody> </table> <p>* increasing dose response; ns = not significant</p>	m. carcinoma	0	2	5	2	0	3	0	3	b. adenoma	4	7	11	9	5	6	4	5	all	4	9	16	11	5	9	4	8		males		females		dose response	(mg/kg bw/d)	0	1400	0	1400	fatal		0	1		ns (m, f)	non-fatal		4	10	5	7	ns	all		4	11			0.038*	
m. carcinoma	0	2	5	2	0	3	0	3																																																							
b. adenoma	4	7	11	9	5	6	4	5																																																							
all	4	9	16	11	5	9	4	8																																																							
	males		females		dose response																																																										
	(mg/kg bw/d)	0	1400	0		1400																																																									
fatal		0	1		ns (m, f)																																																										
non-fatal		4	10	5	7	ns																																																									
all		4	11			0.038*																																																									

No human data are available on carcinogenicity.

The carcinogenic potential of formic acid was investigated in rats and mice. The formic acid salt, potassium diformate ("Formi"), was used as test material as it allowed to achieve high dose levels of the formate ion with the feed due to less irritating potency than formic acid itself.

The effects on the incidence and morphology of tumours was investigated in the **rat** following oral administration in the feed of potassium diformate ("Formi") at 0, 50, 400, and 2000 mg/kg bw/d (0, 35, 280, 1400 mg formate/kg bw/d) for 104 weeks (DocIIIA6.5.-01/ FA_BPR_Ann_II_8_9_3_01 and DocIIIA6.7.-01/ FA_BPR_Ann_II_8_11_1_02: ██████████ 2002a/b). Other parameters than non-neoplastic and neoplastic lesions are discussed in detail in section 3.7.1. That the systemic bioavailability of the test substance was considerable was reflected in the increased formate plasma levels of approx. 90 to 160 mg formate/l that were regularly found after the nocturnal feed intake of the rats in the high-dose group over the entire feeding period (see DOC-IIIA6.4.1_01/ FA_BPR_Ann_II_8_9_2_01, section 3.6.1.).

Non-neoplastic treatment-related changes were observed in the stomach, duodenum, salivary gland and kidney. In the stomach of high dose animals, there were treatment-related increased incidences of nodules, raised focus and thick stomach when compared with controls. These correlated with microscopic findings. A decrease in subcutis masses was noted in high-dose females. Compared to controls, findings in the stomach included: (1) increased incidence and severity of basal cell/squamous cell hyperplasia at the limiting ridge in mid and high dose animals. This correlated with the macroscopic findings described above; (2) acanthosis, hyperkeratosis, formation of variably sized and shaped rete pegs and papillae; associated with minor inflammatory cell infiltration in lamina propria and submucosa; (3) foveolar epithelial hyperplasia in high dose animals; (4) mild inflammatory lesions in the glandular stomach of high dose animals. The NOAEL was 35 mg/kg bw/d. Acinar cell hypertrophy of the salivary gland was similar to that observed in the interim-kill animals (52 weeks, see 3.7.1). Brunner's gland hypertrophy

characterised by large acinar cells was noted in the duodenum of high-dose animals. In the kidney, there was a lower incidence of pelvic and papillary mineralisation and of pyelitis in high-dose groups. In females, there was a decrease in acinar hyperplasia in the mammary gland, decrease in neuropathy in the sciatic nerve, cardiomyopathy in the heart and cysts in the ovary. Notably in high-dose males, there was a decrease in hepatocyte vacuolisation and of eosinophilic and basophilic foci.

The spectrum of **neoplasia** was consistent with that expected in rats of this strain. A reduced incidence of fibroadenoma in the mammary gland was noted in the high dose females. There were no tumours of unusual nature or incidence indicative of specific target organ carcinogenicity on the stomach or any other tissue.

The effects on the incidence and morphology of tumours was investigated in the **mouse** following oral administration in the feed of potassium diformate ("Formi") at 0, 50, 400, and 2000 mg/kg bw/d (0, 35, 280, 1400 mg formate/kg bw/d) for 80 weeks (DocIIIA6.7.-02, FA_BPR_Ann_II_8_11_2_01: ██████████ 2002b). The animals were examined for mortality, clinical signs of toxicity and body weight. Haematological, but no clinical-chemical parameters were evaluated. The surviving animals were subjected to necropsy, and tissue slices prepared for histopathology. There were no treatment-related clinical signs, morbidity or mortality. Body weight gain was slightly but significantly lower in high-dose males, although with a very slight trend in increased food consumption in the high-dose males. There were no treatment-related effects on the red and white blood cell counts. Local irritation effects in the stomach caused thickening of the stomach but without dose-response relationship in the males, and with little correlation with microscopic findings. There was an increased incidence of limiting ridge hyperplasia in the forestomach of high-dose males. This was characterized by a minor increase of thickness and folding of the squamous epithelium at the limiting ridge, with a slightly more basophilic basal layer. This finding was considered to indicate an adaptive change to minor local irritation by the test substance. Minor limiting ridge hyperplasia was seen in all group including controls. Increased incidences of Grade 1 (minimal) and Grade 2 (slight) were seen in high-dose males. There was no evidence of progression to neoplasia. The spectrum of neoplasia was generally consistent with that expected in mice of this strain. However, there was a higher incidence of primary lung tumours (bronchiolo-alveolar adenomas and carcinomas) in high dose males than in controls. One primary tumour of the stomach was seen in one control female. According to the authors of this study, primary lung tumours are common background tumours in mice of this strain, and the incidence in the high dose males was within the background range of the laboratory. The incidence of the control males was slightly lower than expected, and the incidences across all treated groups showed no dose-related trend. Therefore, the slight background variation seen in high dose males was not considered to be of toxicological relevance, despite the statistical significance. In conclusion, the dietary administration of potassium formate to mice at dose levels up to 1400 mg formate/kg bw/d) for 80 weeks was well tolerated without treatment-related clinical effects or mortality. Treatment-related changes were limited to high-dose males and included decreased body weight, (not significant) increased food consumption, and an increased incidence of limiting ridge hyperplasia in the forestomach. The NOAEL for local/systemic toxicity was 280 mg formate/kg bw/d. There was no evidence of a tumorigenic effect in the stomach or any other tissue.

Conclusion used in Risk Assessment – Carcinogenicity

Value/conclusion	There is no carcinogenic potential in rats and mice fed potassium diformate.
Justification for the value/conclusion	The effects on the incidence and morphology of tumours were investigated in rats and mice following oral administration in the feed of potassium diformate (0, 35, 280, 1400 mg formate/kg bw/d). Gastric irritation was observed in the rat and the mouse. However, non-neoplastic lesions were more pronounced in the rat than the mouse. In the rat, non-neoplastic treatment-related changes were observed in the stomach, duodenum, salivary gland and kidney, in the stomach with a clear correlation with stomach thickening. In the mouse non-neoplastic changes were observed in the stomach, low-grade limiting ridge hyperplasia in the forestomach, but with little correlation with the thickening of the stomach. There was no evidence of progression to neoplasia. NOAEL rat = 35 mg formate/kg bw/d, NOAEL mouse = 280 mg formate/kg bw/d, based on gastric hyperplasia. There was no evidence of a tumorigenic effect in the stomach or any other tissue. However, in the mouse there was a higher incidence of primary lung tumours, bronchiolo-alveolar adenomas and carcinomas, in the 1400 mg formate /kg bw/d males. Although there was a background variation, the incidence of the high dose group was within the historical range for this mouse strain in the test laboratory. This was not considered to be of toxicological relevance. In conclusion, the studies provided evidence that there was no cancerogenic potential in rats and mice fed potassium diformate.
Classification according to CLP and DSD	none

Data waiving	
Information requirement	Carcinogenicity testing of formic acid
Justification	A carcinogenicity study is available using potassium diformate. The use of potassium diformate is justified because it is transformed into formic acid (DocIIIA6.2-01; FA_BPR_Ann_II_8_8_01: ██████████ 1997).

3.10 REPRODUCTIVE TOXICITY

3.10.1 Developmental toxicity

Summary table of animal studies on adverse effects on development						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/ group	Test substance Dose levels, Duration of exposure	NOAELs, LOAELs (also for maternal effects)	Results	Remarks (e.g. major deviations)	Reference
OECD 414 GLP: yes Rel. 1	Rat Wistar female 25/group	sodium formate [CAS 141-53-7] purity >99% 0, 59, 236, 945 mg/kg bw/d = 0, 40, 160, 640 mg formate/kg bw/d Oral, gavage Exposure period day 6-19 p.c.	NO(A)EL teratogenicity embryotoxicity 945 mg/kg bw/d = 640 mg formate/kg bw/d LO(A)EL teratogenicity embryotoxicity >945 mg/kg bw/d = >640 mg formate/kg bw/d	<u>Dams:</u> no maternal systemic toxicity reached <u>Foetuses:</u> no influence on gestation parameters no evidence of teratogenetic or embryotoxic effects <u>Morphological effects:</u> - External malformation (anophthalmia of the left eye): 1/213 high dose foetuses in 1/24 litters - Skeletal malformation (misshapen humerus): 1/213 control foetuses in 1/24 litters - External variations: none - Soft tissue variations (dilated renal pelvis with/without dilated ureters): no relation to dosing		BPD ID A6.8.1_01 FA_BPR_Ann_II_8_10_3_01 [REDACTED] 2005

			<p>NO(A)EL maternal 945 mg/kg bw/d = 640 mg formate/ kg bw/d</p> <p>LO(A)EL maternal >945 mg/kg bw/d = >640 mg formate/ kg bw/d</p>	<table border="1"> <thead> <tr> <th>mg/kgbw</th> <th>0</th> <th>40</th> <th>160</th> <th>640</th> </tr> </thead> <tbody> <tr> <td>%</td> <td></td> <td>5.0</td> <td>3.8</td> <td>6.1</td> <td>1.9</td> </tr> <tr> <td>tot #</td> <td></td> <td>5</td> <td>4</td> <td>5</td> <td>2</td> </tr> </tbody> </table> <p>- Skeletal variations: broad range in all groups, no relation to dosing</p>	mg/kgbw	0	40	160	640	%		5.0	3.8	6.1	1.9	tot #		5	4	5	2																				
mg/kgbw	0	40	160	640																																					
%		5.0	3.8	6.1	1.9																																				
tot #		5	4	5	2																																				
OECD 414 GLP: yes Rel. 1	Rabbit Himalayan female 25/group	<p>sodium formate [CAS 141-53-7] purity 100% 0; 100; 300; 1000 mg/kg bw/d =0, 68, 203, 677 mg formate/kg bw/d</p> <p>Oral, gavage Exposure period day 6– 28 p.i.</p>	<p>NO(A)EL teratogenicity embryotoxicity 1000 mg/kg bw/d = ~670 mg formate/ kg bw/d NO(A)EL maternal 1000 mg/kg bw/d = ~670 mg formate/ kg bw/d</p>	<p><u>Dams:</u> no maternal systemic toxicity reached</p> <p><u>Foetuses:</u> no influence on gestation parameters no evidence of teratogenetic or embryotoxic effects</p> <p><i>Morphological effects:</i> - external, soft tissue, skeletal malformations:</p> <table border="1"> <thead> <tr> <th>mg/kgbw</th> <th>0</th> <th>68</th> <th>203</th> <th>677</th> </tr> </thead> <tbody> <tr> <td>litter</td> <td></td> <td>24</td> <td>23</td> <td>22</td> <td>23</td> </tr> <tr> <td>foetuses</td> <td></td> <td>163</td> <td>169</td> <td>137</td> <td>139</td> </tr> <tr> <td>foetal incidence</td> <td></td> <td>5</td> <td>4</td> <td>5</td> <td>9</td> </tr> <tr> <td>litter incidence</td> <td></td> <td>5</td> <td>4</td> <td>4</td> <td>9</td> </tr> <tr> <td>affected</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	mg/kgbw	0	68	203	677	litter		24	23	22	23	foetuses		163	169	137	139	foetal incidence		5	4	5	9	litter incidence		5	4	4	9	affected							<p>BPD ID A6.8.1_02 FA_BPR_Ann_II_8_10_1_01 ██████████ 2008a</p>
mg/kgbw	0	68	203	677																																					
litter		24	23	22	23																																				
foetuses		163	169	137	139																																				
foetal incidence		5	4	5	9																																				
litter incidence		5	4	4	9																																				
affected																																									

				foet/litter 3.8 2.6 3.1 6.7		
				- external, soft tissue, skeletal variations:		
				mg/kgbw 0 68 203 677		
				litter 24 23 22 23		
				foetuses 163 169 137 139		
				foetal incidence 92 116 90 93		
				litter incidence 24 22 21 22		
				affected foet/litter 58.0 66.1 67.2 66.6		

No human data are available on adverse effects on development.

The potential teratogenicity of formic acid was studied in rats and rabbits.

In **rats**, teratogenicity was studied at dose levels of 0, 40, 160, 640 mg formate/kg bw/d administered by oral gavage as sodium formate from day 6 to day 19 post coitum (DocIIIA6.8.1.-01, FA_BPR_Ann_II_8_10_3_01; ██████████, 2005). Sodium formate was applied to avoid unspecific maternal toxic effects through the corrosive action of formic acid, while its potential bioavailability was expected to be equal to that of formic acid itself. No maternal toxicity was observed at any dose level. There were no treatment-related clinical signs or mortality observed, nor changes seen for body weight and food consumption in the dams. Intrauterine growth and survival were unaffected up to and including the highest dose level. The type and incidence of malformations and developmental variations did not indicate a treatment-related finding and were considered to be of spontaneous origin. Therefore no developmental toxicity and teratogenicity was observed up to and including the highest dose level tested i.e. 640 mg formate/kg bw/d. $NOAEL_{maternal} = 640$ mg formate/kg bw/d, $NOAEL_{developmental, teratogenicity} = 640$ mg formate/kg bw/d.

In **rabbits**, teratogenicity was studied at dose levels of 0, 68, 203, 677 mg formate/kg bw/d administered by oral gavage of sodium formate from day 6 to day 28 post insemination (DocIII6.8.1.-02, FA_BPR_Ann_II_8_10_1_01; ██████████ 2008a). No treatment-related effects were observed in the dams concerning mortality, clinical signs, food consumption, (corrected) body weight (gain), uterus weight, and necropsy findings. With regard to reproduction, no dose-related effects were observed including conception rate, mean number of corpora lutea, total implantations, pre-and postimplantation losses, resorption, live foetuses, and foetal sex ratio. Marginally, but not statistically significant lower foetal body weights were observed at the highest dose tested. Examination of the foetuses revealed external, soft tissue and skeletal malformations in all test groups including the control. They did neither show a consistent pattern since a number of morphological structures of different ontogenic origin were affected nor a clear dose-response relationship. Findings appeared at incidences which were generally similar

to historical control data. One external (paw hyperflexion), three soft tissue (absent lobus inferior medialis, dilated cerebral ventricle and malpositioned carotid branch), and a broad range of skeletal variations (e.g. incomplete ossifications of different bony structures) occurred in all test groups including the control. There was no relation seen to dosing, and a comparable frequency was seen in the historical control data. Therefore no maternal and developmental toxicity and teratogenicity was observed up to and including the highest dose level tested i.e. 670 mg formate/kg bw/d. NOAEL_{maternal} = 670 mg formate/kg bw/d, NOAEL_{developmental, teratogenicity} = 670 mg formate/kg bw/d.

Conclusion used in Risk Assessment – Effects on development	
Value/conclusion	No developmental toxicity and teratogenicity was observed for formate in rats and rabbits. Rats: NOAEL _{maternal, developmental, teratogenicity} = 640 mg formate/kg bw/d Rabbits: NOAEL _{maternal, developmental, teratogenicity} = 670 mg formate/kg bw/d
Justification for the value/conclusion	In rats, the type and incidence of malformations and developmental variations did not indicate treatment-related findings. In rabbits, no maternal and developmental toxicity and teratogenicity was observed up to and including the highest dose level tested.

Data waiving	
Information requirement	Adverse effects of formic acid on development
Justification	Sodium formate was applied to avoid unspecific maternal toxic effects through the corrosive action of formic acid.

3.10.2 Fertility

Summary table of animal studies on adverse effects on fertility						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/ group	Test substance Dose levels, Duration of exposure	NOAELs, LOAELs	Results	Remarks (e.g. major deviations)	Reference

OECD 416 GLP: yes Rel. 1	Rat Wistar, m/f 25/group	sodium formate [CAS 141-53- 7] purity 100% 0, 100, 300, 1000 =0, 68, 203, 677 mg formate/kg bw/d continuous, 7 d/week, exposure period: Before mating: at least 75 days Duration of exposure in general: from beginning of the study until sacrifice of parent F1, F2 generation	NOAEL _{sys} 200 mg formate/kg bw/d For F0 and F1 parental rats NOAEL _{fertility, reprod performance} 670 mg formate/kg bw/d For F0 and F1 parental rats NOAEL _{developmental} 670 mg formate/kg bw/d For F1 and F2 progeny	<u>Parental F1 males:</u> ↓ food consumption during 7/15 study weeks (↓ 5-9%) ↓ bw (up to 6%) from week 9 till end of study ↓ bw gain (up to 34%) , average bw gain ↓ 9% <u>F1, F2 generation pups:</u> No adverse effects not reprotoxic, not developmental toxic	BPD ID A6.8.2_01 FA_BPR_Ann_II_8_10_2_01 [REDACTED] 2008b
--------------------------------	-----------------------------------	---	---	--	---

No human data are available on adverse effects on fertility.

Considering the toxicity to fertility of formic acid, a two-generation reproduction toxicity study was conducted in the **rat** at dose levels of 0, 68, 203, 677 mg formate/kg bw/d administered orally in the feed as sodium formate over two parental (F0, F1) generations (DocIIIA6.8.2.-01, FA_BPR_Ann_II_8_10_2_01; ██████████ 2008b). At least 75 days after the beginning of treatment, F0 animals were mated to produce a litter (F1). Mating pairs were of the same dose groups and F1 animals selected for breeding were continued in the same dose group as their parents. Groups selected from F1 pups were to become F1 parental generation, were offered diets containing test substance post weaning, and the breeding program was repeated to produce F2 litter.

No treatment-related clinical signs or mortality were observed. Signs of **systemic toxicity** were observed in the F1 male parental generation at the highest dose. Food consumption and body weight gain were dose-dependently decreased. This resulted in secondary weight changes of brain and liver, but without correlating histopathological findings. Pathological examinations revealed no test-substance-related changes in organ weight, gross lesions, changes in differential ovarian follicle counts or microscopic findings. There were no indications that the **fertility or reproductive performance** of the F0 or F1 parental animals were affected. Estrous cycle data, mating behaviour, conception, gestation, parturition, lactation, weaning, and sperm parameters, sexual organ weights, and gross and histopathological findings of these organs (including differential ovarian follicle counts in the F1 females) were comparable between all test groups and ranged within the historical control data of the test facility. All data recorded during gestation and lactation (embryo/foetal/pup development) gave no indications of any **developmental toxicity** in the F1 and F2 offspring up to the highest dose level. Pup viability, pup body weight, sex ratio, sex maturation were not affected.

In conclusion,

NOAEL_{systemic} = 200 mg formate/kg bw/d for the F0 and F1 parental rats,

based on adverse effects on food consumption and bw gain at 670 mg formate/kg bw/d in the F1 parental males.

NOAEL_{fertility, reproductive performance} = 670 mg formate/kg bw/d for the F0 and F1 parental rats,

based on the lack of adverse effects at the highest dose.

NOAEL_{developmental} = 670 mg formate/kg bw/d for the F1 and F2 progeny,

based on the lack of adverse effects at the highest dose.

There were no negative findings on reproductive or on developmental parameters. The number and developmental of the pups was normal and comparable to the control. Formate, administered in the feed of rats as sodium formate, was not toxic with regard to reproduction or development.

In addition, there were no effects on fertility observed in the 13-week inhalation studies performed with formic acid vapours (0, 15, 30, 61, 122, 244 mg/m³). For a more detailed discussion, see section 3.6.3: DocIIIA6.4.3-01/ FA_BPR_Ann_II_8_9_2_03 and DocIIIA6.4.3-02/ FA_BPR_Ann_II_8_9_2_04: Thompson, 1992. There were no effects on measures of sperm motility, density, or testicular or epididymal weights, and no changes in the length of the estrous cycle. However, no functional fertility parameters were studied.

The effects of potassium diformate (oral feed for 140 d or 300d) in breeding sows was studied by ██████████ 2004 (DocIIIA6.4.1-02/ FA_BPR_Ann_II_8_9_2_02) and ██████████ 2003 (DocIIIA6.5.-02/ FA_BPR_Ann_II_8_9_4_0_JNS). For a more detailed discussion, see section 3.6.1/3.7.1. The studies focused on effects on fertility and, therefore, did not provide the full range of pathological and histopathological data which would be expected to be contained in guideline studies pertaining to chronic toxicity, reproduction toxicity, or developmental toxicity. However, the study provides additional data because the metabolic capability to dispose of formate is more limited in pigs. No treatment-related effects were observed for maternal toxicity (clinical signs, mortality, body weight, feed consumption), nor on ovulation, fertility, gestation parameters, number of live born piglets, piglet viability and weight gain until weaning up to doses of 301 mg formate/kg gbw/d for over 300 days.

Conclusion used in Risk Assessment – Fertility	
Value/conclusion	No adverse effects on fertility were observed for formate in rats. NOAEL _{parental, syst F0, F1} ~ 200 mg formate/kg bw/d; NOAEL _{fertility, reprod performance, developmental} ~ 670 mg formate/kg bw/d
Justification for the value/conclusion	There were no negative findings on reproductive or on developmental parameters. The number and development of the pups was normal and comparable to the control. Formate, administered in the feed of rats as sodium formate, was not toxic with regard to reproduction or development. Signs of systemic toxicity were observed in the F1 male parental generation at the highest dose.

Data waiving	
Information requirement	Adverse effects of formic acid on fertility
Justification	Sodium formate was applied to avoid unspecific toxic effects through the corrosive action of formic acid.

3.10.3 Effects on or via lactation

Conclusion used in Risk Assessment – Effects on or via lactation	
Value/conclusion	No adverse effects on or via lactation are expected for formic acid.
Justification for the value/conclusion	<p>Crossing of barriers as blood/brain, blood/testes, blood/placenta, and exposure via the breastmilk: It may be deduced from the physico-chemical properties of formic acid that the possibility of formate to cross the mentioned barriers is low. The substance is highly soluble in water and the logKow is around -2.0. The pKa is 3.70 at 20°C, and therefore formic acid (and the related salt potassium diformate) is almost exclusively present in the ionised form at physiological pH (DocIIIA6.2-01, FA_BPR_Ann_II_8_8_01). It is known that only the unionised form is likely to cross biological membranes, and that substances with a logP of 2-4 would likely cross membranes. The physico-chemical properties of formic acid differ largely, hence it is unlikely that formate would cross biological membranes. This does not preclude the uptake by means of active transport systems. Penetration into (and through) membranes may occur in minor quantities because the small size of the formate molecule. Transfer into breast milk may be given due to the high solubility in water. In this context it should also be mentioned that endogenous formic acid is produced in the intermediary metabolism in humans, and that the C1-fragment is required in the biosynthesis of amino acids and nucleic acids (DocIIIA6.2-09, FA_BPR_Ann_II_8_8_08), i.e. there is a need in the developing fetus. Excess blood formate is rapidly metabolised to background levels in humans, i.e. formate does not accumulate. Finally, there were no adverse effects noted in the testes, the brain, or the development of offspring, in any of the numerous studies requiring repeated dosing. This includes all subchronic and chronic repeated dose studies, carcinogenicity studies, multigeneration reproduction and teratogenicity studies, conducted in several species (rat, mouse, rabbit, pig) with either sodium formate or potassium diformate. Neurotoxicity is known to occur in humans only in the optical nerve following severe methanol intoxication leading to very high blood formate levels over an extended period of time (DocIIIA6.9). Thus, though formate crossing of the blood/brain, blood/testes, blood/placenta barriers, and the exposure via the breast milk cannot be fully excluded, no adverse effects were seen in the parental animals and their progeny of several species following high-level long-term dosing, or dosing during reproduction and development, of either sodium formate or potassium diformate.</p>

3.10.4 Overall conclusion on reproductive toxicity

Conclusion used in the Risk Assessment – Reproductive toxicity	
Value	<p>Two-generation study, rat:</p> <p>NOAEL_{parental} = 200 mg formate/kg bw/d</p> <p>NOAEL_{offspring} = 670 mg formate/kg bw/d</p> <p>NOAEL_{reproduction parameters} = 670 mg formate/kg bw/d</p> <p>Teratogenicity studies, rat, rabbit:</p> <p>NOAEL_{maternal} = 640 mg formate/kg bw/d</p> <p>NOAEL_{developmental} = 640 mg formate/kg bw/d</p>
Justification for the selected value	<p>The reproductive toxicity of formic acid was studied in a two-generation study in the rat administered orally in the feed as sodium formate (0, 68, 203, 677 mg formate/kg bw/d). The developmental toxicity of formic acid, administered by gavage as sodium formate, was studied in the rat (0, 40, 160, 640 mg formate/kg bw/d) and the rabbit (0, 68, 203, 677 mg formate/kg bw/d) teratogenicity studies.</p> <p>The two-generation study involving oral administration by feed of sodium formate in the rat showed that formate exerts no effect on the different reproduction parameters examined and induces no malformations in the selected dose range.</p> <p>NOAEL_{parental} = 200 mg formate/kg bw/d (based on reduced food consumption and body weight gain in F1 parental males at 670 mg formate/kg bw/d)</p> <p>NOAEL_{offspring} = 670 mg formate/kg bw/d</p> <p>NOAEL_{reproduction parameters} = 670 mg formate/kg bw/d</p> <p>The teratogenicity studies involving gavage administration of sodium formate in the rat and the rabbit showed that formate exerts no foetotoxic or teratogenic effects. No treatment-related effects were noted on the type and incidence of malformations and developmental variations in the selected dose range.</p>

	NOAEL _{maternal} = 640 mg formate/kg bw/d NOAEL _{developmental} = 640 mg formate/kg bw/d
Classification according to CLP and DSD	none

3.11 NEUROTOXICITY

Summary table of animal studies on neurotoxicity						
Method, Guideline, GLP status, Reliability	Species, Strain, Sex, No/group	Test substance, Dose levels, Duration of exposure	NOAEL, LOAEL	Results	Remarks (e.g. major deviations)	Reference
Mechanistic study GLP: no Rel. 2	Rat Long Evans, males 6/group	Methanol [CAS 67-56-1] purity unknown 4 g methanol/kg bw (20% w/v in saline) by i.p., followed by supplemental doses of 2 g/kg bw at 12-24 hour intervals Pre-treatment rats: exposed to a subanaesthetic concentration of nitrous oxide (N ₂ O/O ₂ 1:1) Route: i.p.		<p>Nitrous oxide inhibited methionine synthetase in pre-treated rats. The hepatic tetrahydrofolate (THF) level and the rate of formate oxidation were reduced to 50% compared to untreated rats (= comparable to the levels observed in monkeys and humans).</p> <p>Methanol intoxication of pre-treated rats resulted in acidosis and blood formate levels which were comparable to those seen in intoxicated monkeys and humans; Blood formate concentrations ranged between 8-15mM for 30-40 hours in the treated rats.</p> <p>Functional tests: Statistically significant changes were seen in both the retinal function (by electroretinogram) and the optical nerve integrity (by flash-evoked cortical potential) at 36 hours after the initial dose until the end of the experiment at 60 hours after initial dosing.</p> <p>Histopathology: The retina of the methanol-intoxicated rats showed diffuse edema and vacuolization at the junction of the inner and outer segments of the</p>		BPD ID A6.10_01 FA_BPR_Ann_II_8_13_5_01 Eells et al., 2000

				photoreceptor cells, and in the retinal pigmented epithelial cells. Mitochondrial cristae swelling was seen in the retinal pigmented epithelium cells and photoreceptors of intoxicated rats. Ultrastructural changes were much less pronounced in the optical nerve than in the retina																						
No guideline, but following scientific standards GLP: no Rel. 2	Monkey Rhesus males 4/group	Sodium formate [CAS 141-53-7] purity unknown Single i.v. 1.25 mmol/kg bw (57.5 mg/kg), followed by continuous infusion of 3.1mEq/kg bw/h = ~140 mg formate/ kg bw/h Rate of infusion such as to produce blood concentrations similar to those seen in methanol-intoxicated monkeys Route: i.v.		<p>Ocular/optic effects:</p> <p><u>Pupillary reflexes</u> were rapidly altered, and in most animals no response to light was observed between 24 and 48 h. Ophthalmology revealed marked <u>optic disc oedema</u> (mainly in the prelaminar region, central portion of the proximal part of the <u>optic nerve</u> without significantly reaching to the distal part. The retina including the ganglion-cell layer were completely normal.</p> <p>Blood pH: maintained between 7.4 and 7.6</p> <p>Blood formate levels:</p> <table border="1"> <thead> <tr> <th>Animal</th> <th>Conc. in blood</th> <th>Time of infusion</th> <th colspan="2">Clinical observations</th> </tr> <tr> <th>No.</th> <th>mg/l (mEq/l)</th> <th>h</th> <th>Pupillary reflex</th> <th>Fundus changes</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>1560 (34)</td> <td>39</td> <td>No response, mydriasis 7 mm</td> <td>Moderate optic disc edema with retinal edema</td> </tr> <tr> <td>2</td> <td>1380 (30)</td> <td>50</td> <td>No response, mydriasis 8 mm</td> <td>Severe optic disc edema</td> </tr> </tbody> </table>	Animal	Conc. in blood	Time of infusion	Clinical observations		No.	mg/l (mEq/l)	h	Pupillary reflex	Fundus changes	1	1560 (34)	39	No response, mydriasis 7 mm	Moderate optic disc edema with retinal edema	2	1380 (30)	50	No response, mydriasis 8 mm	Severe optic disc edema		BPD ID A6.2_05 FA_BPR_Ann_II_8_13_2_02 Martin-Amat et al., 1978
Animal	Conc. in blood	Time of infusion	Clinical observations																							
No.	mg/l (mEq/l)	h	Pupillary reflex	Fundus changes																						
1	1560 (34)	39	No response, mydriasis 7 mm	Moderate optic disc edema with retinal edema																						
2	1380 (30)	50	No response, mydriasis 8 mm	Severe optic disc edema																						

				3	920 (20)	41	No response, mydriasis 8 mm	Severe optic disc edema		
				4	550 (12)	25	Normal 3 mm	Moderate optic disc edema		

No regulatory neurotoxicity studies were made available for formic acid.

Formic acid, or formate, is associated with optical nerve and photoreceptor toxicity which is observed in humans and animals following methanol intoxication (DOCIIIA6.2_05, FA_BPR_Ann_II_8_13_2_02: Martin-Amat et al., 1978; DocIIIA6.10_01, FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000). See also Section 3.1.

The lesion may occur under conditions which allow formate to accumulate far above the background level, thus leading to high formate blood concentrations for extended periods of time. The blood levels of formate that correlate with the emergence of pathological changes are high. In a review by Eells et al. (2000) the following values after accidental and experimental methanol intoxication were summarised (see Review, Table 2):

TABLE 2. Blood formate, pH and bicarbonate concentrations in methanol-intoxicated rats, monkeys and humans.

Species	Blood Formate (mM)	Blood Bicarbonate (mEq/L)	Blood pH
N ₂ O - Treated Rats ^a	16.1 ± 0.7	7.7 ± 1.2	6.91 ± 0.06
Monkeys ^b	11.4 ± 1.2	6.5 ± 0.5	7.19 ± 0.02
Humans ^{c,d}	19.3 ± 4.4	3.2 ± 0.4	6.93 ± 0.02

Note: Methanol-intoxicated rats were exposed to a mixture of N₂O/O₂ (1:1) for 4 hours prior to methanol administration (4 g/kg at zero time followed by 2g/kg at 12-hour intervals) and exposure to the gas mixture was continued throughout the experiment. Blood formate concentrations and blood gas measurements were determined 60 hours after the initial dose of methanol. Each value represents the mean ± SE for 6 rats. Rodent data was compiled from studies by Eells *et al.*, (1996)^a. The monkey data was compiled from studies by Martin-Amat *et al.*, (1977)^b and the human data was compiled from studies conducted by McMartin *et al.*, (1980)^c and Eells *et al.*, (1991)^d.

Formate accumulated in all non-human primates (Rhesus monkey) 10 hours after an initial i.v. load of 57.5 mg formate/kg bw, followed by a continuous intravenous infusion of another 140 mg formate/kg bw/h. Maximum blood levels in the range 550 to 1560 mg/l were seen at 25 to 50 hours after the infusion had been started. The ophthalmological examinations revealed ocular problems evidenced by the lack of the light reflex, and moderate to severe retinal and optic disk edema. It is noteworthy that the blood pH was not changed by this treatment (BPD ID A6.2_05, FA_BPR_Ann_II_8_13_2_02; Martin-Amat et al., 1978).

Critical blood concentrations of 8 – 15 mM formate (= 360 – 680 mg/l) maintained over 30 – 40 hours were considered potentially detrimental, producing experimental ocular toxicity in monkeys (DocIIIA6.2_05, FA_BPR_Ann_II_8_13_2_02: Martin-Amat et al., 1978) and were associated with visual toxicity in acute cases of human methanol intoxication (DocIIIA6.10_01, FA_BPR_Ann_II_8_13_5_01: Eells et al., 2000).

In a review on methanol toxicity published by the CERHR Expert Panel (DocIIIA6.2_04, FA_BPR_Ann_II_8_8_03; NTP/USA, 2004), the background blood methanol and formate levels in humans have been reported to range between 0.6 and 2 mg methanol/l and between 3.8 and 11.2 mg formate/l. The blood methanol levels were increased in exposed males and females. However, inhalation exposure of 200 ppm methanol for 4 to 6 hours resulted in blood methanol levels of approx. 2 to 8 mg/l but had no influence on the blood formate levels (3.6 to 9.5 mg/l). It was further reported that the rate of formate oxidation in rats exceeds the maximal rate at which methanol is converted to formate: 1.6 versus 0.9 mmol/kg bw/h, respectively, whereas in non-human primates receiving moderately high doses the formate formation can exceed the oxidation of formate: 1.5 versus 0.75 mmol/kg bw/h, respectively.

An estimate of the methanol concentration that saturates the human folate pathway is 11 mM or 210 mg methanol/kg (DocIIIA6.2_04, FA_BPR_Ann_II_8_8_03: NTP/USA, 2004; BPD ID A6.2_12, FA_BPR_Ann_II_8_8_13: Kavet & Nauss, 1990). The latter would be equivalent to approx. 12.5 g methanol for a 60-kg adult.

The metabolic rate of 0.75 mmol formate/(kg bw*h) in pigtail monkeys is equivalent to approx. 34 mg formate/(kg bw*h) (DocIIIA6.2_04, FA_BPR_Ann_II_8_8_03: NTP/USA, 2004; BPD ID A6.2_12, FA_BPR_Ann_II_8_8_13: Kavet & Nauss, 1990). A metabolic saturation would occur only at higher intake rates. This finding is in line with the rapid and complete metabolism of formic acid or sodium formate observed in humans receiving 1, 2, or 3 g formic acid equivalents (DocIIIA6.2_07, FA_BPR_Ann_II_8_8_06; Malorny, 1969b).

The concept that ocular problems are associated with increased formate levels over an extended time period is supported by the findings of a mechanistic study (DocIIIA6.10_01, FA_BPR_Ann_II_8_13_5_01; Eells et al., 2000). Pretreatment of rats with a subanaesthetic concentration of nitrous oxide lowered the rat's folate pool and hence the formate oxidation rate (see review, Table 1), and rendered the rats susceptible to methanol poisoning, as evidenced by blood formate levels of 8 to 15 mM for 30 to 40 hours and functional and morphological changes of the photoreceptor and the optical nerve.

TABLE 1. Hepatic Folate Concentrations, Hepatic Tetrahydrofolate Concentrations and Rates of Formate Oxidation in Rats, N₂O-Treated Rats, Monkeys and Humans.

Species	Total Hepatic Folate (nmole/g)	Hepatic Tetrahydrofolate (nmole/g)	Rate of Formate Oxidation (mg/kg/hr)
Untreated Rats ^a	26.9 ± 3.3	14.2 ± 0.9	69 ± 1.6
N ₂ O-Treated Rats ^a	28.5 ± 1.2	8.5 ± 0.8	34 ± 1.0
Cynomolgus Monkeys ^a	25.5 ± 0.5	8.1 ± 0.2	34 ± 2.0
Humans ^b	15.8 ± 0.8	6.5 ± 0.3	N.D.

Note: Data compiled from studies conducted by Eells *et al.*, (1981, 1982)^a and Johlin *et al.*, (1987)^b.

Similar blood formate concentrations over these time periods have been shown to produce ocular toxicity in monkeys and are associated with visual toxicity in human methanol intoxication. Blood levels of 10 - 20 mM formate would be equivalent to 450 to 900 mg formate/l, based on the formate approx. molecular weight (approx. 45). Statistically significant changes were seen in both the retinal function (by electroretinogram) and the optical nerve integrity (by flash-evoked cortical potential) at 36 hours after the initial dose until the end of the experiment at 60 hours after initial dosing. The retina of the methanol-intoxicated rats showed diffuse edema and vacuolization at the junction of the inner and outer segments of the photoreceptor cells, and in the retinal pigmented epithelial cells. Mitochondrial cristae swelling was seen in the retinal pigmented epithelium cells and photoreceptors of intoxicated rats. Ultrastructural changes were much less pronounced in the optical nerve than in the retina (DocIIIA6.10_01, FA_BPR_Ann_II_8_13_5_01; Eells *et al.*, 2000).

The common pathophysiological basis of the so-called toxic optical neurotoxicity was recently reviewed by Altiparmak (2013; FA_BPR_Ann_II_8_13_5_03). Formate inhibits the mitochondrial cytochrome oxidase which results in disrupted energy supply and generation of reactive oxygen species (ROS). The prelaminar portion of the optic nerve has a higher number of mitochondria and a high oxygen demand; consequently, this portion is more vulnerable.

Waiver for further studies on neurobehavioral and neuropathological effects of formic acid:

It is known from methanol intoxications that methanol cause selective optical nerve toxicity. This toxicity likely occurs through a direct effect of formic acid (metabolite of methanol in the body). Although no effect on the optical nerve was seen in the toxicological studies with formic

acid or its salts, two studies specifically investigating these effects were added in the dossiers to account for the effects of formic acid when formed as an exclusive sequel of acute methanol intoxication.

The 2 animal studies on neurotoxicity provided are limited to investigations of the optical nerve and eye. A further study investigating neurobehavioral and neuropathological effects (in general) after single and repeated exposure is not available. However, neurotoxicity is part of the ADS. Further studies investigating neurobehavioral and neuropathological effects are only necessary if there is an indication, or knowledge from acute or repeated dose studies that the active substance may have neurotoxic properties.

Though the acute oral and inhalation toxicity studies show some behavioural changes, the repeated dose toxicity studies (Thompson '92, ██████████ '98, ██████████ 2002a/b) do not give rise to requesting additional neurotoxicity data as the main effects seen seem to be related mostly to irritation of the GIT and RT.

Human data on neurotoxicity:

In all human volunteer studies where formic acid or formate salts play a role, and in all human case reports, the single observation related to neurotoxicity is that formic acid, or formate, is associated with optical nerve and photoreceptor toxicity, which is frequently noted in humans following methanol intoxication. The aspect is addressed in more detail within the context of the toxico-kinetics and metabolism of formate in Section 3.1.

Conclusion used in Risk Assessment – Neurotoxicity	
Value/conclusion	Classification/labelling of the active substance 'formic acid' for neurotoxicity according to the criteria in Regulation 1272/2008/EC: none
Justification for the value/conclusion	<p>In methanol poisoning the metabolic capacity to dispose of formate is exceeded. The subsequent formate accumulation is characterized by very high blood formate levels in the range of 8 to 20 mM (i.e. approx. 350 to 900 mg/l) for more than 24 hours. Under such conditions, formate was demonstrated to cause functional and morphological changes of the retina and the optical nerve.</p> <p>It is conceivable that the ingestion of large doses of formate salts could have comparable results. The ingestion of large doses of formic acid would also cause high blood formate levels, but the acute effects, i.e. corrosivity</p>

and systemic toxicity, would prevail. Smaller doses of formate salts or formic acid are unlikely to saturate the metabolic rate which is 34 mg formate/kg bw/h in non-human primates.

Overall, lesions of the optical nerve and the photoreceptors are expected to occur only at formate doses, or formate precursor doses, which exceed by far the folate pathway saturation and thus cause high formate levels for an extended period of time. The proper use of biocidal products containing formic acid is unlikely to be associated with exposures that are sufficiently high to exceed the metabolic rate of approx. 34 mg formate/kg bw/h.

No further neurotoxicity testing is required because formate accumulation and adverse effects on the optical nerve and photoreceptors are considered to be an exclusive sequel of acute methanol intoxication in primates. Repeated dose toxicity studies do not give rise to requesting additional neurotoxicity data as the main effects seen seem to be related mostly to irritation of the GIT and RT.

3.12 IMMUNOTOXICITY

No data are available on immunotoxicity.

Conclusion used in Risk Assessment – Immunotoxicity	
Conclusion	There are no indications that Formic Acid has the potential to induce adverse effects involving the immune system.
Justification for the conclusion	There is no evidence from skin sensitisation, repeated dose or reproduction toxicity studies, that formic acid may have immunotoxic properties.

Data waiving	
Information requirement	Immunotoxicity study on Formic Acid
Justification	There is no evidence from skin sensitisation, repeated dose or reproduction toxicity studies, that formic acid may have immunotoxic properties. Hence, no specific study is required according to ECHA (2014) Guidance on the Biocidal Products Regulation v 1.1: Volume III: Human health - Part A: Information Requirements.

3.13 DISRUPTION OF THE ENDOCRINE SYSTEM

To assess potential effects on the endocrine system of formic acid the analysis of available information was conducted by implementing the assessment strategy outlined in the "Guidance for the identification of endocrine disruptors in the context of Regulations (EU) No 528/2012 and (EC) No 1107/2009" (ECHA/EFSA, 5 June 2018) referred hereafter as the "guidance on ED".

STEP 1 - Gathering of all relevant information

Level 1: existing data and existing or new non-test information

Formic acid is the simplest carboxylic acid. The formate anion is the common metabolite of formic acid and formate salts in aqueous solutions at physiological pH values. Formic acid and its conjugate base, formate, are also naturally occurring in virtually all living organisms as essential endogenous metabolites critical for one-carbon metabolism [Lamarre et al. 2013]. Formate is formed from precursors in the intermediary metabolism and is used as an important constituent of the C1 intermediary metabolism which is required for the biosynthesis of amino acids and nucleic acid bases (purines and pyrimidines). As a critical endogenous metabolite, formate is not assumed to be inherently endocrine active.

Endocrine activity was investigated using in silico methods. None of the endocrine activity related profilers of the OECD QSAR Toolbox V4.1 showed an alert for formic acid. In fact, formic acid was grouped into the category "non-binder, non-cyclic structure". Furthermore, binding to either oestrogen receptor (ER) or androgen receptor (AR) was estimated using in silico models implemented in OASIS TIMES (V2.27.19.13). None of the three models predicted a binding of formic acid to ER (with or without metabolisation of parent compound) and AR (without metabolisation). Please note that formic acid and formate have no structural similarity to intrinsic endocrine active substances (e.g. oestrogen, androgen). Altogether, based on in silico data it is very unlikely that formic acid exerts an endocrine/EATS-specific effect based on an endocrine mode of action.

Level 2: In vitro assays providing data about selected endocrine mechanism(s) /pathways(s)

Formic acid was not tested in any of the listed in vitro receptor binding or transactivation assays. Only some in vitro information is available from other studies (with reliability 3) but not specific of endocrine activity. Therefore they are not considered relevant regarding ED activity.

Level 3 In vivo assays providing data about selected endocrine mechanism(s) /pathway(s)

No information on such in vivo assays is available for formic acid.

Level 4 & 5 In vivo assays providing data on adverse effects on endocrine relevant endpoints

Table 3.1 Summary table of animal data on endocrine disruption*					
Summary table of animal data on endocrine disruption					
Method, Duration of exposure, Route of exposure, Guideline, GLP status, Reliability, Key/supportive study	Species, Strain, Sex, No/group	Test substance (including purity), Vehicle, Dose levels,	Results	Remarks (e.g. major deviations)	Reference
Two-generation reproduction Toxicity Oral OECD 416(2001) GLP Reliability 1	Rat Wistar rats, strain Crl:WI(Han) Male & Female 25 animals/dose group	Sodium formate Purity 100% Doses: 0; 100; 300; 1000 mg/bw sodium formate =0, 68, 203, 677 mg formate/kg bw/d	<u>For EAS-mediated:</u> No effect on : Age at preputial separation, Age at vaginal opening, Anogenital distance, Cervix histopathology, Coagulating gland weight and histopathology, Epididymis histopathology, Oestrus cyclicity, Genital abnormalities, Ovary histopathology, Oviduct histopathology, Prostate	NOAEL _{parental, syst F0, F1} ~ 200 mg formate/kg bw/d NOAEL _{fertility, reprod performance, developmental} 670 mg formate/kg bw/d	BPD ID A6.8.2_01 FA_BPR_Ann_II_8_10_2_01 ██████████ 2008b

			<p>histopathology and weight, Seminal vesicles histopathology and weight, Sperm morphology, motility and number, Testis histopathology and weight, Uterus histopathology and weight, Vagina histopathology and smears,</p> <p>Increase of relative cauda epididymis weight (300) (no effect on absolute weight and no effect in higher dose)</p> <p>Increase of relative ovary weight (300 and 1000) (no effect on absolute weight)</p>		
--	--	--	---	--	--

			<p><u>For Thyroid-mediated:</u> No effect on Thyroid histopathology and weight.</p> <p><u>For parameters sensitive to, but not diagnostic of, EATS:</u> No effect on: Adrenals histopathology and weight Pituitary histopathology, Live birth index, Male and Female fertility index, Gestation index and length, Lactation index, Litter size and viability, Number of implantations, Number of live births, Number of ovarian follicles, Post-implantation loss,</p>		
--	--	--	--	--	--

			<p>Pup weight, clinical observation and mortality, Male and female mating index, Sex ratio and Time to mating</p> <p>Increase (20%) of pituitary weight (68 and 677 mg) but no dose-response relationship</p> <p><u>For general toxicity:</u> Decrease of food consumption, body weight and absolute liver weight at 677 mg</p>		
<p>Subchronic oral toxicity in rodents</p> <p>13 weeks, Oral</p> <p>OECD 408</p> <p>GLP</p>	<p>Rat, CrI:CDBR</p> <p>Male & Female</p> <p>10 animals/sex/dose group</p>	<p>Potassium diformate Purity 95%</p> <p>Doses : 0, 600, 1200, 3000 mg Formi/kg</p>	<p><u>For EAS-mediated:</u> No effect on: Epididymis histopathology and weight, Macroscopic examination of mammary gland (M & F)</p>	<p>NOAEL_{Systemic}: as formate: 840 mg/kg bw/d LOAEL_{Systemic}: as formate: 2100 mg/kg bw/d</p>	<p>BPD ID A6.4.1_01</p> <p>FA_BPR_Ann_II_8_9_2_01 [REDACTED] 1998</p>

Reliability 1		bw/d (nominal) = 0, 420, 840, 2100 mg formate/kg bw/d	Ovary histopathology, Testis histopathology and weight, Uterus histopathology, Vagina histopathology <u>For Thyroid- mediated:</u> No effect on Thyroid histopathology <u>For parameters sensitive to, but not diagnostic of, EATS:</u> No effect on: Adrenals histopathology and Pituitary histopathology Decrease of adrenals weight (but not dose related and not statistically significant) <u>For general toxicity:</u>		
---------------	--	---	--	--	--

			Decrease of food consumption and body weight at 600 mg.		
<p>Subchronic inhalation toxicity</p> <p>13 weeks, Inhalation</p> <p>Similar to OECD 413 GLP</p> <p>Reliability 1</p>	<p>Rat, Fischer 344/N rats</p> <p>Male & Female</p> <p>10 animals/sex/dose group</p>	<p>Formic acid Purity 95%</p> <p>Dose: 0, 8, 16, 32, 64 and 128 ppm</p>	<p><u>For EAS-mediated:</u></p> <p>No effect on: Epididymis histopathology and weight, Oestrus cyclicity, Ovary histopathology, Prostate histopathology, Seminal vesicles histopathology, Sperm morphology, motility and numbers, Testis histopathology and weight, Uterus histopathology.</p> <p><u>For Thyroid-mediated:</u></p> <p>No effect on Thyroid histopathology</p> <p><u>For general toxicity:</u></p>	<p>NOAEL_{Systemic}: 244 mg/m³ (highest dose tested)</p>	<p>BPD ID A6.4.3_01; FA_BPR_Ann_II_8_9_2_03 Thompson, 1992</p>

			No effect		
<p>Subchronic inhalation toxicity</p> <p>13 weeks, Inhalation</p> <p>Similar to OECD 413</p> <p>GLP</p>	<p>Mouse, B6C3F1</p> <p>Male & Female</p> <p>10 animals/sex/dose group</p>	<p>Formic acid Purity 95%</p> <p>Dose: 0, 8, 16, 32, 64 and 128 ppm</p>	<p><u>For EAS-mediated:</u></p> <p>No effect on:</p> <p>Epididymis histopathology and weight, Oestrus cyclicity, Ovary histopathology, Prostate histopathology, Seminal vesicles histopathology, Sperm morphology, Testis histopathology, Uterus histopathology.</p> <p>Decrease in sperm motility (32 ppm) compared to controls but within the historical range for control mice. No dose-response relationship</p> <p>Increase sperm number (up to 33%) dose-</p>	<p>NOAEL_{Systemic}: 122 mg/m³</p> <p>LOAEL_{Systemic}: 244 mg/m³</p>	<p>BPD ID A6.4.3_01; FA_BPR_Ann_II_8_9_2_04 Thompson, 1992</p>

			<p>response relationship</p> <p>Increase relative testis weight (128 ppm) (no effect on absolute weight)</p> <p><u>For Thyroid-mediated:</u> No effect on Thyroid histopathology</p> <p><u>For general toxicity:</u> Decrease of body weight and absolute liver weight at dose 128 ppm. Increase of absolute liver weight (8.4%) at 32 and 64 ppm but not at 128 ppm (M) and decrease at 64 and 128 ppm (F)</p>		
<p>Combined chronic toxicity and oncogenicity study in the rat</p> <p>104 weeks, Oral</p>	<p>Rat, Wistar: CrI:HanWist(Glx:BRL)BR</p> <p>Male & Female</p>	<p>Potassium formate</p> <p>Purity: 98% and 99%</p>	<p><u>For EAS-mediated:</u> No effect on: Epididymis histopathology and weight, Macroscopic examination of</p>	<p>NOAEL_{Systemic}: as formate: 280 mg/kg bw/d LOAEL_{Systemic}: as formate: 1400 mg/kg bw/d</p>	<p>BPD ID A6.5_01/ BPD ID A6.7.-01 FA_BPR_Ann_II_8_9_3_01 FA_BPR_Ann_II_8_11_1_02 ██████████ 2002a/b</p>

<p>Similar to OECD 451-3</p> <p>GLP</p> <p>Reliability 1</p>	<p>20 animals/sex/dose group</p>	<p>Doses 0, 50, 400, 2000 mg/kg bw/d = 0, 35, 280, 1400 mg formate/kg bw/d</p>	<p>mammary gland (M & F), Ovary weight, Prostate histopathology, Testis histopathology and weight, Uterus histopathology and Vagina histopathology</p> <p>Decrease of incidence of fibroadenoma on mammary gland.</p> <p>Decrease in Ovary cysts in high dose females.</p> <p><u>For Thyroid mediated:</u> No effect on Thyroid histopathology</p> <p><u>For parameters sensitive to, but not diagnostic of, EATS</u> No effect on:</p>		
--	----------------------------------	--	---	--	--

			<p>Adrenals histopathology and weight Pituitary histopathology</p> <p><u>For general toxicity:</u> Decrease of food consumption, body weight and incidence of basophilic foci in liver at 1400 mg.</p>		
<p>Prenatal Developmental Toxicity Study</p> <p>Oral, day 6 to 29 post insemination</p> <p>OECD 414(2001) GLP</p> <p>Reliability 1</p>	<p>Rabbit, Himalayan Rabbit</p> <p>Females</p> <p>25 animals/dose group</p>	<p>Sodium formate, Purity 100%</p> <p>Doses: 0, 100, 300, 1000 mg/kg bw/day</p>	<p><u>For parameters sensitive to, but not diagnostic of, EATS:</u> No effect on: Foetal mortality and weight, Live foetus, Number of implantations, Pre and Post implantation loss, Placental weight, Resorption and Sex ratio</p> <p>Increase of fetal malformations (dose 1000) but within the historical range.</p>	<p>NO(A)EL teratogenicity embryotoxicity =670 mg formate/ kg bw/d</p> <p>NO(A)EL maternal =670 mg formate/ kg bw/d</p>	<p>BPD ID A6.8.1_02 FA_BPR_Ann_II_8_10_1_01 [REDACTED] 2008</p>

			For general toxicity: No effect		
<p>Prenatal Developmental Toxicity Study</p> <p>Oral, day 6 to 20 post coitum</p> <p>OECD 414(2001) GLP</p> <p>Reliability 1</p>	<p>Rat, Wistar: CrI:HanWist(Glx:BRL)</p> <p>Females</p>	<p>Sodium formate Purity >99%</p> <p>Doses: 0, 40, 160, 640 mg formate/(kg bw*d)</p>	<p>For parameters sensitive to, but not diagnostic of, EATS:</p> <p>No effect on: Foetal development, mortality and weight, Conception rate, Live foetus, Placental weight, Number of implantations, Pre and post implantation loss, Resorption and Sex ratio</p> <p>For general toxicity: No effect</p>	<p>NO(A)EL teratogenicity embryotoxicity =640 mg formate/kg bw/d</p> <p>NO(A)EL maternal = 640 mg formate/ kg bw/d</p>	<p>BPD ID A6.8.1_01 FA BPR Ann II 8_10_3_01 ██████████ 2005</p>
<p>Prenatal Developmental Toxicity Study</p> <p>Oral, 140 days</p> <p>No guideline GLP</p>	<p>Pig, Large White x Landrace hybrid Female</p> <p>6 animals/ dose group</p>	<p>Potassium diformate purity 98.7%</p> <p>Doses: 0, 157, 384, 753 mg/kg bw/d</p>	<p>For parameters sensitive to, but not diagnostic of, EATS:</p> <p>No effect on: Reproduction parameters and</p>		<p>BPD ID A4.4.1_02 B ██████████ (2004)</p>

Reliability 2			development of piglets at birth and until weaning. <u>For general toxicity:</u> No effect		
---------------	--	--	--	--	--

STEP 2 - Assemble and assess lines of evidence for endocrine activity and adversity

	Grouping	Lines of evidence	Species	Exposure, length	Route of exposure	Effect dose	Observed effects (positive or negative)	Assessment of each line of evidence	Assessment of the integrated line of evidence	Modality
Integrated line of evidence for endocrine adversity	EATS-mediated parameter	Thyroid histopathology	Rat	13 weeks	Oral	n.a.	No effect	No evidence of adversity		
				13 weeks	Inhalation					
				132 days	Oral					
				104 weeks	Oral					
		mouse	13 weeks	inhalation						
		Thyroid weight	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
Age at preputial separation	rat	132 days	oral	n.a.	No effect	No evidence of adversity				

		Age at vaginal opening	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Anogenital distance	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Cervix histopathology	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Coagulating gland histopathology	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Coagulating gland weight	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Epididymis histopathology	rat	13 weeks	oral	n.a.	No effect	No evidence of adversity	
			mouse	13 weeks	inhalation				
			rat	13 weeks	inhalation				
			rat	132 days	oral				
			rat	104 weeks	oral				
		Epididymis weight	rat	132 days	oral	203 mg (formate)/kg bw/day	Increase of relative cauda epididymis weight Due in part to the	Overall no evidence of adversity.	

							decrease of body weight (No effect on absolute weight). Moreover there is no dose-response relationship.		
			rat	13 weeks	oral	n.a.	No effect		
			mouse	13 weeks	inhalation				
			rat	13 weeks	inhalation				
			rat	104 weeks	oral				
		Estrus cyclicity	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of adversity	
			rat	13 weeks	inhalation				
			rat	132 days	oral				
		Genital abnormalities	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Macroscopic examination of mammary gland	rat	13 weeks	oral	n.a.	No effect	No evidence of adversity	
			rat	104 weeks	oral				

		Mammary gland histopathology	rat	104 weeks	oral	1400 mg(formate)/ kg bw/day	Decreased incidence of fibroadenoma. This is a known secondary effect of low body weight and is described commonly in the literature ¹	Overall no evidence of adversity.	
		Ovary histopathology	mouse	13 weeks	inhalation	n.a.	No effect	Overall no evidence of adversity.	
			rat	13 weeks	inhalation				
			rat	13 weeks	oral				
			rat	132 days	oral				
			rat	104 weeks	oral	1400 mg(formate)/ kg bw/day	Decrease in cysts in high dose females, related to a lower body weight.		
		Ovary weight	rat	132 days	oral	203 mg (formate)/kg bw/day	Increase relative ovary weight Due in part to the	Overall no evidence of adversity	

							decrease of body weight (No effect on absolute ovary weight). Moreover there is no dose-response relationship.		
			rat	104 weeks	oral	n.a.	No effect		
		Oviduct histopathology	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Prostate histopathology	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of adversity	
			rat	13 weeks	inhalation				
			rat	132 days	oral				
			rat	104 weeks	oral				
		Prostate weight	rat	132 days	oral	n.a.	No effect	No evidence of adversity	
		Seminal vesicles histopathology	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of adversity	
			rat	13 weeks	inhalation				
			rat	132 days	oral				

		Seminal vesicles weight	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		Sperm morphology	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of adversity		
			rat	13 weeks	inhalation					
			rat	132 days	oral					
		Sperm motility	Mouse	13 weeks	inhalation	6 ppm	Decrease of sperm motility with no dose-response relationship. Moreover the values for exposed mice fall well within the historical range for control mice	Overall no evidence of adversity		
			rat	13 weeks	inhalation	n.a.	No effect			
			rat	132 days	oral					
		Sperm numbers	Mouse	13 weeks	inhalation	32 ppm	Increase of concentration at 32 and 128 ppm	No evidence of adversity		

			rat	13 weeks	inhalation	n.a.	No effect			
			rat	132 days	oral					
		Testis histopathology	rat	13 weeks	oral	n.a.	No effect			No evidence of adversity
			mouse	13 weeks	inhalation					
			rat	13 weeks	inhalation					
			rat	132 days	oral					
			rat	104 weeks	oral					
		Testis weight	Mouse	13 weeks	inhalation	128 ppm	Increase of the relative testis weight at the higher dose. Related to a lower body weight (No effect on absolute testis weight)			Overall no evidence of adversity.
			rat	13 weeks	oral	n.a.	No effect			
			rat	13 weeks	inhalation					
			rat	132 days	oral					
			rat	104 weeks	oral					

	Uterus histopathology	rat	13 weeks	oral	n.a.	No effect	No evidence of adversity		
		mouse	13 weeks	inhalation					
		rat	13 weeks	inhalation					
		rat	132 days	oral					
		rat	104 weeks	oral					
	Uterus weight	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Vagina histopathology	rat	13 weeks	oral	n.a.	No effect	No evidence of adversity		
		rat	132 days	oral					
		rat	104 weeks	oral					
	Vaginal smears	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
Parameter sensitive to, but not diagnostic of EATS	Adrenals histopathology	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		rat	13 weeks	oral					
		rat	104 weeks	oral					
	Adrenals weight	rat	13 weeks	oral	2100 mg	Decrease of adrenals weight in the highest dose in females. Related to a lower	Overall no evidence of adversity		

						terminal body weight.				
		rat	104 weeks	oral	n.a.	No effect				
		rat	132 days	oral						
	Live birth index	rat	132 days	oral	n.a.	No effect	No evidence of adversity			
	Birth index	pig	>150 days	oral	n.a.	No effect	No evidence of adversity			
	Male and Female Fertility index	rat	132 days	oral	n.a.	No effect	No evidence of adversity			
	Fetal development	rat	17 days	oral	n.a.	No effect	Overall no evidence of adversity			
		rabbit	22 days	oral	1000 mg	Increase of fetal malformatio ns at the highest dose but within the historical control range.				
	Fetal mortality	rat	17 days	oral	n.a.	No effect	No evidence of adversity			
		rabbit	22 days	oral						
		pig	>150 days	oral						
	Fetal weight	rat	17 days	oral	n.a.	No effect				

			rabbit	22 days	oral			No evidence of adversity		
	Gestation index		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Conception rate		rat	17 days	oral	n.a.	No effect	No evidence of adversity		
	Gestation length		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Lactation index		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Litter size		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Litter viability		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
	Live fetus		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
			rat	17 days	oral					
			rabbit	22 days	oral					
	Number of implantations		rat	132 days	oral	n.a.	No effect	No evidence of adversity		
			rat	17 days	oral					
			rabbit	22 days	oral					

		Number of live births	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		Number of ovarian follicles	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		Pituitary histopathology	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
			rat	13 weeks	oral					
			rat	104 weeks	oral					
		Pituitary weight	rat	132 days	oral	100 & 1000 mg	Increase (20%) of pituitary weight but no dose-response relationship	Overall no evidence of adversity		
		Placental weight	rat	17 days	oral	n.a.	No effect	No evidence of adversity		
			rabbit	22 days	oral					
		Post implantation loss	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
			rat	17 days	oral					
			rabbit	22 days	oral					
		Pre implantation loss	rat	17 days	oral	n.a.	No effect	No evidence of adversity		
			rabbit	22 days	oral	n.a.	No effect			
		Pup development	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		Pup mortality	rat	132 days	oral	n.a.	No effect			

			pig	>150 days	oral			No evidence of adversity		
		Male and Female mating index	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
		Resorption	rat	17 days	oral	n.a.	No effect	No evidence of adversity		
			rabbit	22 days	oral					
		Sex ratio	rat	17 days	oral	n.a.	No effect	No evidence of adversity		
			rabbit	22 days	oral					
			rat	132 days	oral					
		Time to mating	rat	132 days	oral	n.a.	No effect	No evidence of adversity		
General toxicity		Body weight	mouse	13 weeks	inhalation	128 ppm	Decrease	Decrease of body weight at high doses only; related to the decrease of food consumption		
			pig	>150 days	oral	n.a.	No effect			
			rabbit	22 days	oral	n.a.	No effect			
			rat	13 weeks	Oral	600 mg	Decrease			
			rat	17 days	oral	n.a.	No effect			
			rat	132 days	oral	1000 mg	Decrease			
			rat	104 weeks	oral	2000 mg	Decrease			
			rat	13 weeks	inhalation	n.a.	No effect			

	Food consumption	pig	>150 days	oral	n.a.	No effect	Decrease of food consumption at high doses only.	
		rabbit	22 days	oral				
		rat	17 days	oral				
		rat	13 weeks	Oral	600 mg	Decrease		
		rat	132 days	oral	1000 mg	Decrease		
		rat	104 weeks	oral	2000 mg	Decrease		
	Liver histopathology	mouse	13 weeks	inhalation	n.a.	No effect	Overall no evidence of liver toxicity, except at high dose in 1 study.	
		pig	>150 days	oral				
		rat	13 weeks	inhalation				
		rat	104 weeks	oral	2000 mg	Increase hepatocyte vacuolisation, eosinophilic and basophilic foci in the liver of high dose males.		
	Liver weight	mouse	13 weeks	inhalation	32 ppm	Absolute liver weight: Increase (8.4%) at 32 and 64 ppm but not at 128 ppm (M) and	Minor effects in liver weight.	

							decrease at 64 and 128 ppm (F)			
			rat	13 weeks	Oral	n.a.	No effect			
			rat	132 days	oral	1000	Decrease (7.5%) Related to the reduced body weight.			
			rat	13 weeks	inhalation	n.a.	No effect			
			rat	104 weeks	oral					
		Kidney histopathology	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of adversity		
			pig	>150 days	oral					
			rat	132 days	oral					
			rat	13 weeks	inhalation					
			rat	104 weeks	oral					
		Kidney weight	mouse	13 weeks	inhalation	64 ppm	Increase of relative kidney weight at 64 ppm (F) and 128 ppm (M&F) Related to the reduced	Minor effects in kidney weight in 2 studies.		

							body weight (no effect on absolute kidney weight)				
			rat	132 days	oral	300 mg	Increase of absolute and relative kidney weight at dose 300 and 1000. (up to 8.1%)				
			rat	13 weeks	oral	n.a.	No effect				
			rat	13 weeks	inhalation						
			rat	104 weeks	oral						
		Brain histopathology	rat	104 weeks	oral	n.a.	No effect	No evidence of adversity			
		Brain weight	rat	132 days	oral	300 mg	Increase of relative brain weight in female F0 (but reduced terminal body weight)	Overall no evidence of adversity			
			rat	104 weeks	oral	n.a.	No effect				
		Mortality	mouse	13 weeks	inhalation	n.a.	No effect	No evidence of			

			pig	>150 days	oral			effect on mortality		
			rabbit	22 days	oral					
			rat	13 weeks	Oral					
			rat	17 days	oral					
			rat	132 days	oral					
			rat	104 weeks	oral					
			rat	13 weeks	inhalation					

¹Ghanta, N. R. et al (1987): Influence of body weight on the incidence of spontaneous tumours in rats and mice of long term studies. American Journal of Clinical Nutrition 45: 252-260

Roe, J. C. (1987): The problem of pseudocarcinogenicity in rodent bioassays.

Banbury Report 25: Nongenotoxic Mechanisms in Carcinogenicity.

STEP 3 - Sufficiency of the dataset

For EAS parameters, according the guidance on ED, a two-generation reproductive toxicity study (OECD TG416, test protocol according to latest version of January 2001) is enough to consider that EAS-mediated adversity has been sufficiently investigated.

A two-generation reproductive toxicity study was performed with sodium formate in 2008 according to OECD TG416 guidelines (2001).

Therefore, **EAS-mediated parameters are considered to be sufficiently investigated.**

Regarding thyroid, the available ED adversity related studies (OECD 416, 414 (two species), 408, 451-3 and 413 (two species)) did not investigate all thyroid parameters, since some of the studies are old and did not recorded mandatory T parameters (T3/T4 and TSH level, HDL/LDL ratio and thyroid weight for 2 key studies are missing).

No effect of formic acid was detected in the investigated parameters (macroscopic aspect, histopathology and weight).

Since no adverse effect on thyroid was recorded in the life time carcinogenicity study or in the others available studies, it was agreed at the 14th ED expert group (4-5 june 2019) to consider that the **data set is sufficient for Thyroid.**

Please also note that further vertebrate testing was not supported because the substance is corrosive to the gastro-intestinal tract at low doses.

STEP 4 - Initial analysis of the evidence

According the available studies, there is no evidence of adversity for either "EATS-mediated" or "sensitive to but not diagnostic of EATS" parameters.

Effects on liver and kidney, were recorded in some studies but they are inconsistent between sex, studies and species and cannot be explained by an endocrine pathway.

According the guidance on ED, page 36, scenario 1a is concluded and therefore **ED criteria are not met for Human Health.**

Conclusion used in Risk Assessment – Endocrine disruption	
Conclusion	ED criteria not met for Human Health
Justification for the conclusion	Scenario 1a : No evidence of EATS-mediated adversity

3.14 FURTHER HUMAN DATA

Summary table of further human data				
Type of data/ report, Reliability	Test substance	Relevant information about the study	Observations	Reference
Report on workplace exposure	Formic acid	Measurement of formic acid at the workplace (8-hour time weighed average) 138 workplace measurements	production, filling, processing, laboratory. The mean values and the 95% percentiles were all far below the threshold limit of 5 ppm or 9.6 mg/m ³ . The highest values were seen at the filling station, but still below the threshold, with the mean value of 1.1 mg/m ³ , and the 50%, 90% and 95% percentiles at 0.65, 2.7 and 8.2 mg/m ³ , respectively.	DocIIIA6.12.1-01 FA_BPR_Ann_II_8_12_1_01 ██████████ 2006
Health records from industry	Formic acid Concentration not stated; presumably 50-85%	Sex: not reported Age: 25, 20, 34 and 53 years Route of exposure: dermal	Lesions of skin and eye following facial splashes (3 cases) during filling operations and transportation; one case of skin lesions following contact with contaminated wood.	DocIIIA6.12.3-01 FA_BPR_Ann_II_8_12_3_01 ██████████ 1994, 2002
Case report	Formic acid 60%	1 male, 27-year-old Route of exposure: oral	Suicidal ingestion, 45-90 ml (decalcifying agent). Clinical signs: vomiting, abdominal pain Blood: pH 6.86, pCO ₂ 70.4 mmHg, HCO ₃ 10.6 mmol/l, base deficit -22 mmol/l, initial serum formate level 370.3 µg/ml, haemolysis Autopsy: ulceration of oesophagus, complete necrosis of gastric mucosa, oedema and necrotic areas in deeper tissue	BPD ID A6.12.2_01 FA_BPR_Ann_II_8_12_2_01 Westphal et al., 2001

			<p>layers of stomach, no perforation, coagulated blood in stomach, necrosis of mucosa duodenum.</p> <p>Post-mortem formate concentrations: 855.4 µg/ml (heart blood) 2712 µg/ml (gastric contents) 1128 µg/ml (hemorrhagic fluid abdominal cavity) 3051 µg/ml (bile) 2664 µg/ml (contents small intestine) 442.7 µg/g (liver) 542.3 µg/g (kidney)</p> <p>Within 30 hours after ingestion: corrosion of the gastrointestinal tract, metabolic acidosis, haemolysis, massive bleeding, hepatic and renal failure, death.</p>	
Case report	Formic acid 50%	1 female, 39-year-old Route of exposure: oral	<p>Suicidal ingestion, 200 ml (descaling product).</p> <p>Clinical signs: severe retrosternal and epigastric pain, dyspnea, cyanotic appearance, vomiting blood (2 h after ingestion)</p> <p>Blood: pH 6.87, pCO₂ 46.1 mm Hg, HCO₃ 8.6 mmol/l, base deficit of -26.4 mmol/l, haemolysis (20 min after admission to hospital)</p> <p>Initial serum formate level 348 µg/ml (7.6 mmol/l), elimination T_{1/2} 2.5 hours</p> <p>Urine: red</p> <p>Gastroscopy: severe lesions oesophagus and stomach, superficial burns duodenum</p> <p>Complications: severe gastrointestinal bleeding, pneumonia, acute tubular necrosis, adult respiratory distress syndrome, peritonitis, sepsis</p> <p>Result:</p> <p>Local: corrosion and massive bleeding, loss of blood pressure</p> <p>Systemic: Severe metabolic acidosis and haemolysis, renal failure</p>	<p>BPD ID A6.12.2_02</p> <p>FA_BPR_Ann_II_8_12_2_02</p> <p>Verstraete et al., 1989</p>

			Death: 6 weeks after ingestion	
Case report	Formic acid conc. not known	30 males 23 females 16 to 46 year-old Route of exposure: oral	<p>Suicidal ingestion, ≥ 10 ml, (rubber workers)</p> <p>Major complications:</p> <p>Gastro-intestinal: facial burns, ulcerations of oral and pharyngeal mucosa, abdominal pain, contractures and keloid formation of affected skin, oesophagus stricture (16/53 cases) requiring reparative surgery</p> <p>Respiratory system: inhalation pneumonitis (45 of 53 patients) with cough dyspnea, cyanosis, could proceed to respiratory infection and failure</p> <p>Vascular hypotension: 17/53 cases</p> <p>Haemolysis, haematuria within few hours of ingestion, rapidly followed by renal failure in severe cases, within a day in less severe cases, in total 20/53 cases</p> <p>Result:</p> <p>Local: corrosion and massive bleeding, loss of blood pressure</p> <p>Systemic: Severe metabolic acidosis and haemolysis, renal failure</p> <p>Death: 15/53 patients</p>	<p>BPD ID A6.12.2_03</p> <p>FA_BPR_Ann_II_8_12_2_03</p> <p>Rajan et al., 1985</p>
Case report	Formic acid 40-55%	1 male 2 females 35, 56, 66 year-old Route of exposure: oral	<p>Suicidal ingestion, estimated volumes 'one mouthful' to 50-100 ml (descaling product)</p> <p><i>35-year-old woman, 40% formic acid, 3 mouthfuls:</i> massive bleeding, haemolysis, died on d14 after shock and massive haematemesis. Ulcerations throughout oesophagus and stomach, tubular necrosis, early thrombosis of the portal vein</p> <p><i>66-year-old woman, 55% formic acid, 55 to 100 ml:</i> massive bleeding, haemolysis, extensive erosion of oesophagus, stomach, duodenum, died on d5</p>	<p>BPD ID A6.12.2_04</p> <p>FA_BPR_Ann_II_8_12_2_04</p> <p>Naik et al., 1980</p>

			<p><i>56-year-old man, mouthful of 55% formic acid:</i> died on d11 due to circulatory failure</p> <p>Result: Local: corrosion and massive bleeding, loss of blood pressure Systemic: Severe metabolic acidosis and haemolysis, renal failure Death</p>	
Case report	Formic acid 44 to 60%	male/female <12 years to adult 45 cases Route of exposure: oral	<p>Accidental and suicidal ingestion</p> <p>Estimated doses: < 10 g (children) to 200 g (adults)</p> <p>Children: accidental ingestion of low doses (≤ 10 g), reversible oropharyngeal burns in 9 children, no deaths</p> <p>Adults: suicidal ingestion (34/36 cases), accidental ingestion (2/36)</p> <p><u>5-30 g</u>: reversible oropharyngeal burns (16); abdominal pain, vomiting, dyspnea, dysphagia (5); hematemesis, pneumonitis, esophageal strictures (2)</p> <p><u>30-45 g</u>: intravascular coagulation, acute renal failure, hematemesis, liver impairment, oesophageal strictures</p> <p><u>45-200 g</u>: corrosive perforations of the abdominal viscera and gastrointestinal hemorrhage, acute renal failure</p> <p>dose up to 45g: 28/29 patients survived dose 45g-200g: 14/16 patients died</p> <p>Result: Local: corrosion and massive bleeding, loss of blood pressure Systemic: Severe metabolic acidosis and haemolysis, renal failure Death</p>	<p>BPD ID A6.12.2_05 FA_BPR_Ann_II_8_12_2_05 Jefferys and Wiseman, 1980.</p>

Case report	Formic acid 87 to 96%	male/female children 183 cases Route of exposure: oral	Accidental ingestion: only small quantities Vomiting (10/183 children) and visible caustic lesions in mouth and throat (28/183 cases) Result: Reversible burns of oesophagus	BPD ID A6.12.2_06 FA_BPR_Ann_II_8_12_2_06 von Muehlendahl et al., 1978
Case report	Formic acid conc. not known	1 male, 35-year-old Route of exposure: dermal	Accidental splash from a container on the maxilla, chin, around mouth, thorax (occupational) Clinical signs: burning pain, sialorrhoea, nausea, vomiting Skin: blisters, necrotic areas Systemic: blood pressure 110/60, pulse and breathing regular, blood gases and acido-balance normal, no formic acid detected in blood and urine Result: skin corrosion	BPD ID A6.12.2_07a FA_BPR_Ann_II_8_12_2_07 Malizia et al.,1977
Case report	Formic acid undiluted, conc. not known	1 female, 15-year-old Route of exposure: dermal	Accidental splash on lower extremities (20% of total body surface) Clinical signs: burns, nausea, vomiting (4 hrs after exposure = start treatment) Skin: depth of burns not determined, became full-thickness. Gross oedema on d2 and d3 without fever, ocular damage or pulmonary complications. Burns surgically revived on d16, grafted several times. Major scarring of burned areas persisted. Urine: brownish, hemoglobinuria Blood: pH 7.23, HCO ₃ 16.7 mmol/l, base deficit 9.5, hemolysis Patient recovered rapidly from metabolic acidosis. Result: Skin corrosion Mild metabolic acidosis	BPD ID A6.12.2_08 FA_BPR_Ann_II_8_12_2_08 Sigurdsson et al., 1983

Case report	Formic acid 90%	1 female, 3-year-old Route of exposure: dermal	<p>Accidental splash on right torso and extremities (35% of total body surface)</p> <p>Clinical signs: severe distress (10 min after exposure = start treatment)</p> <p>Skin: full-thickness second- and third-degree burns. Required several skin grafts during several months</p> <p>Urine: initially dark red, hemoglobinuria resolved within few days without kidney failure</p> <p>Blood: pH 6.85, HCO₃ 16.7 mmol/l, base deficit -29.7 on 100% oxygen, bicarbonate 6mEq/l; initial serum formate level 400 µg/ml, hemolysis</p> <p>Patient recovered rapidly from metabolic acidosis.</p> <p>Result:</p> <p>Skin corrosion</p> <p>Metabolic acidosis</p>	BPD ID A6.12.2_09 FA_BPR_Ann_II_8_12_2_09 Chan et al., 1995
Case report	Formic acid 98%	1 male, 39-year-old Route of exposure: inhalation	<p>Accidental spray (aerosol) into the face with concomitant inhalation (occupational)</p> <p>Clinical signs: facial burns (3% of total body surface), dyspnea</p> <p>Nasopharyngoscopy: mild supraglottic erythema, normal vocal cords</p> <p>Skin: second-degree burns</p> <p>Pulmonary function tests: Vital capacity reduced on d1, recovered largely within 14 days. Complains of dyspnea till d15</p> <p><u>Day 1</u></p> <p>FVC (L): 3.74 (79% predicted) FEV₁ (L): 2.86 (73% predicted) FEV₁/FVC: 76.38 (92% predicted) FEF_{25%-75%} (l/sec): 2.32 (56% predicted)</p> <p><u>Day 15</u></p> <p>FVC (L): 4.35 (92% predicted) FEV₁ (L): 3.62 (92% predicted)</p>	BPD ID A6.12.2_10 FA_BPR_Ann_II_8_12_2_10 Yelon et al., 1996

			<p>FEV₁/FVC: 83.09 (101% predicted) FEF_{25%-75%} (l/sec): 3.82 (92% predicted)</p> <p>Result: Reversible Pulmonary dysfunction: Reactive Airway Dysfunction Syndrome</p>	
Case report	Fumes from formic acid (85%) and carbon monoxide (concentration not known)	1 male, 22-year-old Route of exposure: inhalation	<p>Suicide by mixing formic acid with concentrated sulphuric acid in a confined space</p> <p>External chemical burns</p> <p>Internal injuries mainly to the respiratory tract. Injury to the oropharyngeal area and trachea, pulmonary edema, and subpleural petechiae. Complete lack of the respiratory epithelium of the trachea, edema of mucosa, and submucosa of the trachea, thrombi, and hemolysis inside the small vessels of the trachea, pulmonary edema, hemolysis, and thrombosis in the lung vessels</p> <p>Death due to CO intoxication; corrosion/irritation of skin, trachea, lungs, stomach due to formic acid fumes.</p>	Bakovic M, et al (2015) FA_BPR_Ann_II_8_12_2_11
Case report	Fumes from formic acid (concentration not reported, amount 950 ml) and carbon monoxide (concentration not known)	1 male, 26-year-old Route of exposure: inhalation	<p>Suicide by mixing formic acid with concentrated sulphuric acid in a confined space. Death. The body showed pronounce bright pink-red lividity. The autopsy was otherwise unremarkable.</p> <p>No further info on formic acid effects.</p>	Lin PT and Dunn (2014) FA_BPR_Ann_II_8_12_2_12
Case report	Fumes from formic acid (98-100%) and carbon monoxide	1 male, 26-year-old; 1male, 53-year-old, 1	<p>Suicide by mixing formic acid with concentrated sulphuric acid in a confined space</p> <p>26-year-old: death. No autopsy</p>	Yang CC et al. (2008) FA_BPR_Ann_II_8_12_2_13

	(concentration not known)	female, 53-year-old Route of exposure: inhalation	53-year-old father: coma, hypoxemia, metabolic acidosis, and a carboxyhemoglobin level of 45.8%. Developed acute respiratory distress syndrome. Transient ulceration of vocal cords. 53-year-old mother: dizziness, headache, carboxyhemoglobin level of 23.0% In addition to the toxicities of carbon monoxide, concomitant inhalation of formic acid fumes can cause severe lung injury, which may complicate the management of carbon monoxide poisoning.	
Retrospective study	formic acid	302 cases Males and females Age: 29.7-55, mean age 42.8 years Route of exposure: Oral, dermal, inhalation	Suicide Mean (SD) quantity consumed: 110 (78) mL The most common symptoms noted at presentation were: vomiting (78.5 %) abdominal pain (56.3%) hematemesis (48.3%) respiratory distress (44 %) haematuria (30.1%) oliguria (24.5%) hypotension (24.5%) melena (22.2%) direct corneal injury (0.007%) Mean (SD) pH of all patients was 7.3 and the bicarbonate concentration was 19.2 (5.1) mEd/L. Leucocytosis was seen in 57.5% of the patients; liver enzymes (GOT, GPT) were elevated above normal values in 62.1% of the patients. The effectivity of medical treatment depends largely on the ingested dose and concentration of FA, the time delay after exposure. Low blood pH and bicarbonate concentration reflect the severity.	Dalus D et al. (2013) FA_BPR_Ann_II_8_12_5_01

			<p>The mortality rate was 35.4%. Bowel perforation, shock, and tracheoesophageal fistula were associated with 100% mortality.</p> <p>A higher blood pH was less likely to result in mortality. Dysphagia was noted in 154 patients, 98 of whom showed oesophageal stricture on evaluation, requiring repeat endoscopic dilatations after discharge. The prevalence of oesophageal stricture among the 195 patients who survived was 50.2%.</p>	
--	--	--	--	--

Medical surveillance on manufacturing plant personnel:

A total of 138 workplace measurements have been conducted during the period 2001-2006, covering all kinds of operations (production, filling, processing, laboratory). All reported results represented 8 hours shift average values (TWA) obtained by personal air sampling. None of the measurements exceeded the threshold limit of 5 ppm or 9.6 mg/m³ (most well below). To prevent direct skin contact, protective gloves (neoprene or nitrile rubber) are used. According to the applicant workplace exposure is low, due to the appropriate protective measures taken. Consequently, medical surveillance on plant personnel is not required (DocIIIA6.12.1-01: ██████████, 2006).

Four cases of accidental skin and eye contact were seen during 14 years (1989-2002) of operation of BASF's production plant. Lesions of skin and eye were seen following facial splashes (3 cases) during filling operations and transportation, and one case of skin lesions following contact with contaminated wood. As concentrated formic acid is corrosive, the employees underwent First Aid measures and required further medical treatment in hospital. Type and duration of medical treatment were not reported, nor the outcome in the health records (DocIIIA6.12.3-01: ██████████ 1994, 2002).

Clinical cases and poisoning incidents (professional operators and the general population), Expected effects of poisoning, aspects of diagnosis of poisoning, prognosis:

Oral ingestion

There are published cases of accidental ingestion of formic acid, but the incidence is relatively low. The suicidal ingestion (34 of 36 cases, i.e. 94%) clearly prevailed over the accidental ingestion (2 of 36 cases) in adults (DocIIIA6.12.2_05, FA_BPR_Ann_II_8_12_2_05: Jefferys and Wiseman, 1980). Easy access to formic acid was considered to promote the suicidal ingestion of formic acid in the State of Kerala, India, among workers of the rubber industry who used formic acid as a coagulant (DocIIIA6.12.2_03, FA_BPR_Ann_II_8_12_2_03: Rajan et al., 1985).

In children, the accidental ingestion occurs generally at low doses, i.e. **up to 10 g formic acid**, which reportedly caused reversible burns of the pharyngeal tract in 9 children, who all survived (DocIIIA6.12.2_06, FA_BPR_Ann_II_8_12_2_06: von Muehlendahl et al., 1978). The consumption of only small quantities might be related to the pungent smell of formic acid.

The doses are much higher in cases of deliberate ingestion by adults. Doses **up to 45 g** formic acid were survived by 28 of 29 patients. Most of the patients died (14 of 16; 88%) after doses between 45 – 200 g formic acid (DocIIIA6.12.2_05, FA_BPR_Ann_II_8_12_2_05: Jefferys and Wiseman, 1980). In a retrospective study with 302 patients who committed suicide, the estimated mean ingested quantity was 110 mL of formic acid. The mortality rate was 35.4% in this study. The prognosis depended largely on the concentration of formic acid and the amount ingested and the lag time until onset of medical treatment (FA_BPR_Ann_II_8_12_8_02: Dalus et al., 2013).

Due to the corrosivity of formic acid, local effects must be expected at all dose levels. The amount ingested and the concentration determine the grade and the location of the effects. Therefore, the observations range from moderate burns around the mouth to severe corrosion of the gastro-intestinal tract with destruction of the esophagus, perforation of the stomach, and corrosion of the small intestine together with massive bleeding and systemic toxicity:

- Nine children accidentally ingested **less than 10 g of formic acid**. They suffered oropharyngeal burns, which were only superficial, and they fully recovered. Two adults accidentally ingested formic acid, whilst 34 deliberately consumed it.
- Consumption, by 23 subjects, of between **5 and 30 g of formic acid** produced no deaths. The majority (16) developed minor superficial oropharyngeal burns only. Five had more severe symptoms including abdominal pain, vomiting, dyspnea and dysphagia, whilst two experienced sustained hematemesis and pneumonitis, and subsequently developed esophageal strictures.
- Ingestion of **30-45 g of formic acid** produced more serious effects. Of the six patients recorded, one died, one had reversible disseminated intravascular coagulation and three had reversible acute renal failure. All suffered hematemesis and had biochemical evidence of liver impairment. Four needed subsequent treatment for esophageal strictures.
- Ingestion of **45 to 200 g of formic acid** was recorded from 16 patients, of whom 14 died; two recovered. Considering the fatalities, the majority (9) died painfully within the first 36 hours from corrosive perforations of the abdominal viscera and from gastrointestinal hemorrhage. The other five developed acute renal failure which contributed to their death (BPD ID A6.12.2_05, FA_BPR_Ann_II_8_12_2_05).
- **Systemic toxicity** was seen after ingestion of **30 g formic acid** or more.

Prognosis is poor after massive oral ingestion (>45 to 200 g formic acid); prognosis is moderate after moderate oral ingestion (approx. 30 to 45 g); lesions, but low mortality, are expected in most cases with low amounts ingested (<30g); persistent lesions due to tissue corrosion must be expected in cases with >10 g formic acid ingested. Tissue destruction of the gastrointestinal tract may result in fatal bleeding, septic shock, or stricture which may require surgical treatment. Reversibility of effects was often seen in cases with low amounts ingested (<10 g formic acid).

Dermal exposure

Due to the corrosivity of concentrated formic acid, local effects must be expected following contact to the skin and to the eyes.

Prognosis: Local burns heal only slowly. Tissue destruction of the skin may result in scarring.

Systemic effects may result after contact of concentrated formic acid to extended areas of the body surface (DocIIIA6.12.2_07, FA_BPR_Ann_II_8_12_2_07: Malizia et al., 1977; DocIIIA6.12.2_08, FA_BPR_Ann_II_8_12_2_08: Sigurdsson et al., 1983; DocIIIA6.12.2_09, FA_BPR_Ann_II_8_12_2_09: Chan et al., 1995).

Prognosis: Systemic effects were reversible within few days without sequelae in cases where the medical treatment was rapid and strict to counteract the metabolic acidosis.

Inhalation exposure

Due to the warning effect of the pungent smell of formic acid, inhalation exposure is generally low.

As **local effect**, pulmonary dysfunction was observed which was reversible within 14 days in one presumably high-dose case (DocIIIA6.12.2_10, FA_BPR_Ann_II_8_12_2_10: Yelon et al., 1996).

Inhalation of fumes created by mixing formic acid with concentrated sulphuric acid leads to injuries to the respiratory tract from formic acid, and deadly carbon monoxide intoxication (Bakovic et al., 2015; Lin & Dunn, 2014; Yang et al., 2008).

Systemic effects are unlikely to occur. An estimate that was presented in the MAK-justification indicated that the uptake of formic acid at the threshold exposure concentration (MAK-value: 5 ppm i.e. 9.5 mg/m³) equals approx. 0.5% of the metabolic rate observed in non-human primates. It was therefore concluded that an effect on the blood pH is unlikely. Formic acid inhalation concentrations from 30 ppm onwards were regarded as being immediately dangerous to life and health (DocIIIA6.12.8_01: Greim, 2003; NIOSH, 1990).

Aspects of diagnosis: Effective treatment requires an examination which provides adequate poisoning information. The case history provides information on the route of exposure and in some cases on the chemical concentration and amount. Clinical signs (mouth or skin affected) support this. The examination should generally comprise (1) and additionally (2) in cases of inhalation exposure:

(1) Blood pressure, blood count, hemolysis, blood gases, acid-balance, urine. Blood and urine formate concentrations.

(2) Inhalation (additionally): Chest radiograph, Lung function tests

First aid measures, therapeutic regimes

The primary goal must be to restore the metabolic acidosis to counteract the systemic toxicity. Second, the burns must be appropriately treated including the use of antibiotics. Special attention requires internal bleeding, due to local corrosion of the gastrointestinal tract after oral ingestion.

After suicidal exposures the doses are often extremely high, and there is no specific treatment in such cases.

Conclusion on prognosis: The prognosis depends on the exposure (concentration of chemical, amount, route of exposure), the rapid onset of treatment, the proper examination on admission to the hospital, and a strict treatment regimen to counteract systemic and local effects.

The prognosis may be good in cases of low oral, dermal, and inhalation exposure, as the systemic toxicity may be low. The prognosis of severe systemic toxicity is often bad. Tissue corrosion due to local effects heals slowly with scarring in most cases.

Conclusion used in Risk Assessment – Further human data

Conclusion

Dermal exposure:

Due to the corrosivity of concentrated formic acid, local effects must be expected following contact to the skin and to the eyes. Local burns heal only slowly. Tissue destruction of the skin may result in scarring. Systemic effects may result after contact of concentrated formic acid to extended areas of the body surface. Occupational and accidental dermal exposure records report skin corrosion and metabolic acidosis.

Oral exposure:

Due to the corrosivity of formic acid, local effects must be expected at all dose levels. The amount ingested and the concentration determine the grade and the location of the effects. Therefore, the observations range from moderate burns around the mouth to severe corrosion of the gastro-intestinal tract with destruction of the esophagus, perforation of the stomach, and corrosion of the small intestine together with massive bleeding and systemic toxicity (Systemic toxicity observed after ingestion of 30 g formic acid or more).

Accidental and suicidal oral exposure records report reversible burns of the oesophagus after ingestion of small quantities (up to 10g). Consumption of between 5 and 30 g of formic acid led to minor superficial oropharyngeal burns or more severe symptoms including abdominal pain, vomiting, dyspnea and dysphagia, hematemesis and pneumonitis, and esophageal strictures. Doses up to 45 g formic acid were survived by most patients. The majority of patients died after doses between 45 – 200 g formic acid. Reported symptoms at high doses were

	<p>corrosion of the gastro-intestinal tract, metabolic acidosis, haemolysis, loss of blood pressure, massive bleeding, hepatic and renal failure, and death.</p> <p>Inhalation exposure:</p> <p>Systemic effects are unlikely to occur. Workplace measurements showed mean values and 95% percentiles far below the threshold limit of 5 ppm or 9.5 mg/m³. Uptake of formic acid at this threshold exposure concentration equals approx. 0.5% of the metabolic rate observed in non-human primates. Therefore, an effect on the blood pH is unlikely. Formic acid inhalation concentrations from 30 ppm onwards are regarded as being immediately dangerous to life and health.</p> <p>One accidental inhalation exposure record reported reversible Pulmonary dysfunction in the form of Reactive Airway Dysfunction Syndrome. Suicidal inhalation exposure records (mixing of formic acid with concentrated sulphuric acid to form carbon monoxide) report death due to CO intoxication alongside corrosion/irritation of skin, trachea, lungs, stomach due to formic acid fumes.</p>
Justification for the conclusion	Workplace measurements, health records from industry, case reports

Data waiving	
Information requirement	Epidemiological studies on formic acid
Justification	None available

3.15 OTHER DATA

Summary table of other data			
Type of data/ report, Reliability	Test substance	Observations	Reference
Proposed acceptable residue levels	Residue definition: Group formic acid and ethyl formate	ADI 3 mg/kg bw/day	European Commission (2005) BPD ID A6.15.4_01a FA_BPR_Ann_II_8_16_1_01
	Residue definition: Group formic acid and ethyl formate	ADI 3 mg/kg bw/day	JECFA (2003) BPD ID A6.15.4_01b FA_BPR_Ann_II_8_16_1_01
	Formic acid, formate	No MRL set	EFSA (2009, 2014, 2015) FA_BPR_Ann_II_8_16_1_01 FA_BPR_Ann_II_8_16_2_0_JNS FA_BPR_Ann_II_8_16_3_0_JNS

When applied as recommended by the biocidal use patterns, no prolonged continuance of formic acid residues on treated surfaces is expected, owing to the volatility of formic acid and the water solubility of the acid and its salts. After uptake, formic acid and formate is readily and completely metabolised with the consequence that no relevant residue quantities are found in meat, milk, eggs, honey, or other products in addition to naturally occurring trace amounts which result from the fact that formic acid does naturally occur in food and plants. Hence, the formate consumer exposure is not increased through the diet.

As to the animal health, formic acid and formate salts (FORMI™ LHS, ammonium formate and sodium formate) showed a positive effect on the intestinal microflora which is beneficial for the treated animals. Therefore, formic acid and formate salts (FORMI™ LHS and sodium formate) were proposed as feed additives. Formic acid, FORMI™ LHS, and sodium formate are approved feed and drinking water additives, whereas ammonium formate was not approved because of the inevitable presence of formamide, a developmental toxicant, while formate was not considered to be problematic (EFSA, 2009; 2014, 2015; cf. outline further below).

The consumer average daily intake of formic acid with the natural food content was estimated to range between 0.1 to 0.43 mg/kg body weight.

Historically, higher intakes must be considered in those European countries where formic acid, or formate salts, was used as approved food preservative until 1998. A group ADI-value (Acceptable Daily Intake) of 3 mg/kg bw was established by JECFA for formic acid and ethyl formate in 1979 and maintained in 1997, and this value was adopted in the latest synoptic document of the EC updated in 2005.

Following ingestion formic acid distributes rapidly, and it is rapidly metabolised to CO₂. Further, it is required for the biosynthesis of purines and pyrimidines in the intermediary metabolism. In the case of unintentional uptake of residual product, no accumulation is expected as formic acid is rapidly removed from blood in all species that have been investigated.

Formic acid, FORMI™ LHS, and sodium formate are approved feed and drinking water additives, and their use in feed (up to 12,000 ppm for pigs, 10,000 ppm for birds, ruminants, and other species) and drinking water (4,000 ppm) as specified in the Scientific Opinions is considered to be safe for the animals, the consumer, and the environment, whereas users might need protective measures (PPE: skin, eye, respiratory protection) because of the corrosivity of formic acid at concentrations >10%. The EFSA panel (FEEDAP) does not expect relevant residue levels and did not propose a MRL value (EFSA, 2009; 2014, 2015).

Conclusion:

When applied as recommended by the biocidal use patterns, no considerable potential or actual exposure of formic acid to animals and /or humans through diet or other means is expected.

Summary of Scientific EFSA Opinions pertaining to formic acid and its salts			
	EFSA (2009) No. 1315	EFSA (2014) No. 3827	EFSA (2015) No. 4113
Reference No.	FA_BPR_Ann_II_8_16_1_01 FA_BPR_Ann_II_8_16_2_0_JNS FA_BPR_Ann_II_8_16_3_0_JNS		
Objective	Re-evaluation	Re-authorisation	Authorisation of new use
Legal basis of evaluation	Request from BASF SE to the EU Commission; technical dossier obtained directly from BASF SE	Request from ACIAC-EEIG consortium to the EU Commission; technical dossier obtained directly from the applicant.	Request from FEFANA/HYFAC to the EU Commission; technical dossier obtained directly from the applicant.

Trade name	FORMI™ LHS	Formic acid	Not appropriate
Chemical	Potassium diformate, min. 98%	Formic acid, min 84.5%	Formic acid min. 84.5% Ammonium formate; min 35%(liquid) Sodium formate min 98% (solid); min 15% (liquid)
Formula	(KCOOH*HCOOH)	HCOOH	HCOOH NH ₄ COOH NaCOOH
Contains	Formic acid, formate	Formic acid	Formic acid, formate salts
Intended use	Feed additive for sows. 0.8 – 1.2% in feed	Feed additive (pigs 1.2%, poultry 1%, ruminants 1%; all other species 1%) Drinking water 0.4%	Feed additive Formic acid: all species except pigs; 1% in feed Formic acid: pigs; 1.2 % in feed Ammonium formate: all species except pigs; 1% in feed Ammonium formate: pigs; 1.2 % in feed Sodium formate: all species except pigs; 1% in feed Sodium formate: pigs; 1.2 % in feed
Conclusions of safety evaluation	Safe at a max. dose of 1.2% in feed (12,000 ppm); MoS = 4	Safe doses: up to 1.2% in feed. No MoS identified.	Safe doses: up to 1.2% in feed. No MoS identified. Ammonium formate: unsafe, due to inevitable presence of formamide (developmental toxicant)
Livestock	Well tolerated by sows; no adverse effects up to 1.2% in feed.	Safe doses: Pig: 1.2% Poultry, ruminants: 1% Other species: 1% (extrapolation)	Safe doses: Pig: 1.2% (both formic acid and sodium formate) Poultry, ruminants: 1% (both formic acid and sodium formate)

			Other species: 1% (extrapolation; both formic acid and sodium formate)
user	FORMI LHS is an eye irritant. Requires protection measures.	Safe concentrations > 10% considered to be corrosive to skin and eyes. Volatile liquid. Inhalation exposure and exposure of skin and eyes present a risk for unprotected workers	Formic acid: cf. EFSA (2014) No. 3827 Sodium formate: mildly irritating to the skin. Safe handling may require PPE. Formic acid, sodium formate, ammonium formate were all considered to be skin sensitizers due to the lack of data (cf. remark 3 in last line)
consumer	Safe. No consumer formate exposure expected, due to rapid and complete metabolism in the pig.	Safe. No contribution to consumer exposure, due to rapid turnover and no accumulation	Safe ((both formic acid and sodium formate). No contribution to consumer exposure, due to rapid turnover and no accumulation
environment	Safe, when used as intended	Safe, when used as intended	Safe, when used as intended
microbiology	MIC values for Gram-positive and Gram-negative bacteria in the range 0.2-0.4%. No incidence of resistance to formic acid has been recorded until now.	MIC values not reported	MIC values mentioned but no details reported
Efficacy	Given at 1.2 % in feed	Recommended concentrations inhibit bacterial growth in feedingstuffs, drinking water, and in silage.	
MRLs (max. residue levels)	None defined. No negative effect on meat quality at proposed dose.	None defined. No negative effect on meat quality at proposed dose.	
Remark 1		ACIAC-EEIG consortium liquidated and rights transferred to FEFANA (includes Addcon Nordic SA; BASF	FEFANA/HYFAC members: Kemira Oyj; Perstorp AB; Selko feed Additives; Andres Pintaluba; BASF SE; Anitox Ltd.

		SE; Kemira Oyj Pestorp AB; Selko BV)	
Remark 2			Formic acid: conclusions from previous opinion reiterated.
Remark 3			<p>Formic acid, sodium formate, ammonium formate were all considered to be skin sensitizers due to the lack of data.</p> <p>It should be noted that formic acid was negative in a valid Buehler test, and that potassium formate was also negative in a valid assay. This result can be read across to sodium formate. Apparently, the applicants did not present data on this endpoint.</p>

Conclusion used in Risk Assessment – Other data

Conclusion	<p>An ADI has previously been set at 3 mg/kg bw/day.</p> <p>No further data on residues on the treated or contaminated food or feeding stuffs including kinetics of disappearance are needed.</p>
Justification for the conclusion	<p>When applied as recommended, neither prolonged remain of formic acid residues in food or feeding stuffs nor significant exposure to animal or human is expected, due to volatilisation, wash-off, and rapid and complete metabolism.</p> <p>The EFSA Feed additive panel (FEEDAP) shares this opinion and concludes the use of feed additives containing formic acid or formate salts is safe for the consumer, the animals, and the environment in three Scientific Opinions.</p>

4 ENVIRONMENTAL EFFECT ASSESSMENT

In aqueous solution and at neutral pH, formic acid and water-soluble formate salts dissociate and are present as the formate anion in solution. Based on this, it is deemed justified to include studies conducted with water-soluble formate salts in the evaluation of the environmental effects of formic acid.

4.1 FATE AND DISTRIBUTION IN THE ENVIRONMENT

4.1.1 Degradation

4.1.1.1 ABIOTIC DEGRADATION

4.1.1.1.1 Hydrolysis

The hydrolytic stability of formic acid at pH 4, 7 and 9 was investigated in a study following OECD 111, covering also Directive 92/69/EEC C.7 and US EPA OPPTS 835.2110.

The test item was dissolved in 50 mL of appropriate buffer solutions to give a final concentration of 400 mg a.i./L. The solutions were incubated at 50 °C and aliquots were taken after certain intervals and analysed in a modular HPLC system with UV/vis detector. After 5 days (120 h) the test was terminated since no hydrolysis was observed at any pH (preliminary test). At test end about 100 % recovery of the parent compound was reached at pH 4, 7, and 9

Conclusion:

Formic acid is considered to be hydrolytically stable, independent of the pH.

Summary table - Hydrolysis							
Method, Guideline, GLP status, Realibility	pH	Temp. [°C]	Initial TS concentration, Co[mol/l]	Half-life, DT ₅₀ [d]	Coefficient of correlation, r ²	Remarks	Reference
OECD TG 311; Directive 92/69/EEC, C.7; US EPA	4	49.9 ± 0.5 °C	8.7 mmol/L (400 mg a.s./L)	> 1 year	Not applicable	/	██████████ (2002) BPD ID A7.1.1.1.1_01 Doc IIIA JOINT: FA_BPR_Ann_II_10_1_1_1_a
	7						
	9						

OPPTS 835.2110 (Hydrolysis as a function of pH); GLP-study; Reliability 1							
--	--	--	--	--	--	--	--

Converted to environmentally relevant conditions (pH 7; 12 °C) the DT50 value becomes > 20.7 years (Guidance on BPR: Volume IV Environment Parts B+C (Version 2.0 October 2017), Equation 28).

Value used in Risk Assessment	
Value/conclusion	DT50 > 1 year (pH 4, 7 and 9; 49.9±0.5 °C) DT50 > 20.7 years (pH 7; 12 °C)
Justification for the value/conclusion	According to Guideline OECD 111 a substance is considered hydrolytically stable if, in the preliminary test at 50 °C, less than 10 % of hydrolysis is observed after 5 days. No additional testing is required at this point. Conversion of DT50 value to 12 °C using Equation 28 of the Guidance on BPR: Volume IV Environment Parts B+C (Version 2.0 October 2017), Equation 28.

4.1.1.1.2 Phototransformation in water

No new data was submitted for this endpoint, instead a justification for non-submission based on other available data (literature) was submitted (*Doc IIIA JOINT: FA_BPR_Ann_II_10_1_1_1_b*).

Direct photolysis

According to the HSDB database (available online at <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>) formic acid does not absorb at wavelengths > 290 nm and therefore is not expected to be susceptible to direct photolysis by sunlight.

Phototransformation with OH-radicals in water

From the literature (Buxton et al., 1988, BPD ID A7.1.1.1.2_01), a rate constant (k) for the reaction of formic acid and the formate ion with OH-radicals in water were compiled:

pH	Molecule	Rate constant (k) [L/mol*sec]
0.4 – 1.0	Formic acid (HCOOH)	4.4×10^5
7.0 – 13.5	Formate ion (HCOO ⁻)	2.1×10^8

In order to be able to derive half-lives from these data, hydroxyl-radical concentrations in water have to be assumed. This is also derived from literature (Zepp et al., 1987, BPD ID A7.1.1.1.2_02), wherein it is described that for the small lake Greifensee in Switzerland, the average OH-radical concentration over the whole water column (14 m) over the whole year is 3.0×10^{-18} mol/L. From this, a half-life for aquatic photolysis can be calculated for the formate ion, which is the relevant form of formic acid in water, of approximately 35 years (34,89 years).

Phototransformation with NO₃-radicals in water

At pH 5 – 9, the rate coefficients for the aqueous reactions of NO₃ with HCOO⁻ at 25 °C were experimentally determined to range from $4.7 \pm 0.6 \times 10^7$ to $5.0 \pm 0.4 \times 10^7$ L/mol*sec. With formic acid the rate constant was $3.3 \pm 0.4 \times 10^5$ L/mol.sec at pH 0.5 and 25 °C. The differences in reactivity of the anion HCOO⁻ compared to HCOOH were explained by the higher reactivity of NO₃ in the charge transfer processes compared to H-atom abstraction (Exner et al., 1994, BPD ID A7.1.1.1.2_03).

Transformation products

Formic acid is a simple C1-molecule which can be degraded chemically to innocuous substances in most environments.

Value used in Risk Assessment	
Value/conclusion	<ul style="list-style-type: none"> Direct photolysis: not expected Photo-oxidation with OH-radicals in water: DT₅₀ HCOO⁻ = 35 years
Justification for the value/conclusion	The information submitted by the applicant was deemed sufficient. Phototransformation will not likely play a role in the degradation of formic acid in the environment.

4.1.1.1.3 Estimated photo-oxidation in air

According to the HSDB database (available online at <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>) formic acid does not absorb at wavelengths > 290 nm and therefore is not expected to be susceptible to direct photolysis by sunlight.

The photo-degradation of formic acid in air was estimated through the modelling program AOP v1.91, included in the EPISUITE program developed by US EPA.

For a 12-hour day, with an OH-radical concentration of 1.5×10^6 OH/cm³, a half-life of 20.6 days or 493.7 hours was estimated.

For a 24-hour day, with an OH-radical concentration of 0.5×10^6 OH/cm³, a half-life of 30.9 days or 740.5 hours was estimated.

Summary table – Photo-oxidation in air					
Model	Light protection (yes/no)	Estimated daily (24h) OH concentration [OH/cm ³]	Overall OH rate constant [cm ³ /molecule sec]	Half-life [hr]	Reference
AOP v.1.91	/	0.5×10^6	5.2×10^{-13}	740.5	██████████ (2006) BPD ID A7.3.1_01 Doc IIIA JOINT: FA_BPR_Ann_II_10_3_1

Furthermore, according to §2.3.6.3 of the Guidance on the BPR: Volume IV Part B on photochemical reactions in the atmosphere, the pseudo-first order rate constant in air can be calculated using the following:

$$kdeg_{air} = k_{OH} \times OHCONC_{air} \times 24 \times 3600$$

$$\Leftrightarrow kdeg_{air} = 5.2 \cdot 10^{-13} \times 5 \cdot 10^5 \times 24 \times 3600$$

$$\Leftrightarrow kdeg_{air} = 0.0225d^{-1}$$

In a monograph on kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds (Atkinson., 1989, BPD ID A7.3.2_01), a unit-weighted average of the rate constants reported in different sources results in a recommended rate constant of 4.5×10^{-13} cm³/mol.sec for formic acid. From this, using the same formula as above, a degradation half-life of 35.7 days or 855.7 hours can be derived.

The latter derived half-life will be used for further risk assessment purposes, since it is more conservative than the half-life estimated through the AOP program.

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ = 855.7 hours Formic acid is only moderately subjected to photodegradation
Justification for the value/conclusion	The information submitted by the applicant was acceptable.

4.1.1.2 **BIOTIC DEGRADATION**

4.1.1.2.1 **Biodegradability (ready/inherent)**

Four studies are available on the aerobic biodegradation of formic acid in fresh water. All four tests document on ready biodegradability.

Two identical studies were performed by [REDACTED] (1988a/b) on formic acid, both using the modified OECD screening test (OECD 301E).

In both tests 20 mg DOC/L of test substance was inoculated with 0.5 ml effluent per litre medium (composition according to OECD). The mixture was aerated in the dark or diffuse light at room temperature ($22^{\circ}\text{C}\pm 2^{\circ}\text{C}$). A reference substance (sodium benzoate; 20 mg/l DOC) was tested in parallel. Both tests were performed in duplicate.

In the first test (BPD ID A7.1.1.2.1_01), samples were taken on day 0, 1, 7, 10, 13 and 14, while in the second test (BPD ID A7.1.1.2.1_02) samples were taken daily, to measure the DOC concentrations with an oxygen electrode.

In the first tests, 4 additional controls were run next to the test substance and reference substance: a control without test substance (blank), a control with reference substance, an abiotic control and a toxicity control. In the second test, the abiotic and toxicity control was omitted, which can be seen as a deficiency.

For the first test, 90-100% of the initial formic acid (20 mg/L DOC) was eliminated from water after 14 days. The 10-day window was reached.

For the second test, 99 % of the initial formic acid (20 mg/L DOC) was eliminated from water after 11 days, also reaching the 10-day window.

With these results, both tests indicate that formic acid is readily biodegradable.

The third and fourth ready biodegradability test are both closed bottle tests (OECD 301D) performed with potassium formate, for which the formate ion is representative for formic acid in water.

The oldest test ([REDACTED] 1992a, BPD ID A7.1.1.2.1_03) was performed according to the principles of GLP.

In this study the test substance and reference substance (sodium benzoate) were tested at respective concentrations of 18 and 3 mg/L. BOD bottles of 250 mL were filled with a standard nutrient medium, the test substance or reference substance and 1 drop/L of activated sewage sludge bacteria. Samples were taken after days 0, 5, 15 and 28 to measure the BOD with an oxygen electrode. Additionally, a blank control, an inoculum control and an inhibition control were run in parallel. The test was performed at 20 °C in a water bath and was performed in duplicate.

92 % of the initial test substance concentration was eliminated from water after 28 days.

Between day 5 (15 % degradation) and day 15 (90 % degradation) more than 60 % degradation related to ThOD was observed. The 14-day window was met.

The second closed bottle test (████████ 2000, BPD ID A7.1.1.2.1_04) confirmed the results of the first study, albeit not being GLP. In this study the test substance and reference substance (aniline) were tested at respective concentrations of 20 and 1.95 mg/L. The preparation of the BOD bottles was identical to that in the first test and samples were taken at appropriate intervals (days 0, 2, 5, 7, 9, 12, 14, 16, 22 and 28) to measure the BOD with an oxygen electrode. Additional controls, such as in the first test, were run in parallel.

82 % of the initial test substance concentration was eliminated from water after 28 days.

Between day 2 (10 % degradation) and day 9 (75 % degradation) more than 60 % biodegradation related to ThOD was observed. The 14-day window was met.

Conclusion:

Overall, considering the 4 ready biodegradability tests performed with the active substance, it can be concluded that Formic Acid is readily biodegradable.

Further screening tests on inherent biodegradability are deemed unnecessary (applicant justification *Doc IIIA JOINT: FA_BPR_Ann_II_10_1_1_2_b*).

Summary table - biodegradation studies (ready/inherent)											
Method, Guideline, GLP status, Realibility	Test type ¹	Test parameter	Inoculum			Additional substrate	Test substance concentration	Degradation		Remarks	Reference
			Type	Concentration	Adaptation			Incubation period	Degree [%]		
Modified OECD Screening Test, 79/831/EE C, Annex V, C3;	Ready	DOC	Effluent municipal STP (lab. culture)	0.5 mL (total batch volume: 900 mL)	no	no	Formic acid; 20 mg DOC/L	28 (terminated after day 14)	90-100 10-day window passed		████████ 1988a BPD ID A7.1.1.2.1_01 <i>Doc IIIA JOINT: FA_BPR_Ann_II_10_1_1_2_a_1</i>

non-GLP study, Reliability 2											
Modified OECD Screening Test, 79/831/EEC, Annex V, C3; non-GLP, Reliability 2	Ready	DOC	Effluent municipal STP (lab. culture)	0.5 mL (total batch volume: 900 mL)	no	no	Formic acid; 20 mg DOC/L	28 (terminated after day 11)	99	10-day window passed	██████████ 1988b BPD ID A7.1.1.2.1_02 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_1_2_a_2</i>
Closed Bottle Test, OECD TG 301D, GLP Reliability 1	Ready	BOD	Activated sewage sludge of municipal STP	1 drop/L	no	Nutrient medium	Potassium formate; 18 mg/L	28 (90% removal after 15 days)	92	14-day window passed	██████████ 1992a BPD ID A7.1.1.2.1_03 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_1_2_a_3</i>
Closed Bottle Test, OECD TG 301D, non-GLP, Reliability 2	Ready	BOD	Activated sludge cultivated on synth. sewage; supplied w. domes. sewage	6.8*10 ⁵ CFU/L (heterotrophic bacteria)	no	Nutrient medium	Potassium formate; 20 mg/L	28 (75% removal after 9 days)	82	14-day window passed	██████████ 2000 BPD ID A7.1.1.2.1_04 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_1_2_a_4</i>

			5 d prior to start								
¹ Test on inherent or ready biodegradability according to OECD criteria											

Value used in Risk Assessment	
Value/conclusion	Ready biodegradable (meeting the 10 or 14-day window)
Justification for the value/conclusion	Based on the available studies, formic acid is well within the pass levels of 70 % DOC and 60 % ThOD removal. The 10-day or 14-day window (depending on test-type) is met each time.

4.1.1.3 **RATE AND ROUTE OF DEGRADATION INCLUDING IDENTIFICATION OF METABOLITES AND DEGRADATION PRODUCTS**

4.1.1.3.1 **Biological sewage treatment**

4.1.1.3.1.1 *Aerobic biodegradation*

Data waiving	
Information requirement	A justification of non-submission of data was submitted by the applicant (<i>Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_1_a</i>), based on the fact that such a test is not a core data requirement and that submitted studies showed formic acid to be readily biodegradable.
Justification	Justification is accepted.

4.1.1.3.1.2 *Anaerobic biodegradation*

A study on the acclimation and degradation of petrochemical wastewater components by methane fermentation was submitted for this data point (Chou et al., 1979, BPD ID A7.1.2.1.2_01). The study dates from 1979 and does not follow a known guideline or is performed according to GLP.

Hungate serum bottles were filled with water and displaced with an inert gas mixture of CO₂ and CH₄. A 50 mL inoculum of acetate enriched cultures (1000 mg/L SS, laboratory culture of domestic sludge fed with acetate for years) was injected into the bottle, together with 100 mL of acetate and 25 mg of test substance (formic acid, amongst others).

Gas production was monitored and test substance was injected with a microliter syringe as needed.

For formic acid, the test showed 89 % of substrate removal after a lag time of 4 days. An overall degradation rate of 286 mg/L.day was established.

Summary table - STP anaerobic biodegradation							
			Inoculum			Degradation	Reference

Method, Guideline, GLP status, Reliability	Test type ¹	Test parameter	Type	Concentration	Adaptation	Additional substrate	Test substance concentration	Incubation period	Degree [%]	Remarks	
No guideline (Hungate serum bottle) Non-GLP Reliability 4	no guideline (anaerobic)	CH ₄ evolution	Acetate enriched cultures (lab. cult. domestic sludge)	1000 mg /L SS	no	Acetate	Formic acid; 500-1000 mg/L (renewed)	unknown (up to 30 days)	89		Chou et al., 1979 BPD ID A7.1.2.1.2_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_3_1_b</i>
¹ Test according to OECD criteria											

The BE eCA assigns a reliability of 4 to this test, since the test report contains insufficient details. Therefore the results of this test can only be considered as indicative.

The test did not follow an official guideline and contains insufficient details in order to assess whether it could be compared to one.

The applicant was asked if they could provide further information, but they could not and accepted the reliability of 4 assigned by the BE eCA.

Since the anaerobic biodegradation is not a strict data-requirement, further testing was not deemed necessary.

Value used in Risk Assessment	
Value/conclusion	Indication that anaerobic degradation may be possible
Justification for the value/conclusion	Test report contains insufficient details, does not follow a known guideline and was not performed according to GLP. Since this endpoint is not strictly a data requirement, no new testing is required at this point.

4.1.1.3.1.3 STP simulation test

Data waiving	
Information requirement	A justification of non-submission of data was submitted by the applicant (<i>Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_1_a</i>), based on the fact that such a test is not a core data requirement and that other submitted studies showed formic acid to be ready biodegradable.
Justification	Justification is accepted.

4.1.1.3.2 Biodegradation in freshwater

4.1.1.3.2.1 Aerobic aquatic degradation

Data waiving	
Information requirement	A justification of non-submission of data was submitted by the applicant (<i>Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_2_a</i>), based on the fact that such a test is not a core data requirement and that other submitted studies showed formic acid to be ready biodegradable.
Justification	Justification is accepted.

4.1.1.3.2.2 Water/sediment degradation test

Data waiving	
Information requirement	A justification of non-submission of data was submitted by the applicant (<i>Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_2_b</i>), based on the fact that such a test is not a core data requirement and that other submitted studies showed formic acid to be ready biodegradable.
Justification	Justification is accepted.

4.1.1.3.3 Biodegradation in seawater

4.1.1.3.3.1 Seawater degradation study

One test to assess the biodegradability in seawater was submitted (██████████ 1994, BPD ID A7.1.1.2.3_01) (*Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_3*). The test was supposedly performed according to OECD guideline 306 and according to the GLP principles, using potassium formate liquor (i.e. potassium formate 75% in water) as test material.

In a closed bottle test, potassium formate liquor was tested at a concentration of 15 mg/L. Sodium acetate was used as a reference substance. The inoculum was a non-specific mixture of marine microbiota, collected in the field.

The percentage biodegradation was determined by comparing the oxygen depletion value (BOD) with the corresponding Theoretical Oxygen Demand (ThOD), which was calculated as 143 mg O₂/g potassium formate liquor. Samples for oxygen analysis were taken at day 0, 7, 14, 21 and 28.

The test concludes that after 28 days 71.3 % of the initial test substance concentration was eliminated. The 60 % mark was reached between days 0 and 7, with 61.5 % degradation at day 7.

BE eCA is however of the opinion that the test report for this test is severely lacking in details. It is unclear what the exact empirical formula of the test material is to arrive at the calculated ThOD of 143 mg/g. Nor are details on for example the number of repetitions, whether or not a blank control was tested, the reason why a larger concentration than the concentration range suggested in the guideline determinable from the original test report. Merely a statement that the test was performed according to OECD 306 seems insufficiently reliable.

Therefore, BE eCA assigns a reliability 4 to this test, which render its result unusable for further risk assessment purposes.

The applicant was asked if they had any more information on this particular study, but the answer thus far was negative and the applicant accepted the reliability assessment made by BE eCA.

Since this endpoint is not a core data requirement, new testing is not required at this time.

Value used in Risk Assessment	
Value/conclusion	No value from this test is retained for the risk assessment
Justification for the value/conclusion	The test report was deemed too summarily to retain the results as a key value. However, at this point, no further testing is required on the basis that such a data point is not a core data.

4.1.1.3.4 Higher tier degradation studies in water or sediment

No available data.

4.1.1.3.5 Biodegradation during manure storage

A study on the characteristics of volatile fatty acids in stored dairy manure before and after anaerobic digestion (Page et al., 2014, Doc IIIA FA_BPR_Ann_II_10_1_3_4) and a study on changes in swine manure during anaerobic digestion (Iannotti et al., 1979, Doc IIIA FA_BPR_Ann_II_10_1_3_4_Iannotti_1979) are submitted for this data point.

In Page et al. (2014), raw dairy manure and raw dairy manure amended with pre-consumer waste were incubated in reactors without aeration and stirring of the manure; thus simulating storage conditions of manure. Formic acid was not added to the manure samples, but the course of the naturally occurring formic acid was monitored over a period of 100 days at 20 °C. The two types of manure were incubated in duplicate reactors. The reactors were sampled every seven days from the top and the bottom layer. The top layer represents aerobic conditions, while the bottom layer is characterized by anaerobic conditions.

In both manure types the degradation of formic acid could be observed. However, there were also phases where the concentration of formic acid was increasing. These fluctuations can be explained by the degradation of other volatile fatty acids and/or other organic substances, which can lead to the formation of formic acid. Over the last 3 to 5 weeks either formic acid was no longer formed or the degradation activity was equal to the formation rate of formic acid as the observed concentrations were at 0 mg/L.

The study shows that formic acid is degraded under aerobic and anaerobic conditions in manure samples (raw dairy manure and amended dairy manure). Based on the graphical representation of the concentration trends, a DT₅₀ for the aerobic top layer of ≤ 7 days and ≤ 10.5 days for the anaerobic bottom layer can be derived for wet manure storage.

Iannotti et al. (1979) investigated changes in swine manure during anaerobic digestion. Swine manure was digested in pilot-size digesters (0.42 m³) which had been in operation for one year. The loading rate was 3.78 g volatile solids (VS)/L/d. The influent waste was from finishing hogs. The digester temperature was 35 °C. The detention time was 15 days.

The digester was fed swine manure with a total of 4.7±0.6 g/d (= influent). Based on an influent volume of 28.2 L, this results in a concentration of 167 mg/L of formic acid in the swine manure. In the effluent no formic acid was detected which is a removal of 100%. Based on the complete removal of formic acid from the influent and its retention time in the digester, a conservative DT₅₀ of 7.5 days can be deduced.

Summary table – Biodegradation during manure storage

			Inoculum			Degradation		Reference
--	--	--	----------	--	--	-------------	--	-----------

Method, Guideline, GLP status, Reliability	Test type ¹	Test parameter	Type	Concentration	Adaptation	Additional substrate	Test substance concentration.	Incubation period	Degree [%]	Remarks	
No guideline ² (reactors without aeration and stirring) Non-GLP Reliability 2	no harmonised guideline available	Concentration of formic acid	Raw dairy manure: R1 & R2 Dairy manure (90.1%) mixed with blood (5.9%) and trap (4.0%): R3 & R4	N/A	N/A	N/A	Formic acid naturally present in manure samples. R1 & R2: < 850 mg/L R3 & R4: ≤ 27 100 mg/L	98 days (at 20 °C)	100 (measured concentration of 0 mg/L)	N/A	Page et al., 2014 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_3_4</i>
No guideline ² (anaerobic pilot-size)	no harmonised guideline available	Concentration of formic acid	Swine manure from finishing hogs	N/A	N/A	N/A	Formic acid naturally present in manure samples	22 weeks (at 35 °C)	100	N/A	Iannotti et al., 1979 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_10_1_3_4_Iannotti_1979</i>

digester) Non-GLP Reliability 2							: 167 mg/L				
¹ Test according to OECD criteria ² No harmonised guideline available											

Page et al. (2014) is selected as key study. Justification on the use of this study as key study is provided in *Doc IIIA JOINT: FA_BPR_Ann_II_10_1_3_4*.

At ENV WG-I-2022, it was agreed that the DT50 of 10.5 days (at 20°C), derived based on cattle manure can be used also for all other animal categories. This DT50 value is derived based on cattle manure (Page et al., 2014) and confirmed for pigs manure (Iannotti et al., 1979). No data is available for poultry manure. However, poultry manure has another consistency compared to cattle and pigs manure and is much dryer. Degradation in such kind of manure tend to be aerobic, in which case the DT50 values are expected to be covered by the DT50 value of 10.5 days (20 °C). Indeed, according to OECD ESD No. 14 (OECD, 2006), DT50 values for degradation in soil can be used as a surrogate for degradation in manure when no other data are available. In the case of poultry manure (aerobic degradation) this would yield a DT50 value of 1 day (see section 4.1.1.3.6 of the CAR).

The swift anaerobic degradation of formic acid in manure is not surprising. Formic acid is the simplest carboxylic acid and is a natural compound occurring at significant concentrations in all environmental compartments (please refer to section 4.3 of the CAR). Several lines of evidence are available to confirm anaerobic degradation. Section 4.1.1.3.1.2 of the CAR (STP anaerobic degradation) contains the study of Chou et al. (1979). Although the publication was rated a reliability of 4, the data indicate that anaerobic degradation of formic acid occurs. In the publication of Page et al. (2014) is stated that methanogens can directly use formic acid. The publication of H ellsten et al. (2005b, see next section), studying aerobic and anaerobic degradation of formate in soil at low temperatures, concludes that [...] *there is a potential for swift aerobic and anaerobic biodegradation of formate in the subsurface of the study site, which is hardly surprising as formate can be utilized by a wide variety of aerobic, facultative, and anaerobic microorganisms.*

Value used in Risk Assessment	
Value/conclusion	DT ₅₀ for biodegradation in manure: ≤ 10.5 days (20 °C) DT ₅₀ for biodegradation in manure: ≤ 19.9 days (12 °C)
Justification for the value/conclusion	Value derived from the graphical representation of concentration trends of formic acid in manure at anaerobic conditions (bottom of reactor). Value agreed at ENV WG-I-2022.

4.1.1.3.6 Biotic degradation in soil

According to the BPR, all tests on fate and behaviour are not part of the core data set. Requirements for such tests only come into play when there is exposure to soil.

For this dossier, the applicant waived all data referred to by BPR Annex II point 10.2. Since no direct exposure to soil is expected from the intended uses of formic acid in PTs 2, 3, 4, 5 and 6, and since formic acid is readily biodegradable, this waiving is accepted.

The applicant submitted nevertheless 3 open literature studies providing indication of rapid biodegradation of formic acid in soil.

- Lissner et al., 2014: Doc IIIA FA_BPR_Ann_II_10_2_a ;
- Hellstén et al., 2005a: Doc IIIA FA_BPR_Ann_II_10_2_b ;
- Hellstén et al., 2005b: Doc IIIA FA_BPR_Ann_II_10_2_c ;
- Glanville et al., 2012 : Doc IIIA FA_BPR_Ann_II_10_2_d

Lissner et al. (2014) is a lysimeter experiment following the degradation of potassium formate executed in Norway. Formate is added to all of the lysimeters together with propylene glycol (PG) as part of a deicing solution in a ratio of 70 g/m² formate and 350 g/m² PG. Due to the presence of PG and the uncertainty to what extent this interferes with the natural fate and behaviour of formate in soil, this study is assigned a reliability of 3.

In Hellstén et al. (2005a), potassium formate was applied to the soil surface of a lysimeter in Finland. Application took place five times (0,68 kg/m² per application) during winter on the snow cover of a lysimeter. The lysimeters were composed of well-graded sand and gravel. The mean formate concentration entering the soil was calculated at 2730 mg/L. The percolated water was collected at 12 dates and analyzed for formate, CO₂, TOC, COD, and other parameters.

The objective of this study was to examine the migration and degradation of potassium formate in the unsaturated zone of a lysimeter in a sandy aquifer in real winter and spring conditions.

The study concluded that formate was effectively removed in a sandy lysimeter after a cold winter period. The disappearance of formate was accompanied by the formation of carbon dioxide and bicarbonate in the percolating water indicating biodegradation of formate.

Hellstén et al. (2005b) investigated the degradability of sodium and potassium formate in soil under aerobic and anaerobic conditions in a set of microcosm experiments using radiolabeled sodium formate. Formate was shown to degrade under aerobic and anaerobic conditions from soil samples (top and subsurface). Given the differences in organic matter content, soil samples at different depths could be considered as

different soil types. Based on the graphical representation of the degradation data, a degradation half-life (DT_{50}) of < 1 day could be derived for all soil samples at temperatures of + 1 and +6 °C.

Glanville et al. (2012) investigated the overall relationship between laboratory-field and inter-annual field studies for mineralization of low molecular weight substrates in soil solution. Soil samples were spiked with ^{14}C -labelled compounds, formic acid being one of the substances. The soil samples were taken from freely draining agricultural grassland from a hyper-oceanic climatic region in North Wales (UK) at a soil depth of 10 cm. Sampling was done in 2009 and 2010. The half-life of formic acid was determined under lab and field conditions to be ≤ 1 day. This value was read from the graphs of the paper.

Summary table – Aerobic biodegradation in soil – laboratory study											
Method, Guideline, GLP status, Reliability	Test type ¹	Test parameter	Test system				Test substance concentr.	Incubation period	Degradation DT_{50}	Remarks	Reference
			Soil origin	Soil type	pH	OM %					
No guideline (microcosm experiments using radiolabelled sodium formate), Non-GLP, Reliability 2	Aerobic mineralisation in soil (no guideline, public literature data)	$^{14}CO_2$ (% of added ^{14}C -formate)	Kauriansalmi study site, Finland. Soil samples taken at various depths.	sandy and gravelly deposits with occasional thin layers of silt	not specified	0.43 (70-80 cm); 5.4 (5-15 cm); 0.70 (50-60 cm); 0.3	Concentration: 100 000 DPM of ^{14}C -formate (Radio-labelled sodium formate (^{14}C): 56 mCi/mmol; Source: Sigma)	up to 45 days	< 1 day for all soil samples at +1 and +6 °C	tested at low temperatures and for various soil depths; two replicates and one blank per sample	Hellstén et al., 2005b <i>Doc IIIA JOINT : FA_BPR_Ann_II_10_2_c</i>

						3 (10 0- 110 cm)					
No guideline (microcosm experiments using radiolabelled formic acid), Non-GLP, Reliability 2	Aerobic mineralisation in soil (no guideline, public literature data)	¹⁴ CO ₂	Abergwyngreyn, Gwynedd, North Wales (53°14'N, 4°1'W)	sandy clay loam (rhizosphere soil)	4.77-5.35	7.3 7- 7.9 7	Formic acid, ¹⁴ C-labelled (Source: Sigma-Aldrich Company Ltd., UK): < 10 nM formic acid	168 h	< 1 day (20 °C)	Experiments were performed in triplicate and in two subsequent years (2009 and 2010).	Glanville et al., 2021 <i>Doc IIIA JOINT : FA_BPR_Ann_II_10_2_d</i>
¹ Test according to OECD criteria											

Summary table – Field dissipation studies

Method, Guideline, GLP status, Reliability	Site	Application rate	Surface	Soil type	Soil texture	Test duration	Degradation DT ₅₀	Degradation DT ₉₀	Remarks	Reference
No guideline (lysimeter experiment with potassium formate),	Oslo airport, Norway	70 g/m ² potassium formate	not specified	Soil 1	silty and sandy deposits with low clay	2 years	not determined	not determined	Potassium formate was applied as part of a deicing solution with	Lissner et al., 2014 <i>Doc IIIA JOINT : FA_BPR_Ann_II_10_2_a</i>

Non-GLP, Reliability 3					content				polypropylene glycol.	
No guideline (lysimeter experiment with potassium formate), Non-GLP, Reliability 2	Southwestern Finland (Oripää lysimeter station, 60°55' N, 22°44' E)	Total potassium formate loading: 3.4 kg/m ² Substance applied by sprinkler irrigation over surface of one of the snow-covered lysimeter in five stages (0.68 kg/m ² per application) between 19 Dec. 2001 and 04 March 2002	Surface covered with local vegetation	Soil 2	well-graded sand and gravel	7 months	not determined	not determined	Experiment conducted in cold climate conditions.	Hellstén et al., 2005a <i>Doc IIIA Joint : FA_BPR_Ann_II_10_2_b</i>
No guideline (polypropylene cylinders (6.1 cm ² soil surface) placed 2 cm deep in soil),	Abergwyngregyn, Gwynedd, North Wales (53°14'N, 4°1'W)	450 µL soil solution spiked with 50 µL ¹⁴ C-formic acid (< 10 nM	The vegetation at the site consists of perennia	Soil 3 (freely draining agricultural grassland from a	sandy clay loam	168 h	< 1 day (13.8-17 °C)	not determined	Experiments were performed in triplicate and in two subsequent	Glanville et al., 2021 <i>Doc IIIA JOINT : FA_BPR_Ann_II_10_2_d</i>

Non-GLP, Reliability 2		formic acid)	l rye grass (Lolium perenne L.) and white clover (Trifolium repens L.) and is subject to intensive sheep grazing (>5 ewe ha ⁻¹) and receives regular fertilizer addition (120 kg N ha ⁻¹ y ⁻¹)	hyper-oceanic climatic region)					years (2009 and 2010).	
---------------------------	--	--------------	---	--------------------------------	--	--	--	--	------------------------	--

Based on the overall evidence available in public literature, it can be concluded that formic acid is expected to rapidly biodegrade in soil, even in sub-optimal conditions (low temperatures), and a DT50 for biodegradation in soil of < 1 day can be derived from the available data. Furthermore, it should be noted that in both Hellstén et al. (2005b) and Glanville et al. (2012), mineralisation was measured, meaning that the DT50 for biodegradation might be even more rapid.

Formic acid is the simplest carboxylic acid and is a natural compound occurring at significant concentrations in all environmental compartments (please refer to section 4.3 of the CAR), and can be utilized by a wide variety of aerobic, facultative, and anaerobic microorganisms (Hellstén et al. (2005b)).

None of the studies fulfil all conditions of the Guidance on the BPR: Volume IV Part A (version 1.2 May 2018), section 1.2 paragraph 12 specifying the conditions for public literature data to be considered as key studies. However, given the fact that:

- Hellstén et al. (2005b) and Glanville et al. (2012) use radiolabeled test material from a well-defined source for which a high purity can be assumed;
- the reference specification of formic acid doesn't contain relevant impurities;
- Hellstén et al. (2005b) investigated biodegradation in different soil layers with different organic matter contents, which could be considered as different soil types;

it was agreed at ENV WG-I-2022 to consider Hellstén et al. (2005b) and Glanville et al. (2012) as key studies and to use a DT50 value for soil of 1 day at 12 °C for the exposure assessment.

Value used in Risk Assessment	
Value/conclusion	DT50 value for soil of 1 day at 12 °C
Justification for the value/conclusion	Value agreed at ENV WG-I-2022.

4.1.2 Distribution

4.1.2.1 ADSORPTION ONTO/DESORPTION FROM SOILS

The adsorption coefficient (Koc) on soil and sewage sludge of formic acid was investigated in a HPLC screening test following OECD 121. The method of analysis was a modular HPLC system with UV/VIS detector under isocratic conditions (██████████ 2002, BPD ID A7.1.1.1.1_01) (*Doc IIIA JOINT: FA_BPR_Ann_II_10_1_2*).

Ten reference compounds were used for the calibration graph. Small amounts were dissolved in 30 vol% acetonitrile (ultrasonic treatment) and the flasks were made up to volume with water. The dead time (t_0) of the HPLC system was measured with formamide. Measurements of the retention times of the reference substances and of formic acid were performed in duplicate at 23 °C.

As formic acid is an ionisable substance with a pKa of 3.70 (Dolich 2007, BPD ID A3_01), two tests were performed with both non-ionised and ionised forms in appropriate buffer solutions (pH 4 and 10). The test item was dissolved in water/acetonitrile (9:1, v/v).

In the test run with the non-ionised formic acid (acidic conditions) the mean retention time (2.1 min) was shorter than the lower limit of the reference interval (acetanilide, 3.5 min) and shorter than the dead time established with formamide (2.2 min). Normally, the OECD test guideline indicates that if the log Koc of the test substance falls outside the calibration interval, the test should be repeated using more appropriate reference substances. However, in this case, the retention time of formic acid is also below the dead time, determined by using a substance (formamide) that does not react with the column, and thus does not have a tendency to adsorb. Knowing this, it can be concluded that formic acid also does not have a tendency to adsorb. For risk assessment purposes, the log Koc could be set to be smaller than that of the lower limit of the reference interval, being 1.25 for acetanilide.

In the test run with the ionised molecule under basic conditions (formate ion) there were no results on retention time at the end of the test, meaning that its retention time is longer than the upper limit of the reference interval (methiocarb, 9.1 min). In this case, the log Koc of formate is higher than 3.10. Sorption of the ionised form of formic acid is thus stronger than that of the non-ionised form, and the log Koc of formic acid of therefore depends on the pH.

It should be noted that the HPLC screening method is not suitable for the estimation of the Koc of formic acid. OECD Test Guideline No. 121 "Estimation of the Adsorption Coefficient (Koc) on Soil and on Sewage Sludge using High Performance Liquid Chromatography (HPLC)" states that the method may not work for moderate organic acids and that only log Koc values ranging from 1.5 to 5.0 can be determined. As both forms of formic acid, the protonated as well as the unprotonated, are not in the time range of the calibrated substances, no further conclusions can be derived.

In addition to this HPLC-method provided by the applicant, BE eCA used the screening programme EPI Suite 4.1 to estimate the Koc based on the structure of formic acid. KOCWIN v2.00, a subprogram included into EPI Suite to estimate the Koc, uses two different models to make an estimation. On the one hand, the Sabljic molecular connectivity method (MCI), estimates a Koc for formic acid of 1 L/kg (log Koc = 0). On the other hand, the program calculates the Koc based on the log Kow. When using the programs default log Kow of -0.54 (experimental database),

a Koc of 0.7195 L/kg is calculated (log Koc = -0.143). The applicant also submitted a study in which a log Kow is experimentally derived (██████████ 2002, BPD ID A7.1.1.1.1_01). When using this log Kow of -2.1 (pH 7), the program calculated a Koc of 0.09866 L/kg (log Koc = -1.00586).

However, given the pKa 3.70 for formic acid, the environmental relevant species is not formic acid but the formate ion. Franco et al. (2009) developed a method to estimate the Koc of monovalent organic acids and bases. The regression considers pH-dependent speciation and species-specific partition coefficients, calculated from the dissociation constant (pKa) and the octanol–water partition coefficient of the neutral molecule (log P_n). The pH-dependent estimation of Koc is provided by the following equation:

$$K_{OC} = \frac{10^{0.54 \cdot \log P_n + 0.11}}{1 + 10^{(pH_{soil} - 0.6 - pK_a)}} + \frac{10^{0.11 \cdot \log P_{ion} + 1.54}}{1 + 10^{(pK_a - pH_{soil} + 0.6)}}$$

where pK_a is the dissociation constant; log P_n the octanol-water partition coefficient of the neutral molecule; and pH_{soil} the pH of the soil.

(note: the equation contains a typo error in the second term: log P_{ion} should be log P_n)

No pH_{soil} is defined in Table 3 (Definition of the standard environmental characteristics) of the Guidance on BPR Volume IV Parts B+C (v2.0 October 2017). Therefore a neutral pH of 7 is assumed.

Provided a pK_a of 3.7, a log P_n of -0.54 (derived from the EPI Suite experimental database, see above)⁷ and a pH_{soil} of 7, a Koc of 30 (log Koc of 1.48) is yielded.

Conclusion:

The HPLC-method to estimate the Koc for formic acid resulted in an indication that the log Koc for formic acid will be below 1.25 and may vary with pH. The results obtained with KOCWIN, a programme to estimate the Koc, was also in line with the results obtained from the HPLC-method.

A theoretical log Koc of 0 (Koc = 1 L/kg) was estimated for formic acid.

⁷ A note regarding the log Kow used in the model of Franco et al.: log P_n in the model is the octanol–water partition coefficient of the neutral molecule, which is estimated to be -0.54 based on the EPI Suite experimental database. The experimentally derived log Kow of -2.1 is determined at a pH of 7, and can therefore not be used in the model because, given a pKa of 3.7, at that pH the predominant species is the ionized molecule (formate).

However, for risk assessment purposes, the environmental relevant species is not formic acid but formate. The method of Franco et al. (2009) was used to estimate a pH-dependent Koc and yielded a slightly higher log Koc of 1.48 to be used for risk assessment purposes assuming a soil with a neutral pH of 7.

Value used in Risk Assessment	
Value/conclusion	log Koc = 1.48 (for a soil with a neutral pH of 7)
Justification for the value/conclusion	Based on the method of Franco et al. (2009) and in line with the results obtained through the HPLC-method and calculations through EPI Suite, this theoretical value is deemed acceptable and no further tests in soil are required at this point.

4.1.2.2 **HIGHER TIER SOIL ADSORPTION STUDIES**

No available data.

4.1.3 Bioaccumulation

4.1.3.1 **MEASURED AQUATIC BIOCONCENTRATION**

Data waiving	
Information requirement	No experimental value is available. The applicant did not submit a justification for non-submission, however the BPR Annex II states that experimental determination may not be necessary if it can be demonstrated on the basis of physico-chemical properties (e.g. log Kow < 3) or other evidence that the substance has a low potential for bioconcentration. This statement is repeated in the Guidance on BPR: Volume IV. Part A, Chapter II: Requirement for Active Substances, §9.1. This exemption of submission of experimental data is the case for formic acid, since the experimental log Kow is well below the cut-off value of 3 (log Kow = -2.10, pH7).
Justification	The applicant did not submit a justification, but based on the guidance/legislation quoted above, no further justification from the applicant is required.

4.1.3.2 **ESTIMATED AQUATIC BIOCONCENTRATION**

To estimate the accumulation of formic acid in aquatic organisms, the applicant submitted an estimation using BCFWIN v.2.17 (██████████ 2007, BPD ID A7.4.2_01), which is an estimation program included in EPA's EPISUITE. Using this model, the bioconcentration of formic acid in aquatic organisms is estimated based on the experimental log Kow of -2.1 (derived for pH 7 or mean for measured log Kow at pH 5, 7 and 9) (██████████ 2002, BPD ID A7.1.1.1.1_01). Since the log Kow is below 1, the program assigns a default log BCF of 0.5 (BCF = 3.162 L/kg_{wwt}) and does not calculate a specific BCF for formic acid. However, this value indicates that formic acid is not expected to bioaccumulate in aquatic organisms, which is in accordance with the hydrophilic nature of formic acid, as well as with the log Kow being smaller than 3.

Additionally, BE eCA also calculated the BCF from the log Kow, according to the linear relationship developed by Veith et al. for substances with a log Kow between 2 and 6; and which is included in the Guidance Volume IV part B as equation 74:

$$\log BCF_{fish} = 0.85 \times \log Kow - 0.70$$

With this equation a log BCF of -2.48 is calculated (BCF = 0.00327 L/kg_{wwt}).

Summary table – Estimated aquatic bioconcentration					
Basis for estimation	Log Kow (measured)	Estimated BCF for fish (freshwater) [L/kg _{wwt}]	Estimated BCF for fish eating bird/predator	Remarks	Reference
BCFWIN v2.17 (reproduced in BCFBAF v3.01)	-2.1	3.162	/	since log Kow is below 1, the program reverts to a default log BCF of 0.5 (BCF = 3.162)	██████████ 2007 BPD ID A7.4.2_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_4_1</i>
BPR guidance Volume IV, Part B, eq.74	-2.1	0.00327	/	/	/

Value used in Risk Assessment	
Value/conclusion	The different estimated methods concur that formic acid will have a low potential to bioaccumulate, which is in line with the hydrophilic nature of formic acid and its log Kow being below 3.
Justification for the value/conclusion	/

4.1.3.3 **MEASURED TERRESTRIAL BIOCONCENTRATION**

Data waiving	
Information requirement	No experimental value is available and the applicant submitted a justification for non-submission (<i>Doc IIIA JOINT: FA_BPR_Ann_II_9_6</i>), stating the low log Kow (-2.1) as indication of formic acid's low potential to bioaccumulate.
Justification	Justification is acceptable and no experimental test is required.

4.1.3.4 **ESTIMATED TERRESTRIAL BIOCONCENTRATION**

The applicant did not submit an estimation for the terrestrial bioconcentration. BE eCA made its own calculations based on the available guidance.

According to the BPR Guidance Volume IV, Part B; bioconcentration can be described as a hydrophobic partitioning between the pore water and the phases inside the organism. It can be modelled according to the equation described by Jager (1998):

$$BCF_{earthworm} = \frac{(0.84 + 0.012K_{ow})}{RHO_{earthworm}}$$

The log K_{ow} for formic acid was experimentally determined as -2.1, giving a K_{ow} of 0.0079 L/kg. $RHO_{earthworm}$ is set by default on a value of 1 kg_{wwt}/L.

This gives a $BCF_{earthworm}$ of 0.84 L/kg_{wwt}, which indeed indicates a low potential of formic acid for bioaccumulation.

Value used in Risk Assessment	
Value/conclusion	Using the equation proposed in the BPR guidance, a $BCF_{earthworm}$ of 0.84 L/kg _{wwt} is determined
Justification for the value/conclusion	/

4.1.4 **Monitoring data**

No available data.

4.2 EFFECTS ON ENVIRONMENTAL ORGANISMS

4.2.1 Atmosphere

The vapour pressure of 42.71 hPa (20 °C; ██████████ BPD ID A3_01) and the Henry's Law Constant of 0.16 Pa.m³/mol (20 °C; ECT Oekotoxicologie GmbH; BPD ID A3_11) indicate low to moderate potential for volatilization and evaporation from water and wet surfaces. The potential of formic acid to be degraded by photo-oxidation in air is moderate with an estimated half-life of 855.7 hours (cfr. §4.1.1.1.3. above).

Besides the anthropogenic sources of emission, formic acid and formate are naturally occurring molecules with normal ("background") concentrations in the range of < 0.3 – 35 µg/m³. Concentration levels are dependent upon location and season (Doc IIIA JOINT: FA_BPR_Ann_II_10_3_2).

No effects on the ozone layer or relevant contribution to global warming and acidification are expected.

4.2.2 Sewage treatment plant (STP)

Two tests on the inhibitory effect of formic acid on microbial activity were submitted.

- The inhibition of oxygen consumption in activated sludge due to **formic acid** was evaluated in a test conducted according to ISO/DIS 8192 Part B, which is similar to OECD 209 (██████████ 1988c, BPD ID A7.4.1.4_01).

The highest concentration tested was 988 mg/L and the test concludes that the EC₂₀ is greater than this concentration.

However, BE eCA is of the opinion that this test cannot be used for the further risk assessment, since reliability cannot be assigned (value of 4) due to a severe lack in details in the original test report.

- A second study on the inhibitory effect of **formic acid** on the respiration rate of aerobic activated sludge, taken from a sewage treatment plant treating predominantly domestic sewage, was submitted by the applicant after the previous study was deemed lacking. The test (██████████ 2016, BPR ID A9.1.5_01) was performed over a contact period of 3 hours in a static test system, according to OECD 209 and following GLP.

Three replicates of each nominal test-concentration of 5, 15.8, 50, 158 and 500 mg/L were tested in parallel with six control replicates and four different concentrations of the reference item 3,5-dichlorophenol. Additionally, the same test-concentrations were repeated with the addition of N-allylthiourea to distinguish between total, heterotrophic and nitrification-related respiration.

The results of the statistical analysis of the respiration data collected, showed no considerable concentration-related inhibition of total, heterotrophic or nitrification-related respiration by formic acid. No EC_x values could therefore be determined at concentrations ≤500 mg/L following this test.

- The effects of **formic acid** on the growth of *Pseudomonas putida* was studied in a test performed according to DIN 38412 part 8 ([REDACTED] 1991, BPD ID A7.4.1.4_02).

Formic acid was tested in Penicillin flasks of 10 mL at nominal concentrations of 0, 7.81, 15.63, 31.25, 62.5, 125, 250, 500 and 1000 mg/L. Four parallel repeats per test concentration were run, including an un-inoculated sample. After an incubation time of 17 hours, the extinction was measured at 436 nm. No analytical monitoring to confirm the nominal test concentrations was performed. The pH was measured in the un-inoculated samples at test start and end, and in the inoculated samples at test end.

The lowest concentration revealing an inhibition is 31.25 mg/L, with an inhibition of 1.86 %. An inhibition of over 99% compared to the control is observed in the test concentrations of 62.5 mg/L and up. This inhibition can be partly due to the acidic pH, but is not confirmed with a test run at neutralised concentrations.

Statistical analysis calculates an EC₁₀ of 33.9 mg/L, an EC₅₀ of 46.7 mg/L and an EC₉₀ of 59.5 mg/L.

Summary table – inhibition of microbial activity										
Method, Guideline, GLP status, Reliability	Test material	Species/ Inoculum	Endpoint	Exposure		Results			Remarks	Reference
				Design	Duration	EC ₁₀ [mg/L]	EC ₅₀ [mg/L]	EC ₉₀ [mg/L]		
ISO/DIS 8192 Part B No GLP Reliability 4	FORMIC ACID	activated sludge	oxygen consumption	respiration inhibition	30 min	EC ₂₀ = >988	/	/	only single concentration no analytical verification abstract report	[REDACTED] 1988c BPD ID A7.4.1.4_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_5_1_a</i>
OECD 209 GLP Reliability 1	FORMIC ACID	activated sludge	oxygen consumption	respiration inhibition	3 h	>500	>500	>500	nominal concentrations, adjusted for pH with NaOH	[REDACTED] 2016 BPR ID 9.1.5_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_5_01_final_28Mar2017</i>

DIN 38412 part 8 no GLP reliability 2	FORMIC ACID	<i>Pseudomona s putida</i>	optical cell density at 436 nm	growth inhibition	17 h	33.9	46.7	59.5	no measured concentratio n No pH adjusted concentratio ns	1991 BPD ID A7.4.1.4_02 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_5</i> _2
--	----------------	--------------------------------	--------------------------------------	----------------------	------	------	------	------	---	--

Value used in Risk Assessment

Value/conclusion	Formic acid: EC ₁₀ > 500 mg/L
Justification for the value/conclusion	Based on available results, the short-term test is preferred, in accordance with the retention time in a STP. The EC ₁₀ value was determined at concentrations >500 mg/L. The 17h test is considered less relevant, since it uses glucose as a substrate.

4.2.3 Aquatic compartment

In aqueous solution and at neutral pH, formic acid and water-soluble formate salts dissociate and are present as the formate anion in solution. The behaviour of chemical dissociation in water has particularly been investigated with potassium diformate (CAS No. 20642-05-1), which served as test compound in several toxicity studies. Based on these physico-chemical properties, it is justified to include kinetic and metabolism studies conducted with water-soluble formate salts in these considerations. In order to provide data for the ecotoxicity of formic acid without effects due to the low pH which is induced by formic acid, study results for ammonium formate and potassium formate were considered. As fish and aquatic invertebrates are sensitive towards ammonium dissolved in water, the results derived from testing with ammonium formate should not be used alone. On the other hand, no effects are expected due to the potassium ion (K⁺) contained in potassium formate.

4.2.3.1 FRESHWATER COMPARTMENT

4.2.3.1.1 Acute toxicity (freshwater)

4.2.3.1.1.1 Fish

Three acute toxicity tests to fish in freshwater were submitted, one using the test substance formic acid and two other using formate salts, meaning that in water the fish are mainly exposed to the formate anion.

- The acute toxicity of formic acid to the golden orfe (*Leuciscus idus L.*, golden variety) was studied following the German Industrial Standard DIN 38412, Part 15 (██████████ 1989, BPD ID A7.4.1.1_01).

The test system was a static system without any analytical monitoring of the test substance concentration. The nominal test concentrations of 0, 10, 21.5, 46.4 and 100 mg/L were tested using 10 fish for each concentration. An additional concentration of 100 mg/L was tested where the pH was neutralised using NaOH, in order to assess the effect of the low pH on the toxicity. The test water was reconstituted water according to the aforementioned guideline.

The fish were checked for symptoms and mortality after 1, 4, 24, 48, 72 and 96 hours. At these times also other parameters, such as temperature, dissolved oxygen and pH were analysed.

No mortality was reported for the control group and at test concentrations ranging from 10 and 46.4 mg/L and in the pH adjusted test concentration of 100 mg/L. In the non-pH adjusted 100 mg/L test concentration, 100 % mortality was reached after 1 hour.

When analysing the measured pH throughout the study, it is noted that the pH in the 100 mg/L test concentration was 3.3, which was probably a factor for the high mortality, since no mortality was reported at the same test concentration with neutralised pH.

However, it should be mentioned that in the 46.4 mg/L test concentration, pH was initially also quite low (4.3), but quickly rose to a neutral 7.2 at the end of the test. This seems to be an indication that the test substance concentration was not maintained throughout the test and

since no analytical monitoring was performed, this can be seen as a major deviation. Therefore BE eCA is of the opinion that the results from this test are not reliable and cannot be used in the further risk assessment.

The applicant was asked to comment on BE's assessment to attribute a reliability of 3 to this test. In their reply they make reference to the acute toxicity test discussed in the next bullet point below. In this test, also performed under static conditions, analytical monitoring showed that the test item concentration remained within the allowed variation. According to the applicant, if it is the case for that static test, it will also be the case for this static test.

BE understands that this perhaps may be some sort of indication, also when considering that the substance is hydrolytically stable (cfr. BPD ID A7.1.1.1.1_01) and the ready biodegradation tests (cfr. BPD ID A7.1.1.2.1_01 and BPD ID A7.1.1.2.1_02) show little degradation in the first couple of days. However, this does not answer the question of the rising pH and without conclusive proof of stability of the test substance, uncertainty remains. Therefore, BE is not inclined to change their assessment of the reliability. Therefore this test remains at reliability 3.

- In a first test, using **ammonium formate** as test material, the acute toxicity effects on zebrafish (*Danio rerio*) were studied according to OECD 203 (██████████ 2005, BPD ID A7.4.1.1_02). In a 96 h static test design, 10 fish each were exposed to nominal test concentrations of 0, 45, 90, 180, 360 and 720 mg/L. Samples for chemical analysis via ion chromatography were taken at test start and test end from all test vessels. Mean recovery values were higher than 80 % of the nominal concentrations, therefore the effect concentrations are based on these nominal concentrations.

Mortality, behavioural abnormalities, temperature, pH and dissolved oxygen were checked for each concentration after 24, 48, 72 and 96 hours. The test conditions - temperature, pH and dissolved oxygen – remained within acceptable limits throughout the test. No mortality was reported in the control group or in the test concentrations up to 90 mg/L. After 96h the lowest test concentration where all fish had died was 180 mg/L. The 96h LC₅₀ was determined using the geometric mean of the LC₀ and LC₁₀₀ resulting in a value of 127.28 mg/L.

- A second test using a formate salt, this time **potassium formate**, was also conducted according to OECD 203 (██████████ 1992e, BPD ID A7.4.1.1_03) using rainbow trout (*Oncorhynchus mykiss*). The test design was a semi-static one, with daily renewal of the test medium to ensure that test concentrations were maintained. However, no analytical monitoring was performed to corroborate this.

Ten fish per nominal test concentration of 0, 1000, 1800, 3200, 5600 and 10000 mg/L was used. Mortality was checked after 3, 6, 24, 48, 72 and 96 hours. Temperature, pH and dissolved oxygen were measured at test start and end, and remained within the acceptable limits. No mortality was reported for the control group or in the test concentrations up to 1800 mg/L. After 96h, the lowest test concentration where all fish had died was 5600 mg/L. The 96h LC₅₀ was determined using the method of Thompson & Weil (1952, moving-average interpolation) and resulted in a value of 3500 mg/L.

4.2.3.1.1.2 Invertebrates (*daphnia magna*)

Three acute toxicity tests on the aquatic invertebrate *Daphnia magna* were submitted, one using formic acid as test substance, while the other two used formate salts, meaning that in water the test animals are mainly exposed to the formate anion.

- The acute toxicity of formic acid to *Daphnia magna* was studied in a test performed according to Directive 79/831/EEC, C.2 (██████████ 1988, BPD ID A7.4.1.2_01).

The test species were exposed during 48-hours in a static test system without any analytical monitoring of the test concentration. Nominal test concentrations of 0, 0.781, 1.56, 3.12, 6.25, 12.5, 25, 50 and 100 mg/L were tested using 20 animals per concentration. Purified water was used as test medium, in which sulphuric acid was used to reduce the buffering capacity of the carbonic acid system and deionized water was added to reduce the total hardness.

At the beginning and after 3, 6, 24 and 48 hours, the swimming inability of the *Daphnia* was checked. Oxygen and pH measurements were performed at test initiation and after 48 hours.

The immobility in the control group was within the validity criterion (< 10 % immobility). In the two highest concentrations, 50 and 100 mg/L, immobility already reached 100% after 6 and 3 hours respectively. At these concentrations, pH was below 5 at the start of the experiment. No pH adjusted concentrations were tested to distinguish between the effect due to low pH and toxicity. In the test concentration of 25 mg/L, 10 % immobility was reached after 48 hours. The 48h EC₅₀ was calculated using the moving average method, resulting in a value of 32.19 mg/L. However, it must be kept in mind that it is unclear if this concentration causes mortality due to toxicity of the test substance or due to a decrease in the pH of the test medium.

- The acute toxicity of ammonium formate to aquatic invertebrates (*Daphnia magna*) was studied in a GLP-study according to OECD 202 (██████████ 2005, BPD ID A7.4.1.2_02).

In this 48-hour, static test, the test organisms were exposed to nominal concentrations of 0, 45, 90, 180, 360 and 720 mg/L. The *Daphnia* were checked for immobility at test start and after 24 and 48 hours. Oxygen content, pH and test item concentrations were determined at the start and end of the test. These parameters were within the acceptable ranges, leading to the use of nominal values for determining the toxicity values.

After 48 hours, no immobilisation was observed in the control and lowest test concentrations up to 90 mg/L, while 100 % immobility was reached in the highest tested concentration of 720 mg/L. The 48h EC₅₀ was determined using the ToxRat software (v2.09) and yielded a value of 365 mg/L.

- A second study, testing the acute toxicity of the formate ion to *Daphnia magna* was done using potassium formate as a test substance. The test was conducted according to OECD 202, conform GLP (██████████ 1992, BPD ID A7.4.1.2_03).

In this 48-hour static test, *Daphnia* were exposed to nominal concentrations of 0, 10, 18, 32, 56, 100, 180, 320, 560 and 1000 mg/L. Oxygen content and pH were measured at test initiation and after 48 hours, and remained within the acceptable intervals. No analytical monitoring of the test concentration was done, on request by the test sponsor. The applicant was asked why this was requested, but no explanation could be given. The test sponsor probably assumed the test concentration could be maintained for the exposure duration of 48 hours.

After 0, 24 and 48 hours, the test species were checked for mobility. At test end, immobilisation in the control group was within the acceptable range. The highest test concentration where no immobilisation was observed after 48 hours was 56 mg/L. No 100 % immobilisation was reached in any of the tested concentrations. The 48h EC₅₀ was determined using the moving average method, resulting in a value of 540 mg/L

4.2.3.1.1.3 Algae

4.2.3.1.1.3.1 Green algae

Three growth inhibition studies on green algae were submitted, one using formic acid as test substance, while the others used a formate salt, meaning that in water the test animals are mainly exposed to the formate anion.

- The inhibitory effect of **formic acid** on cell multiplication of the unicellular green algae *Desmodesmus subspicatus* was studied in a test performed according to German Industrial Standard DIN 38412, part 9 (██████████ 1988, BPD ID A7.4.1.3_01).

Algal exposition was performed in test tubes of 10 mL with flat bottom. The initial cell density of *Desmodesmus subspicatus* was 10⁴ cells/mL, which is higher than what is recommended according to OECD 201. The algae were exposed to nominal concentrations of 0, 0.781, 1.56, 3.125, 6.25, 12.5, 25 and 50 mg/L. No analytical monitoring of the test concentrations were done, but pH was measured in the uninoculated test concentrations at test start and after 96h and in the inoculated concentrations after 96h. Fluorescence measurements were performed after 0, 24, 48, 72 and 96 hours.

An inhibitory effect on the algal growth rate of 3 % was seen starting at the test concentration of 12.5 mg/L and 100 % inhibition was reached at the 50 mg/L test concentration, the highest concentration tested. The inhibition observed at the higher test concentrations might also be due to the low pH (4.9 in inoculated sample after 96h) and since no neutralized concentrations were tested it is not possible to distinguish between effect due to the pH or due to toxicity. After statistical analysis of the results through ToxRatPro, it is concluded that the E_rC₅₀ is 30.21 mg/L, the E_bC₅₀ is 26.92 mg/L, the E_rC₁₀ is 24.52 mg/L, the E_bC₁₀ is 17.71 mg/L and the NOE_rC is 6.25 mg/L.

- The inhibitory effect of **ammonium formate** on the growth of the unicellular green algae *Pseudokirchneriella subcapitata* was studied a 72h test performed according to OECD 201 (██████████ 2005, BPD ID A7.4.1.3_02).

Algal exposures were performed in 250 mL flasks containing 100 mL test solutions at the nominal test concentrations of 0, 76.8, 192, 480, 1200 and 3000 mg/L. The initial cell density was 10⁴ cells/mL and cell number determinations were performed after 24, 48 and 72 hours. Test item concentrations and pH were determined at the start and end of the test. These parameters were within the acceptable ranges, leading to the use of nominal values for determining the toxicity values. The test results were statistically analysed using the software ToxRat. NOEC was determined by the Welch t-test.

An inhibitory effect of algal growth of 3.4 % was already seen in the lowest test concentration of 76.8 mg/L. At the highest concentration (3000 mg/L) growth inhibition reached 39.8 %. Inhibition of biomass integral showed a 12.6 % inhibition at 76.8 mg/L, while at 3000 mg/L

an inhibition of biomass of 85.4 % was reached. Statistical analysis revealed a 72h E_rC₅₀ of 1240 mg/L, a 72h NOE_rC of less than 76.8 mg/L and a 72h E_bC₅₀ of 320 mg/L.

- A limit test on the inhibitory effect of **potassium formate** on the growth of the unicellular green algae *Scenedesmus subspicatus*, now known under the name *Desmodesmus subspicatus*, was performed according to OECD 201 (██████████ BPD ID A7.4.1.3_03).

Only a single concentration was tested, namely 1000 mg/L, and compared with the untreated control to determine effect. Algal exposure was performed in 250 mL flasks containing 100 mL test solution. The initial cell density was 9.2 x 10⁴ cells/mL. Measurements of fluorescence were performed at 0, 24, 48 and 72 hours. The nominal test concentration was not analytically verified, on request by the test sponsor. The applicant was asked for comment, but could not elaborate on the reasoning. The pH was measured at test initiation and end; and remained within the acceptable range.

The test concentration of 1000 mg/L had an inhibitory effect of 10 % on the algal growth rate (24-48 h), but an increase of 19 % in biomass was reported compared to the control. Since only one test concentration was tested, no EC₅₀ can be determined and it can only be stated that it will be higher than the concentration that was tested.

4.2.3.1.1.3.2 Cyanobacteria or diatoms

According to the Guidance on the Biocidal Product Regulation, Volume IV Part A, on information requirements, tests on the effect on growth rate of cyanobacteria or diatoms are required for phytotoxic and/or antimicrobial substances and should preferably be studied in a fresh water species.

The applicant did not submit a test on a freshwater species, but submitted a justification for non-submission (cfr. Doc IIIA JOINT: FA_BPR_ANN_II_9_1_3_JNS_21Sep2016). Therein they argument that an additional study will not provide additional information to address the risk to algae. This justification for non-submission was deemed acceptable.

Summary table - acute aquatic toxicity										
Method, Guideline, GLP status, Reliability	Test material	Species	Endpoint	Exposure		Results			Remarks	Reference
				Design	Duration	L(E)C ₀ [mg/L]	L(E)C ₅₀ [mg/L]	L(E)C ₁₀₀ [mg/L]		
Fish										

DIN 38412, GLP-study, Reliability 3	FORMIC ACID	<i>Leuciscus idus</i>	mortality	static	96h	46 ⁿ	67.82 ^g	100 ⁿ	no measured concentrations, only nominal	██████████ 1989 BPD ID A7.4.1.1_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_1_1</i>
OECD 203 GLP-study, Reliability 1	Ammonium formate	<i>Danio rerio</i>	mortality	static	96h	90 ⁿ	127.28 ^g	180 ⁿ	mean measured concentrations at test start and test end were >80 % of the nominal concentrations	██████████ 2005 BPD ID A7.4.1.1_02 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_1_2</i>
OECD 203 GLP-study Reliability 2	Potassium formate	<i>Oncorhynchus mykiss</i>	mortality	semi-static	96h	1800 ⁿ	3500 ^t	5600 ⁿ	semi-static conditions with daily renewal	██████████ 1992e BPD ID A7.4.1.1_03 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_1_3</i>
Invertebrates										
79/831/EE C, C.2 no GLP Reliability 2	FORMIC ACID	<i>Daphnia magna</i>	immobility	static	48h	25 ⁿ	32.19 ^t	50 ⁿ	no measured concentration No pH adjusted concentrations	██████████ 1988 BPD ID A7.4.1.2_01 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_2_1_1</i>
OECD 202 GLP study Reliability 1	Ammonium formate	<i>Daphnia magna</i>	immobility	static	48h	90 ⁿ	365	720 ⁿ	mean measured concentrations at test start and	██████████ 2005 BPD ID A7.4.1.2_02

									test end were >80 % of the nominal concentrations	<i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_2_1_2</i>
OECD 202 GLP study Reliability 2	Potassium formate	<i>Daphnia magna</i>	immobility	static	48h	56 ⁿ	540 ^t	>1000 (no 100% reached at highest test concentration)	no measured concentrations at request of test sponsor	██████████ 1992 BPD ID A7.4.1.2_03 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_2_1_3</i>
Algae (growth inhibition)						NOEC/ErC₁₀	E_bC₅₀¹	ErC₅₀²		
DIN 38412, part 9 no GLP Reliability 2	FORMIC ACID	<i>Desmodesmus subspicatus</i>	growth inhibition	static	72h	NOEC = 6.25 ⁿ ErC ₁₀ = 24.52 EbC ₁₀ = 17.71	26.92 ⁿ	30.21 ⁿ	no measured concentrations No pH adjusted concentrations	██████████ 1988 BPD ID A7.4.1.3_01 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_3_1_1</i>
OECD 201 GLP study Reliability 1	Ammonium formate	<i>Pseudokirchneriella subcapitata</i>	growth inhibition	static	72h	<76.8 ⁿ	320 ⁿ	1240 ⁿ	mean measured concentrations at test start and test end were >80 % of the nominal concentrations	██████████ 2005 BPD ID A7.4.1.3_02 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_3_1_2</i>
OECD 201 GLP study	Potassium formate	<i>Desmodesmus subspicatus</i>	growth inhibition limit test	static	72h	≥1000 ⁿ	>1000 ⁿ	>1000 ⁿ	no measured concentrations at	██████████ 1992 BPD ID A7.4.1.3_03

Reliability 2									request of test sponsor	<i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_3 _1_3</i>
<p>ⁿ (based on) nominal concentrations</p> <p>^g geometric mean of LC₀ and LC₁₀₀</p> <p>^t using Thompson & Weil method (moving-average interpolation)</p> <p>^l using a linear model</p> <p>¹ calculated from the area under the growth curve</p> <p>² calculated from growth rate</p>										

Conclusion on acute toxicity (freshwater)

- FORMIC ACID:

An aquatic acute toxicity test on formic acid was submitted for each trophic level. However, the test submitted for fish was deemed unreliable (3). The growth inhibition test on algae and the test on *Daphnia magna* can be of some value, but it must be born in mind that the endpoints derived in these studies are nominal values and that no distinction can be made between the effect due to acidity and effect due to the intrinsic toxicity of formic acid.

- AMMONIUM FORMATE:

An aquatic acute toxicity test using ammonium formate as a test substance was submitted for each of the three required trophic levels. All three tests were assessed with a reliability of 1. The resulting L(E)C₅₀'s between the three trophic levels are each in the same order of magnitude, with the 96h LC₅₀ of 127.28 mg/L for fish being the smallest recorded value.

- POTASSIUM FORMATE:

Aquatic acute toxicity tests using potassium formate are available for each of the three required trophic levels and are all considered reliable (2).

Value used in Risk Assessment	
Value/conclusion	<ul style="list-style-type: none"> • 96h LC₅₀ fish = 3500 mg/L • 48h EC₅₀ daphnia = 540 mg/L

	<ul style="list-style-type: none"> 72h ErC₅₀ algae > 1000 mg/L 72h NOEC algae = 1000 mg/L <p>Even though the order of magnitude of the endpoints derived from the studies submitted on the three trophic levels is not so different between the species, <i>Daphnia</i> are considered the more sensitive species with the lowest reported EC₅₀ of 540 mg/L</p>
Justification for the value/conclusion	Studies conducted with formate salts are considered acceptable to assess the toxicity of formic acid without the effects due to the low pH. Since fish, aquatic invertebrates and algae are known to be sensitive towards ammonium dissolved in water, the results derived from testing with potassium formate are considered more relevant, since no effects are expected due to the potassium ion (K ⁺).

4.2.3.1.2 Chronic toxicity (freshwater)

No chronic toxicity tests were submitted for fish or other aquatic plants. Based on the results obtained in the acute toxicity tests, it was concluded by the applicant that *Daphnia* were the most sensitive of the three trophic levels, and they therefore submitted a chronic tests using *Daphnia magna*.

- The chronic effect of **formic acid** on the reproduction of *Daphnia magna* was tested in a study performed according to OECD 211 (██████████ 2007, BPD ID A7.4.3.4_03).

Nominal test concentrations of 0, 1.0, 3.2, 10, 32 and 100 mg/L were tested. Because the pH of the two highest test concentrations was below the suitable range, these test concentrations were neutralized using NaOH. The actual concentrations were verified and remained within the acceptable range, so that results are based on the nominal concentrations.

Final results on statistical evaluations of the parameters reproduction, length and weight indicate that no effects were observed up to the highest concentrations of 100 mg/L. The corresponding NOEC is > 100 mg/L.

Summary table - chronic aquatic toxicity								
Method, Guideline, GLP status, Reliability	Test material	Species	Endpoint/ type of test	Exposure		Results LOEC/NOEC/EC ₁₀ [mg/L]	Remarks	Reference
				Design	Duration			

Fish								
No test submitted								
Invertebrates								
OECD 211 GLP-study Reliability 1	FORMIC ACID	<i>Daphnia magna</i>	reproduction, length, weight	semi- static (renewal every 2- 3 days)	21d	>100	pH was neutralized in the concentrations indicating a too low pH mean measured concentrations at test start and test end were >80 % of the nominal concentrations	██████████ 2007 BPD ID A7.4.3.4_03 <i>Doc IIIA JOINT:</i> <i>FA_BPR_Ann_II_9_1_6_2_a</i>
Other aquatic plants								
No additional studies were submitted (cfr. acute toxicity tests on algae)								

Value used in Risk Assessment	
Value/conclusion	21d NOEC aquatic invertebrates \geq 100 mg/L
Justification for the value/conclusion	See test results (with reliability 1) above.

4.2.3.2 ***SEDIMENT COMPARTMENT***

4.2.3.2.1 **Acute toxicity (freshwater sediment)**

Data waiving	
Information requirement	None
Justification	

4.2.3.2.2 **Chronic toxicity (freshwater sediment)**

Data waiving	
Information requirement	None
Justification	

4.2.3.3 **MARINE COMPARTMENT**

4.2.3.3.1 **Acute toxicity (seawater)**

4.2.3.3.1.1 *Fish*

In the environment, formic acid will mostly be present in its formate form. To test the effect of the formate anion on marine fish species, one study was submitted.

- An acute toxicity test, was performed with synthetic seawater, using **potassium formate** as the test material and juvenile turbot (*Scophthalmus maximus*) as test species. The test was conducted according to Guidelines of the UK Ministry of Agriculture, Fisheries and Food (██████████ 1992d, BPD ID A7.4.1.1_04), using a semi static test design, with daily renewal of the test medium. However, no analytical monitoring was performed to confirm that the test concentrations were indeed maintained throughout the test.

Ten fish each were exposed to the nominal test concentrations of 0, 320, 560, 1000, 1800 and 3200 mg/L. Temperature, pH and dissolved oxygen were checked at test start and after 24, 48, 72 and 96 hours, and the results show that these parameters remained within the acceptable limits. Test concentrations were checked for mortality after 3, 6, 24, 48, 72 and 96 hours. No mortality was reported in the control group or in test concentrations up to 1000 mg/L. After 96 h, the lowest test concentration that had a 100 % mortality rate was 3200 mg/L. The 96h LC₅₀ was determined using a linear model and resulted in a value of 1720 mg/L.

4.2.3.3.1.2 *Invertebrates (other species)*

In addition to the acute toxicity tests on *Daphnia magna*, the applicant submitted two supplementary studies on the acute toxicity effect of the formate ion on two marine invertebrate species.

- The acute toxicity of **potassium formate** to brown shrimp (*Crangon crangon*) was studied in a 96-hour semi-static test following Guidelines of the Ministry of Agriculture, Fisheries and Food, UK (██████████, 1992c, BPD ID A7.4.1.2_04).

Twenty shrimp each were exposed to nominal test concentrations of 0, 1000, 1800, 3200, 5600 and 10000 mg/L. Synthetic seawater was used as a test medium. No chemical analysis was carried out. Temperature, oxygen content and pH were measured at test start and after 24, 48, 72 and 96 hours, and remained within the acceptable ranges.

The test species were checked for moulting and mortality after 3, 6, 24, 48, 72 and 96 hours. No mortality was reported in the control group and in the test concentrations up to 1000 mg/L. After only 3 hours, all test species had died in the highest concentration of 1000 mg/L. After 96h all shrimp in the test concentration of 1800 mg/L and up had died. The 96h LC₅₀ was calculated according to a quadratic model and yielded the value of 1308 mg/L.

- The acute toxicity of **potassium formate liquor** (i.e. potassium formate 75% in water) to the marine copepod *Acartia tonsa* was studied in a 48-hour static test according to a guideline proposal to ISO TC147/SC5/WG2 (██████████ 1994, BPD ID A7.4.1.2_05).

Twenty copepods each were exposed to nominal test concentrations of 0, 56, 100, 320, 560 and 1000 mg/L. Natural seawater was used as a test medium. No chemical analysis was carried out. Temperature, salinity, oxygen content and pH were measured at test start and end in the control group and in the group testing 1000 mg/L. Based on these measurements, these parameters remained within the acceptable range.

The test species was checked for mortality after 24 and 48 hours. Mortality in the control group was within the acceptable limits. After 48 hours, no or insignificant mortality occurred in test concentrations up to 320 mg/L. In the 560 mg/L concentration 20 % and in the 1000 mg/L concentration 65 % of the animals had died after 48 hours. The 48h LC₅₀ was graphically estimated as 531 mg/L.

4.2.3.3.1.3 Algae (diatoms)

The effect of the formate ion on the growth of marine diatoms was demonstrated by the submission of one test.

- The inhibitory effect of **potassium formate liquor** (i.e. potassium formate 75% in water) on cell multiplication of the marine diatom *Skeletonema costatum* was studied according to ISO/DIS 10253 (██████████ 1994, BPD ID A7.4.1.3_04).

Exponentially growing algae were exposed to nominal concentrations of 0, 56, 100, 320, 560, and 1000 mg/l; using 250 mL flasks containing 200 mL of the test medium. Natural seawater was used as in preparing the culture medium and the initial cell density of the *Skeletonema costatum* was 10⁴ cells/mL. No analytical monitoring of the test substance concentrations was performed throughout the test. The pH was measured at the start and end of test and the results show that this parameter remained within the acceptable range.

Cell density measurements were performed after 24, 48 and 72 hours. The EC₅₀ values were estimated using a logarithm linear or logarithm-probit plot of concentration and percent growth inhibition. At the highest tested concentration, 6 % inhibition of the growth rate and 20 % inhibition of the biomass integral was calculated after 72 hours. The 72-hour EC₅₀ could therefore only be estimated as being larger than 1000 mg/L.

Summary table - acute aquatic toxicity										
Method, Guideline, GLP status, Reliability	Test material	Species	Endpoint	Exposure		Results			Remarks	Reference
				Design	Duration	L(E)C ₀ [mg/L]	L(E)C ₅₀ [mg/L]	L(E)C ₁₀₀ [mg/L]		
Fish										
UK Ministry of Agriculture,	Potassium formate	Scophthalmus maximus	mortality	semi-static, marine	96h	1000 ⁿ	1720 ^l	3200 ⁿ	marine species	██████████ 1992d

Ficheries and Food guideline GLP study Reliability 2									semi-static conditions with daily renewal	BPD ID A7.4.1.1_04 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_1_4</i>
Invertebrates										
Guidelines of the Ministry of Agriculture, Fisheries and Food, UK GLP study Reliability 2	Potassium formate	<i>Crangon crangon</i>	mortality	semi-static marine	96h	1000 ⁿ	1308	1800	marine species no measured concentrations	██████████ 1992c BPD ID A7.4.1.2_04 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_2_2_1</i>
ISO TC147/SC5/WG2 GLP study Reliability 2	Potassium formate liquor	<i>Acartia tonsa</i>	mortality	static marine	48h	320 ⁿ	531	>1000 (no 100% reached at highest test concentration)	marine species no measured concentrations	██████████ 1994 BPD ID A7.4.1.2_05 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_2_2_2</i>
Algae (growth inhibition)						NOE_rC/E_rC₁₀	E_bC₅₀¹	E_rC₅₀²		
ISO/DIS 10253 (draft 1991) GLP study Reliability 2	Potassium formate liquor	<i>Skeletonema costatum</i>	growth inhibition	static marine	72h	Not reported	>1000 ⁿ	>1000 ⁿ	marine species no measured concentrations	██████████ 1994 BPD ID A7.4.1.3_04 <i>Doc IIIA JOINT: FA_BPR_Ann_II_9_1_3_2</i>
ⁿ (based on) nominal concentrations ^g geometric mean of LC ₀ and LC ₁₀₀ ^t using Thompson & Weil method (moving-average interpolation) ^l using a linear model ¹ calculated from the area under the growth curve										

² calculated from growth rate

Value used in Risk Assessment	
Value/conclusion	<ul style="list-style-type: none"> • 96h LC₅₀ fish = 1720 mg/L • 48h EC₅₀ invertebrates = 531 mg/L • 72h ErC₅₀ algae > 1000 mg/L <p>Just as with the studies in fresh water, the order of magnitude of the toxicity values derived from the studies submitted on the three trophic levels is not so different between the species, Daphnia are considered the more sensitive species with the lowest reported marine EC₅₀ of 531 mg/L</p>
Justification for the value/conclusion	Studies conducted with formate salts are considered acceptable to assess the toxicity of formic acid without the effects due to the low pH. No effect on the test species is expected due to the potassium ion (K ⁺).

4.2.3.3.2 Chronic toxicity (seawater)

Data waiving	
Information requirement	None
Justification	

4.2.3.4 SEA SEDIMENT COMPARTMENT

4.2.3.4.1 Acute toxicity (sea sediment)

Data waiving	
Information requirement	None

Justification	
----------------------	--

4.2.3.4.2 Chronic toxicity (sea sediment)

Data waiving	
Information requirement	None
Justification	

4.2.3.5 **HIGHER TIER STUDIES ON AQUATIC ORGANISMS**

None such studies for formic acid or the formate ion were submitted or required at this point.

4.2.4 Terrestrial compartment

Formic acid is soluble in water and has a low adsorption potential ($\log K_{oc} = 1.48$). In soil formic acid will be mobile and present in the pore and ground water. The compound is however readily biodegradable and no long-term exposure of soil organisms to formic acid in soil is expected.

No specific results of ecotoxicity tests on terrestrial organisms are available for the risk assessment.

Data waiving	
Information requirement	No specific information submitted, but not required.
Justification	Equilibrium partitioning method will be used in the risk assessment.

4.2.5 Groundwater

No data on groundwater was submitted.

4.2.6 Birds and mammals

No studies on birds were submitted.

The available literature data show a low intrinsic toxicity of formic acid or formate to birds (*Doc IIIA JOINT: FA_BPR_Ann_II_9_4_1*), with a reported LD₅₀ ≥ 111 mg/kg_{bw} for wild-trapped redwinged blackbirds and no adverse effects on body weight, feed utilisation or liveability up to 1.0%_{w/w} Formic Acid and 1.45 % calcium formate in the diets of male broilers.

For oral studies on mammals, please see paragraphs 3.6.1 and 3.7.1 above.

Summary table –toxicity to birds and mammals									
Method, Guideline, GLP status, Reliability	Species	Endpoint	Exposure		Results [mg a.i./kg bw or feed]			Remarks	Reference
			Design	Duration	LD/LC ₅₀	LOEL/ LOEC	NOEL/ NOEC		
Birds									
No test submitted									
Mammals									
OECD 408 GLP: yes Rel. 1	Rat (≥ 6 weeks)	sub-chronic repeated oral toxicity Systemic values	OECD 408	90 days	/	2100	840	study with potassium diformate as test substance	██████████ 1998 BPD ID A6.4.1_01 <i>Doc IIIA JOINT: FA_BPR_Ann_II _8_9_2_01</i>
Comparable to 94/40/EEC GLP: yes Rel. 1	Rat (≥ 6 weeks)	long-term repeated oral toxicity Systemic values	Compara ble to 94/40/E EC	104 weeks	/	1400	280		██████████ 2002a BPD ID A6.5_01 <i>Doc IIIA JOINT: FA_BPR_Ann_II _8_11_1_02</i>

Value used in Risk Assessment	
Value/conclusion	NOAEL _{bird} = no value available NOAEL _{mammal, oral_chr} = 280 mg/kg _{bw} .day
Justification for the value/conclusion	Data on the avian toxicity of formic acid is not required. Data on the toxicity of formic acid on mammals was submitted for the human health part (see §3.6.1 and 3.7.1 on sub-chronic and long-term toxicity)

4.2.7 Primary and secondary poisoning

4.2.7.1 *PRIMARY POISONING*

Data waiving	
Information requirement	None
Justification	/

4.2.7.2 *SECONDARY POISONING*

Data waiving	
Information requirement	No
Justification	Formic acid is not expected to bioaccumulate based on the experimentally derived log Kow of -2.1 (23 °C, pH7) and the calculated BCF (see §4.1.3 above). Therefore, secondary poisoning of formic acid in either the aquatic or terrestrial food chain is considered not relevant.

4.3 ENDOCRINE DISRUPTING PROPERTIES

No specific vertebrate tests to assess the endocrine disrupting (ED) properties of formic acid/formate for other non-target organisms were submitted by the applicant.

The 'Guidance for the identification of endocrine disruptors in the context of Regulations (EU) No 528/2012 and (EC) No 1107/2009' (ECHA/EFSA, 7 June 2018)⁸ states:

There may be cases in which due to the knowledge on the physico-chemical and (eco)toxicological properties of the substance an ED assessment does not appear scientifically necessary or testing for this purpose not technically possible (BP Regulation, Annex IV or PPP Regulation, Annex, Point 1.5).

The Annex IV, section 1.2 of the BPR states:

There may be sufficient weight of evidence from several independent sources of information leading to the assumption/conclusion that a substance has or does not have a particular dangerous property, while the information from each single source alone is considered insufficient to support this notion. [...] Where consideration of all the available data provides sufficient weight of evidence for the presence or absence of a particular dangerous property:

- *further testing on vertebrates for that property shall not be undertaken,*
- *further testing not involving vertebrates may be omitted.*

The following discussion focusses on a weight of evidence based argumentation to determine whether an ED assessment for formic acid and its salts, and the subsequent vertebrate testing appear scientifically necessary.

Formic acid is the simplest carboxylic acid. The formate anion is the common metabolite of formic acid and formate salts in aqueous solutions at physiological and environmental pH values. The water soluble formic acid and formate salts rapidly dissociate in aqueous solutions (fresh and salt water, body fluids) to formate and a cation (H⁺ or Na⁺, K⁺, NH₄⁺, etc.). Formic acid and formate are both readily biodegradable in freshwater, producing only water and CO₂. Formate is also biodegradable in seawater (Please refer to 4.1 Fate and distribution in the environment). Formic acid has no potential for bioaccumulation (indeed log Kow is - 2.1 at pH 7 and BCF calculated value is 3.2).

Formic acid is a natural compound occurring at significant concentrations in all environmental compartments. Formic acid has been identified as a major contributor to acidic rain in remote environments [Galloway *et al.* 1982; Chameides and Davis, 1983]. Known major sources of formic acid in the atmosphere include fossil fuel and biofuel combustion [Kawamura and Kaplan, 1985], biomass burning [Andreae and Merlet, 2001], plants [Gabriel *et al.* 1999] and photochemical oxidation of volatile organic precursors [Neeb *et al.* 1997]. Stavrou *et al.* (2012)

⁸ Referred to as 'Guidance on ED'.

showed that 90% of the formic acid produced is biogenic in origin, and largely sourced from tropical and boreal forests. The authors suggest that terpenoids – volatile organic compounds released by plants – are the predominant precursors.

In soil, low molecular-weight organic acids (including formic acid) are commonly present and are constantly released from root exudates and decayed plant litter, and through microbial organic matter decomposition. Takata *et al.* (2011) reported measured formic acid concentrations in soil ranging from 0.088 to 0.217 mg/kg_{dwt} on arable land and from 0.072 to 0.444 mg/kg_{dwt} in an adjacent oak forest. In a soil incubation experiment with poorly-drained soil conducted by Tete *et al.* (2015), formate concentrations in soil up to 0.18 mg/kg_{dwt} (at field capacity) and 1.89 mg/kg_{dwt} (in waterlogged soil) were observed. Van Hees *et al.* (2008) determined formate concentrations in the soil solution in different horizons of two coniferous forest soils. The authors observed the highest concentrations in the top soil (O1 horizon), ranging from 0.152 mg/L (3.3 µM, mean of 6 values) at Heden, Sweden to 0.354 mg/L (7.7 µM, mean of 6 values) at Nyänget, Sweden. Formic acid has a rapid turnover in soil. Half-lives in soil under aerobic conditions of ≤1 day were observed in studies conducted by Glanville *et al.* (2012) and Hellstén *et al.* (2005b).

Formic acid is also reported to be present in manure (up to 1415 mg kg dry matter in fresh dairy manure) [Baziramakenga and Simard, 1998; Spoelstra, 1979; Iannotti *et al.*, 1979] and surface water (up to 155 µg/L) [Murtaugh and Bunch, 1965; Hama and Handa, 1981].

Besides their presence in the environment, formic acid and its conjugate base, formate, are also naturally occurring in virtually all living organisms as essential endogenous metabolites critical for one-carbon metabolism [Lamarre *et al.* 2013]. Formate is formed from precursors in the intermediary metabolism and is used as an important constituent of the C1 intermediary metabolism which is required for the biosynthesis of amino acids and nucleic acid bases (purines and pyrimidines). As a critical endogenous metabolite, formate is not assumed to be inherently endocrine active.

Endocrine activity was investigated using *in silico* methods. None of the endocrine activity related profilers of the OECD QSAR Toolbox V4.1 showed an alert for formic acid. In fact, formic acid was grouped into the category “non-binder, non-cyclic structure”. Furthermore, binding to either oestrogen receptor (ER) or androgen receptor (AR) was estimated using *in silico* models implemented in OASIS TIMES (V2.27.19.13). None of the three models predicted a binding of formic acid to ER (with or without metabolism of parent compound) and AR (without metabolism). Please note that formic acid and formate have no structural similarity to intrinsic endocrine active substances (e.g. oestrogen, androgen). Altogether, based on *in silico* data it is very unlikely that formic acid exerts an endocrine/EATS-specific effect based on an endocrine mode of action.

In the mammalian dataset, no pattern related adverse effects in endocrine-sensitive organs or endpoints was identified in the available OECD Level 4 & 5 *in vivo* toxicity studies. Based on that mammalian dataset, it is concluded that formic acid does not meet the endocrine disruptor criteria for humans regarding E,A, S and T modalities (see **§Erreur ! Source du renvoi introuvable.**).

The Guidance on ED states that due to the high level of conservation of the endocrine system and receptor homology across the vertebrates, as well as the key enzymes involved, the mammalian data may also be relevant for other vertebrates.

Considering all above mentioned arguments, it was agreed by the Biocides Environment Working Group Meeting IV-2019 (ENV WG-IV-2019) that no further vertebrate testing is needed to conclude on the endocrine disruptor criteria for other non-target organisms. Based on the evaluation of available data in a weight-of-evidence based approach, it is concluded that formic acid does not meet the endocrine disruptor criteria for non-target organisms regarding E,A, S and T modalities.

Value used in Risk Assessment	
Value/conclusion	Formic acid does not meet the endocrine disruptor criteria for both human health and non-target organisms.
Justification for the value/conclusion	Conclusion agreed by the ENV WG-IV-2019 based on the evaluation of available data in a weight-of-evidence based approach.

4.4 DERIVATION OF PNECS

Compartment	PNEC	Remarks/Justification
Freshwater	$PNEC_{\text{freshwater}}: \geq 2 \text{ mg/L}$	<p><u>Organism:</u> <i>Daphnia magna</i></p> <p><u>Endpoint:</u> 21d NOEC $\geq 100 \text{ mg/L}$</p> <p><u>Assessment factor:</u> 50</p> <p><u>Extrapolation method:</u> assessment factor</p> <p><u>Justification:</u> The three taxonomic groups (fish, invertebrates, algae) are covered in short term data, of which <i>Daphnia</i> is considered as the most sensitive. A long-term NOEC for <i>Daphnia</i> is also available, and consequently the NOEC derived from the algal growth inhibition test is considered as an additional long-term study. An assessment factor of 50 is thus justified.</p>
Freshwater sediment	$PNEC_{\text{sediment}}: \geq 2.87 \text{ mg/kg}_{\text{wwt}}$ (converts to $\geq 13.2 \text{ mg/kg}_{\text{dwt}}$)	<p><u>Extrapolation method:</u> Equilibrium partitioning method</p> <p><u>Justification:</u> No specific data available or required</p> <p><u>Note:</u> Since also the PEC_{sediment} is calculated from the $PEC_{\text{freshwater}}$ using this method, the risk assessment and PEC/PNEC-ratio for the freshwater compartment are considered to cover the sediment compartment as well.</p>
Saltwater	$PNEC_{\text{seawater}}: > 0.2 \text{ mg/L}$	<p><u>Organism:</u> <i>Daphnia magna</i></p> <p><u>Endpoint:</u> 21d NOEC $> 100 \text{ mg/L}$</p> <p><u>Assessment factor:</u> 500</p> <p><u>Extrapolation method:</u> assessment factor</p> <p><u>Justification:</u> short term data for the basic three taxonomic groups (fish, invertebrates, algae) are available for both freshwater and saltwater species. No difference in sensitivity between the aquatic species in both media was observed. Long-term effect data</p>

Compartment	PNEC	Remarks/Justification
		(NOEC/EC ₁₀) are available for two trophic levels (algae and crustaceans) covering the most sensitive trophic level (= crustaceans). Therefore, the use of an assessment factor of 500 is justified.
Saltwater sediment	PNEC _{marine-sediment} : > 0.143 mg/kg _{wwt}	<u>Extrapolation method</u> : Equilibrium partitioning method <u>Justification</u> : No specific data available or required <u>Note</u> : Since also the PEC _{marine-sediment} is calculated from the PEC _{seawater} using this method, the risk assessment and PEC/PNEC-ratio for the marine compartment are considered to cover the sediment compartment as well.
Soil	PNEC _{soil} : ≥ 1.29 mg/kg _{wwt} (converts to ≥ 1.47 mg/kg _{dwt})	<u>Extrapolation method</u> : Equilibrium partitioning method <u>Justification</u> : No specific data available or required <u>Note</u> : The LOQ of the analytical method for soil established in the APCP section of this CAR is above the PNEC value for the soil compartment. Although not ideal, this is not a problem in the present case: the PNEC _{soil} is determined using the equilibrium partitioning method (and not based on measured test concentrations), and the risk assessment is based on calculated PEC values.
Groundwater	Not applicable	General drinking water limit: 0.0001 mg/L
Air	Not determined	Not relevant
STP	PNEC _{STP} : > 50 mg/L	<u>Organism</u> : activated sludge <u>Endpoint</u> : 3h EC ₁₀ > 500 mg/L <u>Assessment factor</u> : 10 <u>Extrapolation method</u> : assessment factor <u>Justification</u> : EC ₁₀ derived from OECD209

Compartment	PNEC	Remarks/Justification
Secondary poisoning birds	Not determined	No available data, but not considered relevant since no accumulation is expected
Secondary poisoning mammals	Not relevant	Risk assessment for secondary poisoning is not considered necessary, since no accumulation is expected.

5 ASSESSMENT OF EXCLUSION CRITERIA, SUBSTITUTION CRITERIA AND POP

5.1 EXCLUSION CRITERIA

5.1.1 Assessment of CMR properties

Criteria (BPR Article 5[1])	Assessment
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, carcinogen category 1A or 1B	Formic acid is not classified and does not meet the criteria to be classified as Carc. Cat. 1A or 1B.
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, mutagen category 1A or 1B	Formic acid is not classified and does not meet the criteria to be classified as Muta. Cat. 1A or 1B.
Active substances which have been classified in accordance with Regulation (EC) No 1272/2008 as, or which meet the criteria to be classified as, toxic for reproduction category 1A or 1B	Formic acid is not classified and does not meet the criteria to be classified as Repr. Cat. 1A or 1B.

Conclusion on CMR properties	The exclusion criteria in BPR Article 5(1)a-c are not met.
-------------------------------------	--

5.1.2 Assessment of endocrine disrupting properties

Criteria (BPR Article 5)	Assessment
Active substances which, on the basis of the criteria specified pursuant to the first subparagraph of paragraph 3 are considered as having endocrine-disrupting properties that may cause adverse effects in humans and to the environment.	The endocrine disrupting properties are assessed in accordance with the scientific criteria set out in COMMISSION DELEGATED REGULATION (EU) 2017/2100. Formic acid is not considered as having endocrine-disrupting properties that may cause adverse effects in humans and to the environment.
Conclusion on ED properties	The exclusion criteria in BPR Article 5(1)d are not met.

5.1.3 PBT Assessment (following Annex XIII to Regulation (EC) No 1907/2006)

5.1.3.1 ASSESSMENT OF PERSISTENCE

5.1.3.1.1 Screening

The available data on degradation reveal that formic acid should be considered readily biodegradable.

5.1.3.2 ASSESSMENT

P Criteria	Assessment
T1/2 > 60 days in seawater, or	no experimental data
T1/2 > 40 days in fresh- or estuarine water, or	no experimental data
T1/2 > 180 days in seawater sediment, or	no experimental data
T1/2 > 120 days in freshwater- or estuarine sediment, or	no experimental data
T1/2 ≤ 120 days in soil.	no experimental data

vP Criteria	Assessment
T1/2 > 60 days in sea-, fresh- or estuarine water water, or	no experimental data
T1/2 > 180 days in seawater-, freshwater- or estuarine sediment, or	no experimental data
T1/2 > 180 days in soil.	no experimental data

Conclusion on P / vP properties	Based on degradation data, formic acid is considered readily biodegradable. Therefore formic acid is considered not P or vP
--	---

5.1.3.3 **ASSESSMENT OF BIOACCUMULATION**

5.1.3.3.1 **Screening**

The log octanol-water partitioning coefficient (Log K_{ow}) for formic acid was determined at -2.10 (23 °C, pH7). Formic acid is considered hydrophilic in nature.

5.1.3.3.2 **Assessment**

B Criteria	Assessment
BCF > 2000	no experimental data

vB Criteria	Assessment
BCF > 5000	no experimental data

Conclusion on B / vB properties	The log K_{ow} for formic acid is well below the screening criterion of 4.5 for bioaccumulation. Therefore formic acid is not considered B or vB.
--	---

5.1.3.4 **ASSESSMENT OF TOXICITY**

5.1.3.4.1 **Screening**

The lowest available short term toxicity value for formic acid is the 48h EC_{50} for daphnia equal to 540 mg/L, which is well above the screening threshold for short-term aquatic toxicity of 0.01 mg/L.

The lowest chronic endpoint is a 21d NOEC for daphnia of equal or greater than 100 mg/L.

5.1.3.4.2 Assessment

T Criteria	Assessment
NOEC/EC10 (long-term) < 0.01 mg/L for freshwater or seawater organisms, or	The lowest chronic endpoint is a 21d NOEC for daphnia of 100 mg/L, which is well above the criterium.
substance meets the criteria for classification as carcinogenic (category 1A or 1B), germ cell mutagenic (category 1A or 1B), or toxic for reproduction (category 1A, 1B or 2) according to the CLP Regulation, or	Formic acid does not meet the criteria for classification as carcinogenic (category 1A or 1B), germ cell mutagenic (category 1A or 1B), or toxic for reproduction (category 1A, 1B or 2) according to the CLP Regulation.
there is other evidence of chronic toxicity, as identified by the substance meeting the criteria for classification: specific target organ toxicity after repeated exposure (STOT RE category 1 or 2) according to the CLP Regulation.	For formic acid there is no other evidence of chronic toxicity, as the substance does not meet the criteria for classification: specific target organ toxicity after repeated exposure (STOT RE category 1 or 2) according to the CLP Regulation.

Conclusion on T properties	Based on the available data, formic acid is considered not T
-----------------------------------	--

5.1.3.5 SUMMARY AND OVERALL CONCLUSIONS ON PBT OR VPvB PROPERTIES

5.1.3.5.1 Summary

- Formic acid is readily biodegradable
- Formic acid is hydrophilic and has no potential to bio-accumulate
- Formic acid is not classified for toxicity

5.1.3.5.2 Overall conclusion:

Based on the assessment described in the subsections above the submission substance is not a PBT / vPvB substance.

5.2 SUBSTITUTION CRITERIA

[Include an assessment if the active substance meets any of the following conditions:]

Substitution criteria (BPR, Article 10)	Assessment
One of the exclusion criteria listed in Article 5(1) is met but AS may be approved in accordance with Article 5(2)	For formic acid, the exclusion criteria in BPR Article 5(1)a-c are not met.
The criteria to be classified, in accordance with Regulation (EC) No 1272/2008, as a respiratory sensitiser is met	For formic acid, the criteria to be classified, in accordance with Regulation (EC) No 1272/2008, as a respiratory sensitiser are not met.
The acceptable daily intake, acute reference dose or acceptable operator exposure level, as appropriate, is significantly lower than those of the majority of approved active substances for the same product-type and use scenario	For formic acid, acceptable daily intake, acute reference dose or acceptable operator exposure level, as appropriate, are not significantly lower than those of the majority of approved active substances for the same product-type and use scenario
Two of the criteria for being PBT in accordance with Annex XIII to Regulation (EC) No 1907/2006 are met	No
There are reasons for concern linked to the nature of the critical effects which, in combination with the use patterns, amount to use that could still cause concern, such as high potential of risk to groundwater, even with very restrictive risk management measures	No
The AS contains a significant proportion of non-active isomers or impurities.	No

Conclusion on substitution criteria	The substitution criteria in BPR Article 10(1)a-f are not met.
--	--

5.3 ASSESSMENT OF LONG-RANGE ENVIRONMENTAL TRANSPORTATION AND IMPACT ON ENVIRONMENTAL COMPARTMENTS

Criteria	Assessment
The active substance or a degradation product is a persistent organic pollutant (POP) listed in Annex I of EC 850/2004	No
Assessment of long-range transport potential (LRTAP): <ul style="list-style-type: none">• Vapour pressure <1000 Pa and• half-life in air > 2 days or• Monitoring data in remote area showing that the substance is found in remote regions or• Result of modelling	No
The active substance or a degradation product is vP/vB or T?	No

Conclusion on LRTAP/POP assessment

Formic acid does not meet the criteria for being a POP or LRTAP.

PART B : EXPOSURE ASSESSMENT AND EFFECTS OF THE ACTIVE SUBSTANCE IN THE BIOCIDAL PRODUCT(S)

6 GENERAL PRODUCT INFORMATION

6.1 IDENTIFICATION OF THE PRODUCT

Name(s) of the product	
Trade name(s) or proposed Trade name(s)	Protectol® FM 85
Manufacturer's development code and number of the product	Not applicable
Formulation type	Water based concentrate / water soluble concentrate (SL)

6.2 COMPLETE QUALITATIVE AND QUANTITATIVE COMPOSITION OF THE BIOCIDAL PRODUCT

Active substance(s)					
ISO or Trivial name	IUPAC name or other accepted chemical name	EC number	CAS number	Composition / all constituents (upper and lower concentration limit in % (w/w))	Concentration in the product in % (w/w)
Formic Acid	Methanoic Acid	200-579-1	64-18-6	Minimum 99% w/w purity (BASF)	85% w/w (pure)

Other components / ingredients of the product					
ISO or Trivial name	IUPAC name or other accepted chemical name	EC number	CAS number	Concentration in the product in % (w/w)	Function
Please refer to BASF PT3 Confidential Annex					

6.3 PHYSICAL, CHEMICAL AND TECHNICAL PROPERTIES

Table 6.3.1: Protectol® FM 85

Property	Result	Test method applied or description in case of deviation	Remarks / Discussion / Justification for waiving	References
Physical state at 20°C and 101.3 kPa (85%)	Liquid	Organoleptic	The biocidal product contains 85 % active substance with no other ingredients than water. These properties are expected to be similar as for the active substance	Study no. 07L00084 ██████████ (2007)
Colour at 20°C and 101.3 kPa (85%)	Colourless	Organoleptic		
Odour at 20°C and 101.3 kPa (85%)	Pungent	Organoleptic		
Acidity / alkalinity (85%)	pH _{85% formic acid} = -1.6 At 1%: pH = 2.2	German Industrial Standard DIN 19268	Potentiometric measurement	Study no. 07L00172, ██████████ (2007)
	90.9530 ± 0.0663 % acidity	CIPAC MT 191	On 85% formic acid in water sample. Since test item is an acid, only acidity was tested.	Study no 16011907G975 ██████████ (2016a)
	pH = 2.18	CIPAC MT 75	At 24.8 °C On 1% aqueous solution of 85% formic acid sample	Study no 16011907G907 ██████████ (2016c)
Relative density (85%)	D ₄ ²⁰ = 1.19522	OECD 109	/	Study no. 02L00109, ██████████ (2002)

Storage stability, stability and shelf-life

Accelerated storage	Waived	-	Protectol® FM 85 is to be regarded as [REDACTED] Since a long term storage test at ambient temperature is available, the BE CA accepted the waiver to not submit an accelerated storage study at this stage, however, for product authorization the applicant will have to provide such study (performed at 40°C).	-
Long term storage at ambient temperature (85%)	Shelf life of 20 months	Storage conditions: transparent glass bottle 1000 ml; illumination: day light; temperature: approx. 24 °C; pressure: 1013 hPa	Acceptable: variation of 0.1 % (85.31 versus 85.24 %) after 20 months long term storage for formic acid based product need to be demonstrated at product authorisation in the commercial packaging.	[REDACTED] (2007b)
Low temperature stability (liquids)	Waived	EC method A.1	Protectol® FM 85 is a liquid at 0 °C and starts to show crystallization not before -10 °C, therefore it is not expected that the storage of the biocidal product at 0 °C will change the stability of the product.	Study no. 02L00109, [REDACTED] (2002) Statement on above study by [REDACTED] on 13/09/2016 (BPR ID 3.4.1.3_01)
Effects on content of the active substance				
Light	No effect	Long term storage at ambient temperature	Sample was stored in transparent glass bottle and subjected to daylight	[REDACTED] (2007b)
Temperature and humidity	Waived	-	Boiling point = 107.3 °C Formic Acid 99% shows no signs of decomposition up to the boiling point	Study no. 02L00109, [REDACTED] (2002) Study no. 07L00084, [REDACTED] (2007)

			Therefore the product can be considered stable at high temp. Humidity is irrelevant since the product is an aqueous solution.	
Reactivity towards container material	<p>Compatible:</p> <ul style="list-style-type: none"> - stainless steel, types 1.4306, 1.4307, 1.4311, 1.4404, 1.4541, 1.4571 - plastics: different types of PE like HD-PE; PP (for plugs and caps) <p>Not compatible:</p> <ul style="list-style-type: none"> - carbon steel, paper, board 	Based on experience with more concentrated solution of formic acid (99.4 %)	<p>Formic acid and solutions of formic acid are acidic. Therefore, materials which are not sufficiently resistant towards acids should not be used to avoid equipment damage and spoilage of products</p> <p>Materials used at BASF for container material (container, bung, gaskets, sealing, venting devices):</p> <ul style="list-style-type: none"> - polyethylene (Lupolen, Hostalen, Lucalen) - copolymer of ethylene and butylacrylate (Lucofin) - polypropylene (Moplen) - ethylene propylene diene monomer rubber (EPDM) - ethylene tetrafluoroethene (ETFE) <p>Plastic parts in contact with product must only be made from virgin material (= without addition of regrind, recycle and production waste) in order to avoid contamination with heavy metals.</p> <p>Applicant should provide suitable data at product authorisation stage.</p>	██████████ (2007a)
Technical characteristics				

Wettability	Waived	-	Not applicable	-
Suspensibility, spontaneity and dispersion stability	Waived	-	Not applicable	-
Wet sieve analysis and dry sieve test	Waived	-	Not applicable	-
Emulsifiability, reemulsifiability and emulsion stability	Waived	-	Not applicable, Protectol® FM 85 is not an emulsion	-
Disintergration time	Waived	-	Not applicable	-
Particle size distribution, content of dust / fines, attrition, friability	Waived	-	Not applicable	-
Persistent foaming	Protectol® FM 85 is a non-foaming liquid solution	Experience in use	Information on persistent foaming would be necessary at product authorisation level if additional formulants are introduced in the composition.	██████████ (2007c)
Flowability, pourability, dustability	Waived	-	Not applicable	-
Burning rate – smoke generators	Waived	-	Not applicable	-
Burning completeness – smoke generators	Waived	-	Not applicable	-

Composition of smoke – smoke generators	Waived	-	Not applicable	-
Spraying pattern - aerosols	Waived	-	Not applicable	-
Other technical characteristics	Waived	-	Not applicable	-
Physical and chemical compatibility with other products including other biocidal products with which its uses is to be authorised				
Physical compatibility	Waived	-	Not applicable, Protectol® FM 85 is not intended to be used in combination with other products	-
Chemical compatibility	Waived	-		-
Degree of dissolution and dilution stability	Waived	-	As the active substance is highly soluble in water, no issue with stability in water is expected.	-
Surface tension	At 20 °C: 71.5 mN/m	OECD 115	Result for solution with 99.4 % formic acid. The [REDACTED] other ingredients of Protectol® FM 85 is [REDACTED] As formic acid and water (at 20 °C: 72.75 mN/m) have almost identical surface tensions, no significant change of this value is expected for dilutions of formic acid	Study no. 07L00084, [REDACTED] (2007)
Viscosity	Dynamic viscosity At 20 °C: 1.80 mPa.s At 40 °C: 1.22 mPa.s Kinematic viscosity	OECD 114	For more concentrated (99.4 %) formic acid	Study no. 07L00084, [REDACTED] (2007)

	At 20 °C: 1.47 mm ² /s At 40 °C: 1.02 mm ² /s			
	Dynamic viscosity At 20 °C: 1.61 mPa.s At 40 °C: 1.10 mPa.s Kinematic viscosity At 20 °C: 1.37 mm ² /s At 40 °C: 0.95 mm ² /s	Ubbelohde viscometer (glas), similar to DIN 51562	For more diluted (75 %) formic acid	Study no. 2014-209.1 ██████████ (2014)
	Dynamic viscosity At 20 °C: 1.71 mPa.s At 40 °C: 1.18 mPa.s Kinematic viscosity At 20 °C: 1.42 mm ² /s At 40 °C: 0.99 mm ² /s	Expert judgement	Estimation for product Protectol® FM 85 with 85 % formic acid	/
Physical hazards and characteristics				
Explosives (85%)	The substance is not explosive	UN Manual of Tests and Criteria (2010)	The substance has no chemical groups indicating explosive properties	██████████ (2006)
Flammable gases	Waived	-	Not applicable	-
Flammable aerosols	Waived	-	Not applicable	-
Oxidising gases	Waived	-	Not applicable	-
Gases under pressure	Waived	-	Not applicable	-

Flammable liquids	Not a flammable liquid Flash point = 73.5 °C Classified as Flammable Liquid 3 (H226)	German Industrial Standard DIN EN ISO 2719, method A (Pensky-Martens closed cup)	For solution with 83 % formic acid	Study no. SIK-No.14/1849, [REDACTED] (2015)
Flammable solids	Waived	-	Not applicable	-
Self-reactive substances and mixtures (85%)	The substance is not self-reactive	UN Manual of Tests and Criteria (2010)	The substance has no chemical groups indicating explosive or self-reactive properties	
Pyrophoric liquids	Waived	-	Not a pyrophoric liquid, based on auto-ignition temperature (528 °C for 99.4 % formic acid) and experience in manufacture and handling	Study no. SIK-Nr.07/1018, [REDACTED] (2007)
Pyrophoric solids	Waived	-	Not applicable	-
Substances and mixtures which in contact with water emit flammable gases	Waived	-	Not applicable	-
Oxidising liquids (85%)	The substance is not an oxidising liquid	UN Manual of Tests and Criteria (2010)	The compound contains oxygen but this element is chemically bonded only to carbon and hydrogen The compound does not contain any halogen atoms	[REDACTED] (2006)
Oxidising solids	Waived	-	Not applicable	-
Organic peroxides	Waived	-	Not applicable	-

Corrosive to metals	Corrosive to steel Not corrosive to aluminium Classified as Corrosive to Metal (H290)	UN Test C.1 (37.4)	On 85% formic acid in water sample	Study no 16011907G979 ██████████ (2016b)
	Compatible materials: - stainless steel, types 1.4306, 1.4307, 1.4311, 1.4404, 1.4541, 1.4571 Not compatible: - carbon steel Classified as Corrosive to Metal (H290)	Based on experience	On 99% formic acid	██████████ (2007a)
Auto-ignition temperature of products (liquid and gas)	Auto-ignition temperature: 528 °C (corrected according to EN 14522)	EC method A.15	Result for solution with 99.4 % formic acid. The ██████████ other ingredients of Protectol® FM 85 ██████████	Study no. SIK-Nr.07/1018, ██████████ (2007)
Relative self-igniton temperature of solids	Waived	-	Not applicable	-
Dust explosion hazard	Waived	-	Not applicable	-

6.4 HAZARD IDENTIFICATION FOR PHYSICAL AND CHEMICAL PROPERTIES

The product **Protectol® FM 85** as manufactured is a colourless liquid with a pungent smell. The relative density of the product is 1.195 at 20 °C. The product has a long term stability of 20 months and is stable under cold storage conditions. Light influence is negligible. The surface tension is expected to be around 72 nN/m and the viscosity around 1.71 mPa.s. Physical and chemical compatibility with other products are not relevant.

6.5 ANALYTICAL METHODS FOR DETECTION AND IDENTIFICATION

Please note that only Formic acid and the formate ion are analysed in the monitoring table presented below

Analytical methods for the analysis of the product as such including the active substance, impurities and residues									
Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ) or other limits	Reference
					Range	Mean	RSD		
Active substance	The test principle for the determination of formic acid is titration of the organic acid with sodium hydroxide using an automated commercial titration system "Titrol alpha plus" from SI Analytics. The test item used for the validation was 100% formic acid (from VWR International GmbH, Darmstadt, Germany; certificate contained in the report), both as	5	$r > 0.99$		-	-	-	-	██████████ (2017)

	<p>pure test item and diluted to 85% with water.</p> <p>In addition, GC-MS analysis was performed to confirm the identity of formic acid and to demonstrate the absence of any other interfering organic acid or other impurity.</p> <p><i>The study was conducted in accordance with SANCO/3030/99 rev. 4 and under GLP conditions. The procedure is sufficiently described in the sections above.</i></p>								
--	---	--	--	--	--	--	--	--	--

Analytical methods for monitoring

			Linearity	Specificity	Recovery rate (%)		Reference
--	--	--	-----------	-------------	-------------------	--	-----------

Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements			Range	Mean	RSD	Limit of quantification (LOQ) or other limits	
Depending on extra data request for ASAS	UV absorption (334, 340 or 365 nm)	7	r2= 0.99981	none	0.2 to 5 mg/L			0.2 mg/L	██████████ (2013)

Analytical methods for soil

Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ) or other limits	Reference
					Range	Mean	RSD		
a.s. (formic acid)	UV absorption after stoichiometric , enzyme-catalyzed reduction of NAD+ to NADH by formic acid Formic acid (formate) is quantitatively oxidized to bicarbonate by nicotinamide adenine dinucleotide (NAD) in the	5- 50 mg/kg (25 number of measurements)	r2= 0.99981 <i>Linearity is given in the range 0.2 mg formic acid /l sample solution to 200 mg formic acid/l sample</i>	The method is specific for formic acid. Acetic acid, propionic acid, oxalic acid and L-ascorbic acid do not influence the determination. Formaldehyde reduces the reaction rate but does not influence the specificity of the method.”	Fortification range 5-50 mg/kg	RR 5 mg/L: 31% RR 10 mg/kg: 85% RR 50 mg/kg: 93%	4.7 (at 50 mg/kg)	10 mg/kg	██████████ (2013)

	<p>presence of formate dehydrogenase (FDH).</p> <p>FDH Formate + NAD⁺ + H₂O → bicarbonate + NADH + H⁺</p> <p>The amount of NADH formed is stoichiometric to the amount of formic acid. The increase in NADH is measured by means of its light absorbance at 334, 340 or 365 nm. The molar extinction coefficient is large at 340 nm [$\epsilon = 6.3 \text{ L}/(\text{mmol} \times \text{cm})$], i.e. the method is most sensitive at this</p>							
--	--	--	--	--	--	--	--	--

	<p>wavelength. The extinction coefficient allows to calculate the formate concentration from the absorbance difference at the start and at the end of the reaction, which is a common method in biochemical laboratories.</p> <p><i>Photometric measurements provide the basis for the majority of quantitative methods in biochemistry and are related to the amount of light absorbed. The temperature range should be 20-25°C, the pH value at approx. 7.5.</i></p>								
--	--	--	--	--	--	--	--	--	--

	<i>The specificity of the method is based on the specificity of the enzyme for its substrate (known as "key-lock principle").</i>								
--	---	--	--	--	--	--	--	--	--

Analytical methods for air									
Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ) or other limits	Reference
					Range	Mean	RSD		
Depending on extra data request for <i>a.s.formic acid</i>	Ion Chromatography <i>Material and conditions: Ion chromatographer DIONEX DX 120 with conductivity detector and autosampler. Pre-column: Micro-Guard Cation H-Cartridge (Bio-</i>	6 (per concentration)	<i>Formic acid, 1.2 to 47.8 mg/L.</i>	Specificity depends on the column and eluant chosen, and also on the separation condition.	94%-95%	95% RR for 0.9mg/m3	9.7% for 0.9 mg/m3 fortification level	<i>Absolute: 0.1µg; relative: 0.12 mg/m3 formic acid for a 140 l air sample, 10 ml absorption volume and 50 µl injection volume</i>	██████████ 2007 (BPD ID A4.1_02)

	<i>Rad, Munich). Column: Aminex HPx-87H (Bio- Rad). Suppressor: AMMS-ICE II P/N 037107 (Dionex). Suppressor solution: Tetrabutyl ammonium hydroxide, 5 mM. Eluent: hydrochloric acid 0.15 mM. Flow rate 0.6 mL/min. Flow rate suppressor: 1 mL/min Injection volume: 50 µL. Temperature: room temperature.</i>								
--	--	--	--	--	--	--	--	--	--

Analytical methods for water									
Analyte (type of analyte e.g. active substanc e)	Analytical method	Fortificatio n range / Number of measureme nts	Linearity	Specific ity	Recovery rate (%)			Limit of quantificat ion (LOQ) or other limits	Referen ce
					Ran ge	Mean	RSD		

Active substance formic acid	UV absorption after stoichiometric, enzyme-catalyzed reduction of NAD ⁺ to NADH by formic acid. Formic acid (formate) is quantitatively oxidized to bicarbonate by nicotinamide adenine dinucleotide (NAD) in the presence of formate dehydrogenase (FDH). FDH Formate + NAD ⁺ + H ₂ O → bicarbonate + NADH + H ⁺	Drinking water: 20 (5 measurements at each of the four fortification levels) and blanks Surface water: 15 (5 measurements at each of the three fortification levels) and blanks	<i>given in the range 0.2 to 5 mg/L. R²= 0.9997 for the regression curve for all measurements given in the range 0.2 to 5 mg/L. R²= 0.99998 for the regression curve for all measurements</i>	(enzyme specific for formic acid)	0.2 to 5 mg/L	Fortification level [mg/L]	Recovery [%] Drinking water	Recovery [%] Surface water	Fortification level [mg/L]	Rel SD [%] Drinking water	Rel SD [%] Surface water	0.2 mg/L in drinking water and surface water	██████████ (2013)
						0.2	103	100					
						0.5	91	n.d.	0.5	2.4	n.d.		
						2	103	81	2	6.6	1.6		
						5	101	78	5	3.7	1.7		

<p>The amount of NADH formed is stoichiometric to the amount of formic acid. The increase in NADH is measured by means of its light absorbance at 334, 340 or 365 nm. The molar extinction coefficient is large at 340 nm [$\epsilon = 6.3 \text{ L}/(\text{mmol} \times \text{cm})$], i.e. the method is most sensitive at this wavelength. The extinction coefficient allows to calculate the formate concentrati</p>								
--	--	--	--	--	--	--	--	--

<p>on from the absorbance difference at the start and at the end of the reaction, which is a common method in biochemical laboratories .</p> <p><i>Photometric measurements provide the basis for the majority of quantitative methods in biochemistry and are related to the amount of light absorbed. The temperature range should be 20-25°C, the pH value at approx. 7.5. The</i></p>								
---	--	--	--	--	--	--	--	--

	specificity of the method is based on the specificity of the enzyme for its substrate (known as "key-lock principle").								
--	--	--	--	--	--	--	--	--	--

Analytical methods for animal and human body fluids and tissues									
Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ) or other limits	Reference
					Range	Mean	RSD		
Active substance formic acid	UV absorption <i>Formic acid (formate) is quantitatively oxidized to bicarbonate by nicotinamideadenine dinucleotide (NAD) in the presence of formate</i>	n.a.	Linearity is given in the range 0.2 mg formic acid/l sample solution to 200 mg formic acid/l	yes	0.2 mg/L to 200 mg/L	100%	0.48-2.40%	0.2 mg/L	Anonymous (2007) UV test for the determination of Formic Acid in foodstuffs and other materials, Roche commercial test

<p><i>dehydrogenase (FDH).</i></p> <p><i>FDH</i> $\text{Formate} + \text{NAD}^+ + \text{H}_2\text{O} \longrightarrow \text{bicarbonate} + \text{NADH} + \text{H}^+$</p> <p><i>The amount of NADH formed is stoichiometric to the amount of formic acid. The increase in NADH is measured by means of its light absorbance at 334, 340 or 365 nm. NADH and NADPH absorb in the long-wave UV-range with a maximum at 340 nm, whilst the oxidized forms (NAD and NADP) do not show any absorption at this wavelength (see Figure 3). Therefore, any reaction in which either NAD(P) is reduced or NAD(P)H is oxidized may be measured by recording the</i></p>		<p>sample solution (cf. full test description in Section A4.1_01).</p>						<p>combination, R-Biopharm, Cat. No. 10 979732 035</p>
---	--	--	--	--	--	--	--	--

	<i>change in absorption in this wave length range.</i>								
--	--	--	--	--	--	--	--	--	--

Analytical methods for monitoring of active substances and residues in food and feeding stuff									
Analyte (type of analyte e.g. active substance)	Analytical method	Fortification range / Number of measurements	Linearity	Specificity	Recovery rate (%)			Limit of quantification (LOQ) or other limits	Reference
					Range	Mean	RSD		
Active substance formic acid	UV absorption Formic acid (formate) is quantitatively oxidized to bicarbonate by nicotinamideadenine dinucleotide (NAD) in the presence of formate dehydrogenase (FDH). FDH $\text{Formate} + \text{NAD}^+ + \text{H}_2\text{O} \longrightarrow \text{bicarbonate} + \text{NADH} + \text{H}^+$ The amount of	16	Linearity is given in the range 0.2 mg formic acid/l sample solution to 200 mg formic acid/l sample	Specific to formic acid	0 to 50 mg/L	recovery 92% at fortification level 10 mg/L and 101% at fortification level 50 mg/L	11% at 10 mg/L and 0.9 % at 50 mg/L	0.2 mg/L	██████████ (2013)

	<p>NADH formed is stoichiometric to the amount of formic acid. The increase in NADH is measured by means of its light absorbance at 334, 340 or 365 nm. The molar extinction coefficient is large at 340 nm [$\epsilon = 6.3 \text{ L}/(\text{mmol} \times \text{cm})$], i.e. the method is most sensitive at this wavelength. The extinction coefficient allows to calculate the formate concentration from the absorbance difference at the start and at the end of the reaction, which is a common method in biochemical laboratories. Photometric measurements provide the basis for the majority of quantitative methods in biochemistry and are related to the</p>								
--	---	--	--	--	--	--	--	--	--

	amount of light absorbed								
--	--------------------------	--	--	--	--	--	--	--	--

Additional remarks:

According to the guidance on residue analysis in soil "The LOQ must be below the PNEC water if technically possible". In the present case it was not technically possible to achieve an LOQ below 5 mg/L.

For drinking water it is suggested that the stringent limit and corresponding analytical LOQ of 0.1µg/L for biocides should not be relevant for formic acid. Formic acid is a naturally occurring substance, which is expected to be present in drinking water from many other, also natural sources other than only via biocide use

Methods analysis for body fluids: Body fluids was not validated as according to the guidance such method is not necessary for substances that are not toxic or very toxic (systemic toxicity)

7 EFFICACY

Products containing FORMIC ACID are intended to be used for PT3 applications as broad spectrum surface disinfectants against bacteria, yeasts, fungi.

The products are intended to be used for disinfection of animal housing (including fogging procedures), animal transportation vehicles (including tyres), of boots, of animal feet and of outsides of machinery connected with livestock farming.

In the context of a decision on the approval of FORMIC ACID for PT3 applications, 2 intended uses are considered : surface disinfection by fogging and by dipping.

In the context of a decision on the approval of FORMIC ACID, in order to assess the microbicide activity of FORMIC ACID-based products, the Applicant **BASF SE** have submitted many documents:

- Among them, a lot of documents are scientific papers with reliability 3-4. Due to lack of critical information or to data so succinctly reported, these documents are not robust enough to state efficacious concentrations usable to perform the risk assessment. Information from these documents is not taken into account and is not reported into the table below, but reported in Doc IIIB as additional information.
- Two scientific publications reviewing some information about mode of action of FORMIC ACID; one scientific publication reviewing the resistance potential of FORMIC ACID and one document giving information about pH of FORMIC ACID solutions (Document BPR_6.7_06"pH measurements of solutions of Protectol® FM85 in hard water - Technical Report BIO15_014-EX"confidential information).
- Among the remaining documents, we could find :
 - One report from efficacy tests performed according to the EN 1040 with reliability 3 due to lack of raw data. Then, these results are not taken into account and are not reported into the table below.
 - Several reports from efficacy tests performed according to EN phase 2/Step 1 EN standards (EN 1656 and EN 1657) and to EN phase 2/Step 2 EN standards (EN 14349 and EN 16438) :
Both efficacy tests have been performed on the product **Protectol® FM 85** and are summarised into the table below.
The results from the efficacy tests performed according to EN phase 2/Step 1 standards (suspension tests – i.e. EN 1656 and EN 1657) are taken into account to support basic efficacy of FORMIC ACID-based products for PT3 claims and the results from the efficacy test performed according to EN phase 2/Step 2 standards (surface tests – i.e. EN 14349 and EN 16438) is taken into account to support efficacy of FORMIC ACID-based products for "surface disinfection" PT3 claims.

7.1 EFFICACY

CONFIDENTIAL INFORMATION : Since the mode of action of Formic Acid is dependent on a low pH and could influence the efficacy of the product, refer to the *PT2 Confidential Annex* to have information about the measured pH-values of the different % of the representative product **Protectol® FM 85**. Confidential data also available in the doc. "BIO15-014-ex_pH measurements", embedded in the *PT2 Confidential Annex (p. 22)*.

Experimental data on the efficacy of the biocidal product against target organism(s)																																															
Function	Field of use envisaged	Test substance	Test organism(s)	Test method	Test system / concentrations applied / exposure time	Test results: effects	Reference																																								
Bactericidal	PT3	Protectol® FM 85 (85% formic acid)	<i>Enterococcus hirae</i> <i>Proteus vulgaris</i> <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i>	EN 1656	<u>Test concentrations</u> : 5.88 ; 3.53 and 1.17% (corresponds to 5; 3 and 1% Formic Acid) <u>Test temperature</u> : +10°C ± 1°C <u>Contact time</u> : 30 min <u>Organic loading</u> : 0.3% BSA (clean conditions)	<table border="1"> <tr> <td colspan="4">Product : <i>Protectol® FM 85</i></td> </tr> <tr> <td colspan="4">Contact time : 30 min</td> </tr> <tr> <td colspan="4">Interfering substance : 0.3% BSA (clean)</td> </tr> <tr> <td colspan="4">Test temperature : +10°C ± 1°C</td> </tr> <tr> <td>Concentration (%)</td> <td>5.88</td> <td>3.53</td> <td>1.18</td> </tr> <tr> <td colspan="4">Test Strain</td> </tr> <tr> <td>S. aureus</td> <td>5.18</td> <td>5.18</td> <td>2.30</td> </tr> <tr> <td>E. hirae</td> <td>5.38</td> <td>2.05</td> <td>0.71</td> </tr> <tr> <td>P. aeruginosa</td> <td>5.39</td> <td>5.39</td> <td>5.39</td> </tr> <tr> <td>P. vulgaris</td> <td>5.27</td> <td>5.27</td> <td>5.27</td> </tr> </table> <p>At +10°C, in suspension under clean conditions (0.3% BSA), the product Protectol® FM 85 is bactericidal in 30 min at 5.88% (5% FORMIC ACID).</p>	Product : <i>Protectol® FM 85</i>				Contact time : 30 min				Interfering substance : 0.3% BSA (clean)				Test temperature : +10°C ± 1°C				Concentration (%)	5.88	3.53	1.18	Test Strain				S. aureus	5.18	5.18	2.30	E. hirae	5.38	2.05	0.71	P. aeruginosa	5.39	5.39	5.39	P. vulgaris	5.27	5.27	5.27	Name of the document : Doc IV-BPR_6.7_04 L+S Code : 0543119 [REDACTED] (2016) <i>"Quantitative suspension test for the evaluation of bactericidal efficacy according to EN 1656"</i> Key study R.1
Product : <i>Protectol® FM 85</i>																																															
Contact time : 30 min																																															
Interfering substance : 0.3% BSA (clean)																																															
Test temperature : +10°C ± 1°C																																															
Concentration (%)	5.88	3.53	1.18																																												
Test Strain																																															
S. aureus	5.18	5.18	2.30																																												
E. hirae	5.38	2.05	0.71																																												
P. aeruginosa	5.39	5.39	5.39																																												
P. vulgaris	5.27	5.27	5.27																																												
Yeasticidal	PT3	Protectol® FM 85 (85% formic acid)	<i>Candida albicans</i>	EN 1657	<u>Test concentrations</u> : 5.88 ; 3.53 and 1.17% (corresponds to 5; 3 and 1% Formic Acid)	<table border="1"> <tr> <td colspan="4">Product : <i>Protectol® FM 85</i></td> </tr> <tr> <td colspan="4">Contact time : 30 min</td> </tr> <tr> <td colspan="4">Interfering substance : 0.3% BSA (clean)</td> </tr> <tr> <td colspan="4">Test temperature : +10°C ± 1°C</td> </tr> </table>	Product : <i>Protectol® FM 85</i>				Contact time : 30 min				Interfering substance : 0.3% BSA (clean)				Test temperature : +10°C ± 1°C				Name of the document : "1089285_1657_Version01" L+S Code : 180411-0321-001																								
Product : <i>Protectol® FM 85</i>																																															
Contact time : 30 min																																															
Interfering substance : 0.3% BSA (clean)																																															
Test temperature : +10°C ± 1°C																																															

					<p><u>Test temperature :</u> +10°C ± 1°C</p> <p><u>Contact time :</u> 30 min</p> <p><u>Organic loading :</u> 0.3% BSA (clean conditions)</p>	<table border="1"> <thead> <tr> <th>Test Concentration (%)</th> <th>5.88</th> <th>3.53</th> <th>1.18</th> </tr> </thead> <tbody> <tr> <td>Test Strain</td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>C. albicans</i></td> <td>4.37</td> <td>4.37</td> <td>0.76</td> </tr> </tbody> </table> <p>At +10°C, in suspension under clean conditions (0.3% BSA), the product Protectol® FM 85 is yeasticidal in 30 min at 3.53% (3% FORMIC ACID).</p>	Test Concentration (%)	5.88	3.53	1.18	Test Strain				<i>C. albicans</i>	4.37	4.37	0.76	<p>██████████ (2018)</p> <p>"Quantitative suspension test for the evaluation of yeasticidal efficacy according to EN 1657"</p> <p>Key study R.1</p>															
Test Concentration (%)	5.88	3.53	1.18																															
Test Strain																																		
<i>C. albicans</i>	4.37	4.37	0.76																															
Bactericidal Yeasticidal	PT3	Protectol® FM 85 (85% formic acid)	<i>Enterococcus hirae</i> <i>Proteus vulgaris</i> <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i> <i>Candida albicans</i>	<p>EN 14349 EN 16438</p> <p><u>Test concentrations :</u> 5.88; 3.53 & 1.18% (corresponds to 5; 3 and 1% Formic Acid)</p> <p><u>Test temperature :</u> +10°C ± 1°C</p> <p><u>Contact time :</u> 30 min</p> <p><u>Organic loading :</u> 0.3% BSA (clean conditions)</p>	<p>Product : <i>Protectol® FM 85</i> Contact time : 30 min Interfering substance : 0.3% BSA (clean) Test temperature : +10°C ± 1°C</p> <table border="1"> <thead> <tr> <th>Test Concentration (%)</th> <th>5.88</th> <th>3.53</th> <th>1.18</th> </tr> </thead> <tbody> <tr> <td>Test Strain</td> <td></td> <td></td> <td></td> </tr> <tr> <td><i>S. aureus</i></td> <td>5.93</td> <td>5.93</td> <td>5.93</td> </tr> <tr> <td><i>E. hirae</i></td> <td>5.82</td> <td>5.82</td> <td>2.15</td> </tr> <tr> <td><i>P. aeruginosa</i></td> <td>5.83</td> <td>5.83</td> <td>5.83</td> </tr> <tr> <td><i>P. vulgaris</i></td> <td>5.93</td> <td>5.93</td> <td>5.93</td> </tr> <tr> <td><i>C. albicans</i></td> <td>4.70</td> <td>4.70</td> <td>1.98</td> </tr> </tbody> </table> <p>At +10°C, on hard/non-porous surfaces in clean conditions (0.3% BSA), the product Protectol® FM 85 is bactericidal and yeasticidal in 30 min at 3.53% (3% FORMIC ACID).</p>	Test Concentration (%)	5.88	3.53	1.18	Test Strain				<i>S. aureus</i>	5.93	5.93	5.93	<i>E. hirae</i>	5.82	5.82	2.15	<i>P. aeruginosa</i>	5.83	5.83	5.83	<i>P. vulgaris</i>	5.93	5.93	5.93	<i>C. albicans</i>	4.70	4.70	1.98	<p>Name of the document : "0702966_14349_16438_engl_Version01" L+S Code: 160804-0106-001 ██████████ (2016)</p> <p>"Quantitative surface test for the evaluation of bactericidal according to EN 14349 and yeasticidal efficacy according to EN 16438"</p> <p>Key study R.1</p>
Test Concentration (%)	5.88	3.53	1.18																															
Test Strain																																		
<i>S. aureus</i>	5.93	5.93	5.93																															
<i>E. hirae</i>	5.82	5.82	2.15																															
<i>P. aeruginosa</i>	5.83	5.83	5.83																															
<i>P. vulgaris</i>	5.93	5.93	5.93																															
<i>C. albicans</i>	4.70	4.70	1.98																															

According to the section #4.2.2.1(p.28) of the BPR guidance (Vol. II - Parts B+C – 2018), an extensive data package and evaluation is not required at this approval stage.

As a conclusion, taking into account the results of all the efficacy tests provided by the Applicant (Phase2/Step1 suspension tests & Phase2/Step2 surface test), the product **Protectol® FM 85** is :

- Bactericidal on hard/non-porous surfaces at 5.88% (5% FORMIC ACID) at +10°C in CLEAN conditions (0.3% BSA) in 30 min
 - Yeasticidal on hard/non-porous surfaces at 3.53 % (3% FORMIC ACID) at +10°C in CLEAN conditions (0.3% BSA) in 30 min
- ⇒ For surface disinfection by dipping, both results from P2S1 & P2S2 tests should be considered and showed that the product **Protectol® FM 85** is bactericidal and yeasticidal at 5.88% (5% FORMIC ACID) in 30 min at +10°C in CLEAN conditions.
- ⇒ For surface disinfection by fogging, since EN 17272 tests are missing, only innate efficacy of the AS Formic Acid has been demonstrated. The BE eCA agreed with BASF that providing EN 17272 tests is beyond what is reasonable to submit at this present active substance Authorisation Stage. Specific P2S2 tests (performed according to the EN 17272 standard) should be provided if needed at Product Authorisation Stage.

However, for RA assessments related to surface disinfection by fogging, the application rates proposed by the APP will be used as follows : Formulated concentrate 45-55% AS

- Dilute to 4.5-5.5 % AS in the RTU solution (⇔ 5.3 % to 6.5 % of Protectol® FM 85) with 1.0 L/100 m³ for ambient temperature fogging
- Dilute to 19 % AS in the RTU solution (= 22.4% of Protectol® FM 85) with 2.25 L/1000 m³ for thermal fogging.

Considering the AS% proven effective from P2S2 efficacy tests, this way-forward is considered as acceptable for AS approval stage since AS% used for fogging are higher.

FOR INFORMATION : addition of co-formulants (surfactants for wetting/cleaning, acids for descaling, ... without impact on efficacy) could likely permit the use of lower FA concentrations.

7.2 MODE OF ACTION

The biocidal activity of Formic Acid, i.e. acidulant action and corrosion which causes enzyme denaturation and inhibition, cellular structure disruption, and impairment of cellular metabolic pathways.

This mode of action is considered to depend on the low pH-value. Secondly, formic acid does inhibit cytochrome C oxidase and thus impairs cellular energy supply. Organisms and tissues with a high energy demand are specifically susceptible :

1. Acidulant : acidification of cytoplasm;
2. Inhibitor for decarboxylases and haemin enzymes such as catalase;
3. Organic acids in general may disrupt the proton-motive force, as well as inhibit substrate transport, energy-yielding processes and macromolecular synthesis.

Acidulant action is responsible for formic acid being most effective at lower pH values (below 3.5), but enzyme inhibition and other modes also provide some antimicrobial action at higher pH values. Enzyme inhibition is less significant in the control of fungi; therefore, higher

concentrations of formic acid are needed to control fungi. The activity of formic acid against some viruses is presumably explained by the action of acid in denaturing polypeptide chains.

- Acidulant action: Organic acids cross cell membranes, leading to acidification of the cytoplasm.

- Formate inhibits cytochrome oxidase (terminal oxidase in electron transport chain), reducing ATP synthesis and thus availability of energy. Inhibition of cytochrome oxidase leads to increased production of reactive oxygen species (ROS), causing oxidative burst and damage of cell compartments. Low concentrations of formic acid were reported to induce apoptosis (-like) programmed cell death in *Saccharomyces cerevisiae* and *Candida* species.

7.3 RESISTANCE

There is no adaptation to cope with acidic pH values or denaturated proteins, nor is there a mechanism known to exist that a sub-lethal energy supply, due to an incomplete cytochrome C oxidase inhibition, would lead to undesired side-effects or resistance against this inhibitor.

No incidence of resistance to formic acid has been recorded until now.

7.4 CONCLUSION ON EFFICACY

In conclusion, the data submitted are sufficient to demonstrate efficacy of FORMIC ACID on clean hard/non-porous surfaces against bacteria (with the exception of spore-forming bacteria and mycobacteria) and yeasts for PT3 intended uses, and are therefore sufficient for the inclusion.

The data submitted are robust enough to state efficacious concentrations usable to perform the risk assessment :

The product **Protectol® FM 85** (based on 85% FORMIC ACID) is BACTERICIDAL (with the exception of spore-forming bacteria and mycobacteria) and YEASTICIDAL **at 5.88 % (⇔ 5% FORMIC ACID)** in 30 min at +10°C on hard/non-porous surfaces in clean conditions (0.3% BSA).

However, at the Product Authorisation Stage, new efficacy tests should be performed according to the requirements mentioned in the Efficacy guidance document.

8 HUMAN EXPOSURE ASSESSMENT

Default values and exposure models were taken from the document 'Biocides Human Health Exposure Methodology' and Recommendation no. 6 of the BPC Ad hoc Working Group on Human Exposure (referred to as 'Recommendation 6'), unless otherwise stated.

For the purpose of the Human Exposure Assessment for PT3, the information pertaining to the intended product concentrations and the users is summarized below. Detailed descriptions are contained in the relevant sections on exposure (8.3 – 8.8).

Intended uses

Protectol® FM 85 (85% formic acid) is used for disinfection in different premises. Prior to application it is diluted with water to the tabulated specified in-use concentrations.

Table 8.1 Overview of intended uses and in-use concentrations

Product type	Field of use envisaged	Users	Likely concentration at which a.s. will be used
PT 3.1 (Veterinary hygiene)	Disinfection of animal housings	Professional ambient temperature fogging thermal fogging	Formulated concentrate 45-55% a.s.; dilute to 4.5-5.5 % a.s. in the ready-for-use solution (= 5.3 % to 6.5 % of Protectol® FM 85); 1.0 L/100 m ³ 19 % a.s. in the ready-for-use solution (= 22.4% of Protectol® FM 85); 2.25 L/1000 m ³
PT 3.2 (Veterinary hygiene)	Disinfection of means of transport, footwear and animal feet*	Professional dipping troughs	Formulated concentrate 45-55% a.s.; dilute to 5%; a.s. in the ready-for-use solution (= 5.9% of Protectol® FM 85), unspec. volume
*though referred to as 'hoof bath disinfection' scenario in Recommendation 6, according to the applicant this use should not be limited to hooved animals.			

PT3.1 Veterinary Hygiene, Disinfection of animal housing

The biocidal product, Protectol® FM 85, is applied to disinfect areas where animals are housed. These areas are first cleaned and then disinfected on a regular basis. Disinfection of the housing includes the disinfection of containers and equipment within the area as well as the surfaces (floors, walls and ceilings) of the housing itself.

Disinfection is usually carried out as a full housing operation. Application is by the use of a fogging system. Fogging systems occur in two forms – ambient temperature fogging and

thermal fogging. Fogging systems are portable devices that can be located outside the housing.

Ambient temperature fogging

Protectol® FM 85 is formulated into the end product as supplied to the farm as an aqueous solution (approximately 45% to 55% formic acid a.s. = 52.9 % to 64.7 % of Protectol® FM 85). This is further diluted typically at 1:10 into the feed tank of a proprietary mist generator. The final concentration of the active substance is therefore 4.5% to 5.5% formic acid (= 5.3 % to 6.5 % of Protectol® FM 85). The application rate of the final dilution is typically 1.0 L/100 m³ air space. The application is controlled remotely from outside the housing being dispersed via a fan. No personnel are present in the housing undergoing disinfection during the process.

Thermal fogging

Protectol® FM 85 is formulated into products that are supplied to farms as aqueous solutions (typically containing 45% to 55% a.s. = 52.9 % to 64.7 % of Protectol® FM 85). This product as supplied is then diluted to a final use concentration of 19% formic acid (= 22.4% of Protectol® FM 85). The application rate of the product is 2.25 L of final solution /1000 m³ of air space. The product is applied via a proprietary fogging machine, which is situated outside the housing. Fogging is achieved via a flexible trunking system. No personnel are present inside the housing during the procedure.

Application frequency

Application occurs at regular (2 – 8) intervals throughout the year, preceded by a cleaning stage (a wash with detergent and drying). There is no rinsing of the area at the end of the disinfectant application. The disinfected housing is closed and allowed to air dry. There is an overnight resting stage before fresh litter is added to the housing. The housing is then ready for re-use or restocking. The resting period is a minimum of 2 hours but this is often extended to 24 –36 hours prior to restocking. In Breeding/Hatchery stock, occasional additional disinfection may take place. The wash / disinfect / rest protocol does not change but frequency may rise up to 12 regular events per year.

PT3.2 Veterinary Hygiene, Means of transport, footwear and machinery

Protectol® FM 85 is applied to disinfect transport vehicles, footwear and animal feet. This may be by the use of troughs containing disinfectant through which vehicles, the footwear of personnel or animals may pass.

Troughs for footwear disinfection, animal feet disinfection or for vehicle drive-trough

Protectol® FM 85 is formulated into end products that are supplied to farms as aqueous solutions (45-55% formic acid active substance corresponding to 52.9 to 64.7% of Protectol® FM 85). This is further diluted (to achieve concentrations of 3% to 5% formic acid = 3.5 % to 5.9 % of Protectol® FM 85) into troughs employed to disinfect footwear, as animal feet baths or for vehicle drive-through. Disinfectant solution is prepared freshly on a daily basis. Vehicles will use a drive through each time that they visit premises. This could be up to eight times per day. Similarly, footwear will be disinfected in a footbath each time that a critical area is visited. This could be several times per day.

The risk assessment will take into consideration 5% FA.

For the representative products in this CAR, secondary or indirect exposure of the general public should be avoided by implementation of appropriate RMM, considering the volatility

and corrosive properties of the a.s., and the fact that PPE and RPE are not applicable for the general public. As access of the general public to treated areas needs to be excluded by RMM, the scenario for secondary exposure will not be considered here.

Animal health:

Animal exposure assessment and measures for animal health are required at product authorisation stage.

8.1 IDENTIFICATION OF MAIN PATHS OF HUMAN EXPOSURE TOWARDS ACTIVE SUBSTANCE FROM ITS USE IN BIOCIDAL PRODUCT

Summary table: relevant paths of human exposure							
Exposure path	Primary (direct) exposure			Secondary (indirect) exposure			
	Industrial use	Professional use	Non-professional use	Industrial use	Professional use	General public	Via food
Disinfection of animal housings by fogging							
Inhalation	n.a.	Yes	n.a.	n.a.	yes	no	n.a.
Dermal	n.a.	Yes	n.a.	n.a.	yes	no	n.a.
Oral	n.a.	No	n.a.	n.a.	no	no	n.a.
Disinfection of means of transport, footwear and animal feet by dipping troughs							
Inhalation	n.a.	Yes	n.a.	n.a.	negligible	negligible	n.a.
Dermal	n.a.	Yes	n.a.	n.a.	negligible	no	n.a.
Oral	n.a.	No	n.a.	n.a.	no	no	n.a.

For Product Type 3, the biocidal product is handled and used by professionals (usually contractors, or experienced farm workers) to disinfect walls, floors and other surfaces in animal housing by fogging (PT3.1), to disinfect transport, animal feet and boots by footbaths, dipping troughs (PT3.2). Users of vehicle troughs could also be farm visitors (consumers).

There is potential for inhalation and dermal exposure to professionals; during use of vehicle troughs, inhalation is possible but unlikely for consumers.

As a result of the inherent properties of the active substance, the required activity to disinfect large areas and the method of application, it is standard practice for the workers to wear respiratory and face protection, waterproof coveralls, gauntlets and impervious boots when contact with formic acid concentrate is possible. The maximum concentration of formic acid in the applied in-use solution is 5.5% for ambient temperature fogging and 19% for thermal

fogging. For disinfection of transport, animal feet and boots by footbaths and dipping troughs, 5% FA in-use solution is considered.

The most significant potential source of secondary exposure is to farm hands restocking disinfected animal housings.

The assessment of exposure towards formic acid as active substance in product type 3 disinfectants is based on information provided by the applicant. Possible gaps are bridged by the Rapporteur using reasonable assumptions. For lack of measurement data, exposure models are applied.

In view of the high vapour pressure of Formic Acid (4271 Pa for 99% formic Acid at 20°C), exposure to vapours should be assessed when relevant for the scenario. eCA BE uses the ConsExpoWeb Exposure to Vapour model, taking into account the in-use dilution concentration and the vapour pressure of the pure active substance. The applicant prefers to use the estimated vapour pressure of the in-use dilution. However, since applying the vapour pressure of the pure active substance is a reasonable-worst-case calculation, use of the pure active substance's vapour pressure should be maintained, at least as a first tier approach.

8.2 LIST OF SCENARIOS

Summary table: scenarios			
Scenario number	Scenario (e.g. mixing/loading)	Primary or secondary exposure Description of scenario	Exposed group (e.g. professionals, non-professionals, bystanders)
1.	Animal house disinfection by fogging	1a.primary exposure during mixing and loading by professional contractors	Professionals: professional contractors
		1b.application of the in use solution by fogging, manipulated from outside	
		1c. cleaning of equipment and disposal of containers	
2.	Disinfection of footwear in dipping troughs	2a.primary exposure during mixing and loading by experienced farm workers in dipping troughs	Professionals: experienced farm workers
		2b.passive use of footbath/trough by experienced farm workers	
		2c.draining of disinfection solution and disposal of containers by experienced farm workers	
3.	Animal feet disinfection and disinfection of transport vehicles	3a.primary exposure during mixing and loading by experienced farm workers in animal feet baths/vehicle troughs	Professionals: experienced farm workers
		3b.passive animal use of animal feet bath, passive use of vehicle trough by experienced farm workers	

		3c.draining of disinfection solution and disposal of containers by experienced farm workers	
4.	Reuse/restocking of disinfected animal housing	Secondary exposure while opening animal housings sealed for treatment (fogging)	Professionals: experienced farm workers

8.3 INDUSTRIAL EXPOSURE

This section has not been evaluated by the CA-BE because the production/formulation process of the active substance is outside the scope of the Biocidal Products Regulation (EU) No 528/2012.

Protectol® FM 85 is manufactured on the BASF SE site in D-67058 Ludwigshafen, Germany. Exposure of manufacturing workers is governed by industrial legislation and controlled by the use of automated processes. The active substance is rigorously contained by production methods and the use of personal protective equipment so that direct exposure of manufacturing workers is prevented.

Formic acid is produced in a production plant and further processed within other operations. Formic acid is produced within a closed system. A total of 138 workplace measurements have been conducted during the period 2001-2006, covering all kinds of operations (production, filling, processing, laboratory). All reported results represented 8 hours shift average values (TWA) obtained by personal air sampling. None of the measurements exceeded the threshold limit of 5 ppm or 9.6 mg/m³ (most well below). To prevent direct skin contact, protective gloves (neoprene or nitrile rubber) must be used. According to the applicant workplace exposure is low, due to the appropriate protective measures taken (DocIIIA6.12.1-01: Eisenbarth, 2006).

Four cases of accidental skin and eye contact were seen during 14 years (1989-2002) of operation of the BASF's production plant. Lesions of skin and eye were seen following facial splashes (3 cases) during filling operations and transportation, and one case of skin lesions following contact with contaminated wood (DocIIIA6.12.3-01: ██████████ 1994, 2002).

Nevertheless, exposure estimates for industrial workers during these stages have not been calculated as they are already addressed by other legislation. Therefore, in accordance with the Commission Document agreed at the 22nd CA meeting in September 2006, detailed information on exposure associated with the manufacturing process is not required for biocidal product risk assessment.

8.4 PROFESSIONAL EXPOSURE

The biocidal product available for professional contractors and experienced farm workers is a concentrated product containing 45-55% formic acid, formulated starting from Protectol® FM 85 (85% FA), and to be further diluted to the recommended use concentration of 5% for dipping troughs, or to the recommended use concentrations of 4.5-5.5% to 19% for fogging. Professionals use products on a prolonged basis, mixing and loading, fogging. Professional contractors are exposed daily to loading fogging machines, experienced farm workers are exposed daily to dipping troughs.

General default values:

parameter	Default value
Body weight adult (prof/consumer)	60 kg
Respiration rate adult	1.25 m ³ /h
Oral absorption	100%
Dermal absorption	100%
Inhalation absorption	100%

PRIMARY EXPOSURE

8.4.1 Scenario 1: animal house disinfection by fogging

This scenario involves the following subscenarios:

- 1a. mixing and loading by professional contractors
- 1b. application of the in use solution by fogging, manipulated from outside
- 1c. cleaning of equipment and disposal of containers

Description of Scenario 1a

Task, exposure model and parameters:

1a. mixing and loading by professional contractors

Manual opening and emptying of containers into a reservoir and dilution with water

Concentration of a.s. in biocidal product: 55%

Density of product: ca. 1200 g/L

Frequency: daily for professional contractors

Duration of exposure: 10 min

Emission duration: 5 min⁽¹⁾

Ventilation rate: 10/h⁽¹⁾

Room volume (M&L): 24 m³ ⁽¹⁾

Volume treated: 2810 m³ ⁽²⁾

Amount of product (55% concentrate) handled: ca. 3 kg ⁽³⁾

Amount of active substance handled: ca. 1.65 kg

Release area: 100 cm² ⁽¹⁾

Exposed worker: professional contractors

Protective equipment: impermeable coveralls, boots, gloves and face protection⁽⁵⁾

Model: EUROPOEM II database, liquid manual loading/pouring (dermal only)

ConsexpoWeb, evaporation, area of release constant⁽⁴⁾

	Parameters*	Value
Tier 1	Hands	8.0 mg/kg a.s.
	body	1.95 mg/kg a.s.
Tier 2	Protection gloves	90%
	Protection impermeable coveralls	95%

⁽¹⁾ defaults taken from Recomm. 6 v3, scenario 12 hoof bath disinfection mixing & loading

⁽²⁾ ENV/JM/MONO(2006)4, Table 5.2, defaults for animal housings, standard poultry housing.

⁽³⁾ Max application rate of in-use solution for fogging is 1L/100m³. For 2810 m³ housing, approx.. 28L of in-use solution is needed. This 5.5% in-use solution is prepared from a 55% concentrate (10-fold dilution) so from 2.8L concentrate. Taking into account its density, approx. 3 kg concentrate is needed.

⁽⁴⁾ Mass transfer coefficient: ConsExpo default of 10m/hr.

⁽⁵⁾ See applicant's SDS for 85% FA, section 8 Exposure controls/personal protection

Calculations for Scenario 1a

Model: EUROPOEM II database, liquid manual loading/pouring

Tier 1:

55% concentrate, dermal exposure:

$$[(8.0 \text{ mg/kg} \cdot 1.65 \text{ kg}) + (1.95 \text{ mg/kg} \cdot 1.65 \text{ kg})] / 60 \text{ kg} = 16.418 \text{ mg/kg bw per shift}$$

Tier 2:

55% concentrate, dermal exposure:

$$[(8.0 \text{ mg/kg} \cdot 1.65 \text{ kg} \cdot 0.1) + (1.95 \text{ mg/kg} \cdot 1.65 \text{ kg} \cdot 0.05)] / 60 \text{ kg} = 0.0247 \text{ mg/kg bw per shift}$$

Model: ConsexpoWeb, evaporation, area of release constant**Tier 1, exposure to vapour:**

For full ConsExpo reports see Appendix II

Mean event concentration 3.8 mg/m³

Peak concentration (TWA 15 min) 3.8 mg/m³

Year average concentration 2.6×10^{-2} mg/m³

External event dose 1.3×10^{-2} mg/kg bw

Internal event dose 1.3×10^{-2} mg/kg bw

Internal year average dose 1.3×10^{-2} mg/kg bw/day

Description of Scenario 1b

Tasks, exposure models and parameters:

1b. application of the in use solution by fogging, manipulated from outside

Ambient temperature or thermal fogging of animal housing in the absence of personnel.

Concentration of a.s. in diluted biocidal product:

Ambient temperature fogging: typically 5.5%

Thermal fogging: 19%

Application rate:

Ambient temperature fogging: 1.0 L per 100 m³

Thermal fogging: 2.25 L per 1000 m³

Density of product: 5.5% FA: ca. 1000 g/L; 19%: ca. 1200 g/L (worst case)

Frequency: daily for professional contractors

Duration of exposure: not relevant, treatment of animal housing in absence of personnel

Exposed worker: professional contractor

Protective equipment: not relevant

Model: not relevant; the application of the product is controlled remotely from outside the housing. The product is dispersed via a fan or fogging is achieved via a flexible trunking system. No personnel are present in the housing undergoing disinfection during the process.

Therefore no calculations are provided for scenario 1b.

Description of Scenario 1c

Tasks, exposure models and parameters:

1c. cleaning of equipment and disposal of containers

Disposal of emptied containers after filling of equipment, and maintenance of equipment after use

Concentration of a.s. in biocidal product: 55%

Concentration of a.s. in diluted biocidal product:

 Ambient temperature fogging: typically 5.5%

 Thermal fogging: 19%

Density of product: 5.5% FA: ca. 1000 g/L; 19%: ca. 1200 g/L (worst case)

Frequency: daily for professional contractors

Duration of exposure: N.A. (see below)

Exposed worker: professional contractors

Protective equipment: impermeable coveralls, boots, gloves and face protection

Model: not relevant: we assume that exposure during the cleaning and disposal step is covered by the mixing & loading step. Prepared fogging solution is assumed to be used up during the application phase.

Therefore no calculations are provided for scenario 1c.

Further information and considerations on scenario 1

Because during fogging activities nobody is inside the building, it is considered that only the mixing and loading phase will be used as the basis for risk assessment for this scenario.

Personal Protective Equipment (PPE) incorporating impermeable coveralls, boots, gloves and face protection is used during the mixing and loading phase and will significantly reduce exposure via the dermal and inhalation routes. As a tier 2, protection factors for gloves and impermeable coveralls have been used. In situations where ventilation is insufficient, appropriate RPE would be used due to the acidity of the formic acid fumes released during mixing and loading of the product.

Exposure to vapour during mixing and loading was calculated with the ConsExpo – exposure to vapour – evaporation scenario. The amount handled was calculated for a standard poultry battery (ENV/JM/MONO(2006)4). Defaults given for application duration, ventilation rate, room volume and release area for the PT hoof bath disinfection model (Recomm. 6, v3, model no. 12) were adopted, as it was supposed that conditions for mixing and loading the in-use solution for fogging would be similar to those for hoof bath disinfection. Exposure to vapour should be reduced by ventilation and other appropriate risk mitigation measures.

The resting period is a minimum of 2 hours but this is often extended to 24-36 hours prior to restocking. No ventilation phase is foreseen before reuse/restocking of the animal housing.

For a graphic representation of the Formic Acid air concentration during mixing and loading, see Appendix II graph II.1.

(Semi-)quantitative assessment for oral, dermal and inhalation routes

Results table exposure to PT3 disinfection of animal housing by fogging						
Exposure subscenario	Tier/PE	Estimated inhalation uptake (mg/kg bw/d)	Estimated dermal uptake (mg/kg bw/d)	Estimated total uptake (mg/kg bw/d)	Local dermal exposure (conc., %)	Local inhalation exposure (mg/m³)
1a M&L 55%	1/none	0.013 (ConsExpo vapour)	16.418	16.431	55	3.8 (vapour)
	2/impermeable coveralls, boots, gloves and face protection	0.013 (ConsExpo vapour)	0.0247	0.0377	55	3.8 (vapour)
1b application 5.5% or 19%	N.A.(**)	-	-	-	-	-
1c cleaning & disposal	N.A.(*)	-	-	-	5,5% - 19%	-

(*) covered by subscenario 1a Mixing & loading

(**) no personnel present

Qualitative local assessment for dermal route

As formic acid is corrosive at or above a 10% dilution, a qualitative risk characterisation is needed for local dermal exposure. This RC is triggered for those BP classified for local effects. In BP where formic acid is present at concentrations that do not trigger classification of the product according to the CLP criteria, RC for local effects is not required.

The concentrate (55% FA) for PT3 professional use is classified as corrosive to the skin, cat. 1B. This classification triggers a qualitative local assessment for the dermal route. We refer to section 12.4.2 for relevant RMM end PPE and the conclusion on the acceptability of the risk.

The in-use dilutions (5.5% and 19% FA) have not been included in the qualitative assessment for the dermal route as no dermal exposure to these dilutions is foreseen.

8.4.2 Scenario 2: disinfection of footwear in dipping troughs

Disinfection of footwear by footbath use.

This scenario involves the following subscenarios:

- 2a. mixing and loading by experienced farm workers in dipping troughs
- 2b. passive use of footbath by experienced farm workers
- 2c. draining of disinfection solution and disposal of containers by experienced farm workers.

Description of Scenario 2a2a. mixing and loading by experienced farm workers in dipping troughs

Manual opening and emptying of containers into a reservoir and dilution with water

Concentration of a.s. in biocidal product: 55%

Concentration of a.s. in diluted product: 5%

Density of concentrated product: ca. 1200 g/L

Frequency: 104 times per year

Duration of exposure: 10 min

Emission duration: 5 min⁽¹⁾

Ventilation rate: 10/h⁽¹⁾

Room volume: 24 m³ ⁽¹⁾

Release area: 100 cm² ⁽¹⁾

Volume loaded: 10L⁽²⁾

Amount of product (55% concentrate) handled:

To attain in-use dilution of 5% FA: 1.09 kg⁽³⁾

Amount handled, weight, pure a.s.:

To attain in-use dilution 5% FA: ca. 0.6 kg

Exposed worker: professional: experienced farm workers, professionals whose main job is not principally related to disinfection

Protective equipment: coated coveralls, boots, gloves and face protection

Model: EUROPOEM II database, Manual loading/pouring volumes up to 20L (dermal only)

ConsexpoWeb, evaporation, area of release constant⁽⁴⁾

	Parameters*	Value
Tier 1	Indicative value hands :	8 mg/kg a.s.
	Indicative value body :	1.95 mg/kg a.s.
Tier 2**	PPE: Coveralls, boots, gloves,	
	Probability of glove penetration	10%
	Probability of clothing penetration	10%

⁽¹⁾ defaults taken from Recomm. 6, model 12 scenario for hoof bath disinfection mixing & loading

⁽²⁾ BHHEM, default for volume of bath for footwear

⁽³⁾ dilute 55% concentrate to 5% in-use solution or 11-fold dilution; 909 ml concentrate needed to prepare 10L in-use solution; considering the density of the concentrate, approx. 1090g is needed to prepare 10L.

⁽⁴⁾ Mass transfer coefficient: consExpo default 10 m/hr.

Calculations for Scenario 2a

amount handled, weight, pure a.s.: ca. 0.6 kg

Model: EUROPOEM II database, Manual loading/pouring volumes up to 20L

Tier 1:

In-use dilution 5%:

55% concentrate, dermal exposure:

$$[(0.6 \text{ kg} * 8 \text{ mg/kg a.s.}) + (0.6 \text{ kg} * 1.95 \text{ mg/kg a.s.})] / 60 \text{ kg} = 0.0995 \text{ mg/kg bw}$$

Tier 2:

In-use dilution 5%:

55% concentrate, dermal exposure:

$$[(0.6 \text{ kg} * 8 \text{ mg/kg a.s.} * 0.1) + (0.6 \text{ kg} * 1.95 \text{ mg/kg a.s.} * 0.1)] / 60 \text{ kg} = 0.00995 \text{ mg/kg bw}$$

Model: ConsexpoWeb, evaporation, area of release constant

Tier 1, exposure to vapour:

For full ConsExpo reports see Appendix II

In-use dilution 5%

Mean event concentration 3.8 mg/m³

Peak concentration (TWA 15 min) 3.8 mg/m³

Year average concentration 7.5×10^{-3} mg/m³

External event dose 1.3×10^{-2} mg/kg bw

Internal event dose 1.3×10^{-2} mg/kg bw

Internal year average dose 3.8×10^{-3} mg/kg bw/day

Description of Scenario 2b

2b. passive use of footbath by experienced farm workers

Dipping of boots in footbath

Concentration of a.s. in diluted product: 5%

Density of product: ca. 1000 g/L

Frequency: several times per day

Duration of exposure: 30 sec per day

Exposed worker: experienced farm workers, professionals whose main job is not principally related to disinfection

Protective equipment: work clothing

Model: not relevant.

Use of the footbath is limited to walking in + soaking boots in the bath; short exposure, negligible dermal contact and inhalation.

Therefore no calculations are provided for scenario 2b.

Description of Scenario 2c

2c. draining of disinfection solution and disposal of containers by experienced farm workers

Pouring of used footbaths to slurry pits, disposal of emptied containers

Concentration of a.s. in diluted product: 5%

Frequency: 104 times per year

Duration of exposure: 10 min per application day (104d)

Volume removed: 10L (worst case)

Amount handled, weight, pure a.s.: ca. 0.6 kg, see scenario 2a

Exposed worker: experienced farm workers, professionals whose main job is not principally related to disinfection

Protective equipment: coated coveralls, boots, gloves and face protection

Model: EUROPOEM II database, Manual loading/pouring volumes up to 20L

For model defaults and calculations, see scenario 2a.

As a worst case, it was assumed that during post-application the worker is exposed to a fully filled trough. It was assumed that the mixing and loading model is also appropriate for draining and disposal, and that exposure is equivalent to the exposure during the mixing & loading phase.

However, inhalation of vapour as calculated for M&L is considered less relevant for disposal. During disposal a dilute is handled with considerably lower % of FA. Therefore inhalation was not taken into consideration for the draining of the diluted solution. Also, a substantial part the formic acid may have evaporated out of the dilution before the disposal step.

Calculations for Scenario 2c

amount handled, weight, pure a.s.: ca. 0.6 kg

Model: EUROPOEM II database, Manual loading/pouring volumes up to 20L

Tier 1:

In-use dilution 5%:

55% concentrate, dermal exposure:

$$[(0.6 \text{ kg} * 8 \text{ mg/kg a.s.}) + (0.6 \text{ kg} * 1.95 \text{ mg/kg a.s.})] / 60 \text{ kg} = 0.0995 \text{ mg/kg bw}$$

Tier 2:

In-use dilution 5%:

55% concentrate, dermal exposure:

$$[(0.6 \text{ kg} * 8 \text{ mg/kg a.s.} * 0.1) + (0.6 \text{ kg} * 1.95 \text{ mg/kg a.s.} * 0.1)] / 60 \text{ kg} = 0.00995 \text{ mg/kg bw}$$

Further information and considerations on scenario 2

Dermal exposure is expected to occur during mixing and loading and emptying of used footbath tubs. Personal Protective Equipment (PPE) incorporating coated coveralls, boots, gloves and face protection will significantly reduce exposure via the dermal route. The TNsG on Human Exposure to Biocidal Products (2002, Part 2, page 36) indicates that 10% clothing penetration is the standard default in lieu of actual or model data based on information from agrochemical uses.

For dermal exposure during mixing & loading as well as during post-application, the EUROPOEM II model for Manual loading/pouring volumes up to 20L was chosen. As a worst case, it was assumed that during post-application the worker is exposed to a fully filled trough; therefore dermal exposure is equivalent to the exposure during the mixing & loading phase. No dermal exposure was calculated for use of the footbath to disinfect footwear as its use is limited to walking through and soaking footwear in the bath; negligible dermal contact is expected.

Inhalation exposure is assumed during the mixing and loading of the concentrate. In situations where ventilation is insufficient, appropriate RPE would be used due to the acidity of the formic acid fumes released during mixing and loading of the product.

Taking into account the volatility of formic acid, exposure to vapour during mixing and loading was calculated with the ConsExpo – exposure to vapour – evaporation scenario. Defaults given for application duration, ventilation rate, room volume and release area for the PT hoof bath disinfection model (Recomm. 6, v3, model 12) were adopted as it was supposed that conditions for mixing and loading would be similar. Refinements for this exposure estimate can be used at product authorisation. In any case, exposure to vapour should be reduced by ventilation and other appropriate risk mitigation measures. No inhalation exposure was calculated for use of the footbath to disinfect footwear. Exposure duration is short; exposure to vapours is considered negligible. Inhalation of vapour was also not taken into consideration for the draining of the diluted solution; a substantial part the formic acid may have evaporated out of the dilution before the disposal step.

For a graphic representation of the Formic Acid air concentration during mixing and loading, see Appendix II graph II.2.

(Semi-)quantitative assessment for oral, dermal and inhalation routes

Results table exposure to PT3 disinfection of footwear in dipping troughs						
Exposure subscenario	Tier/PP E	Estimated inhalation uptake (mg/kg bw/d)	Estimated dermal uptake (mg/kg bw/d)	Estimated total uptake (mg/kg bw/d)	Local dermal exposure (conc., %)	Local inhalation exposure (mg/m³)

2a M&L 55%, in-use dilution 5% FA	1/none	0.013 (ConsExpo vapour)	0.0995	0.1125	55	3.8 (vapour)
	2/coated coveralls, boots, gloves and face protection	0.013 (ConsExpo vapour)	0.00995	0.0230	55	3.8 (vapour)
2b application 5%	N.A.(*)	-	-	-	-	-
2c draining & disposal 5%	1/none	-	0.0995	0.0995	5	-
	2/coated coveralls, boots, gloves and face protection	-	0.00995	0.00995	5	-
2 disinfection of footwear, total	1/none	0.013 (ConsExpo vapour)	0.199	0.212	55 & 5	3.8 (vapour)
	2/coated coveralls, boots, gloves and face protection, respirator	0.013 (ConsExpo vapour)	0.0199	0.0329	55 & 5	3.8 (vapour)

(*) exposure considered negligible

Qualitative local assessment for dermal route

As formic acid is corrosive at or above a 10% dilution, a qualitative risk characterisation is needed for local dermal exposure. This RC is triggered for those BP classified for local effects. In BP where formic acid is present at concentrations that do not trigger classification of the product according to the CLP criteria, RC for local effects is not required.

The concentrate (55% FA) for PT2 professional use is classified as corrosive to the skin, cat. 1B. Some in-use dilutions (2-10% FA) are classified as skin and eye irritant cat. 2. These classifications trigger a qualitative local assessment for the dermal route. We refer to section 12.4.2 for relevant RMM end PPE and the conclusion on the acceptability of the risk.

8.4.3 Scenario 3: animal feet disinfection and disinfection of transport vehicles

Animal feet disinfection by hoof bath use & disinfection of vehicles in troughs for vehicle drive-through.

This scenario involves the following subscenarios:

3a. mixing and loading of the substance in the hoof bath/vehicle trough by experienced farm workers

3b. walking through of livestock in hoof bath, passive use of vehicle trough by experienced farm workers

3c. draining of disinfection solution and disposal of containers by experienced farm workers.

Description of Scenario 3

Manual opening and emptying of containers into a reservoir and dilution with water. Passive disinfection of animal feet. Draining of disinfection solution and disposal of containers.

Concentration of a.s. in biocidal product: 55%

Concentration of a.s. in diluted product: 5%

Density of concentrated product: ca. 1200 g/L

Density of diluted product: ca. 1000 g/L

Frequency: daily

Application duration: 5 min for M&L, 35 min for application, 10 min for post-application ⁽¹⁾

Ventilation rate & room volume: 10/h, 24 m³ for M&L; 2/h, 9630 m³ for (post)application ⁽¹⁾

Release area: 100 cm² for M&L, 3 m² for (post)application ⁽¹⁾

Volume of trough: 1000 L ⁽²⁾

Amount of product (55% concentrate) handled:

For in-use dilution of 5 % FA: ca. 110 kg ⁽³⁾

Amount of dilute solution exposed to: ca. 1000 kg

Exposed worker: professional: experienced farm workers, professionals whose main job is not principally related to disinfection

Protective equipment: coveralls, boots, gloves, face protection ⁽⁵⁾

Models: dermal exposure: Mixing & loading Model 4 (TNsG 2002) for M&L and (post)application

Inhalation exposure: ConsexpoWeb, evaporation, area of release constant⁽⁴⁾

	Parameters*	Value
Tier 1	Indicative value dermal exposure, hands only :	0.5 mL b.p./loading
	Indicative nr. of loadings:	1 loading for M&L, 4 for post-application
Tier 2**	PPE: Coveralls, boots, gloves	
	Probability of glove penetration	10%

⁽¹⁾ defaults taken from Recomm. 6 v3, model 12 scenario hoof bath disinfection

⁽²⁾ BHHM, default for volume of hoof bath

⁽³⁾ in-use dilution 5% FA: dilute 55% concentrate to 5% in-use solution or 11-fold dilution; 91L concentrate needed to prepare 1000L in-use solution; considering the density of the concentrate, approx. 110kg is needed to prepare 1000L.

⁽⁴⁾ Mass transfer coefficient ConsExpo default 10 m/hr

⁽⁵⁾ See applicant's SDS for 85% FA, section 8 Exposure controls/personal protection

Calculations for Scenario 3

amount handled, weight, 55% concentrate.: ca. 110 kg

Model: Mixing & loading Model 4 (TNsG 2002)

Tier 1:

55% concentrate, dermal exposure:

0.5 mL b.p./loading = ca. 0.6 g b.p./loading

$(0.6 \text{ g/loading} * 5 \text{ loadings} * 0.55)/60\text{kg} = 0.0275 \text{ mg/kg bw}$

Tier 2:

55% concentrate, dermal exposure:

$(0.6 \text{ g/loading} * 5 \text{ loadings} * 0.55 * 0.1)/60\text{kg} = 0.00275 \text{ mg/kg bw}$

Model: ConsexpoWeb, evaporation, area of release constant

Tier 1, exposure to vapour during M&L:

For full ConsExpo reports see Appendix II

	In-use dilution 5% FA
Mean event concentration	3.5 mg/m ³
Peak concentration (TWA 15 min)	3.5 mg/m ³
Year average concentration	$1.2 \times 10^{-2} \text{ mg/m}^3$
External event dose	$6.1 \times 10^{-3} \text{ mg/kg bw}$
Internal event dose	$6.1 \times 10^{-3} \text{ mg/kg bw}$

Internal year average dose	6.1×10^{-3} mg/kg bw/day
----------------------------	--------------------------------------

Tier 1, exposure to vapour during application:

	In-use dilution 5% FA
Mean event concentration	1.0 mg/m ³
Peak concentration (TWA 15 min)	1.5 mg/m ³
Year average concentration	2.5×10^{-2} mg/m ³
External event dose	1.3×10^{-2} mg/kg bw
Internal event dose	1.3×10^{-2} mg/kg bw
Internal year average dose	1.3×10^{-2} mg/kg bw/day

Tier 1, exposure to vapour during post-application:

	In-use dilution 5% FA
Mean event concentration	3.8×10^{-1} mg/m ³
Peak concentration (TWA 15 min)	3.8×10^{-1} mg/m ³
Year average concentration	2.6×10^{-3} mg/m ³
External event dose	1.3×10^{-3} mg/kg bw
Internal event dose	1.3×10^{-3} mg/kg bw

Internal year average dose 1.3×10^{-3} mg/kg bw/day

Further information and considerations on scenario 3

Exposure models and calculations as presented here are based on data for hoof bath disinfection. It can be assumed that exposure to vehicle troughs will not be higher than exposure during use of hoof baths. Experienced farmers are exposed to hoof baths; vehicle troughs can also be used by consumers but exposure is considered negligible.

Dermal exposure is expected to occur during mixing and loading and emptying of used hoof baths, particularly to hands. Personal Protective Equipment (PPE) incorporating coveralls, boots, gloves and face protection will significantly reduce exposure via the dermal route. 10% glove penetration is the standard default used (BHHM).

For inhalation exposure, taking into account the volatility of formic acid, calculations were done following the ConsExpo – exposure to vapour – evaporation scenario for the mixing & loading, application and post-application phase. Defaults given for application duration, ventilation rate, room volume and release area for the PT3 hoof bath disinfection model (Recomm. 6 v3, scenario 12) were followed. Further refinements for this exposure estimate can be used at product authorisation. In any case, exposure to vapour should be reduced by ventilation and other appropriate risk mitigation measures. In situations where ventilation is insufficient during mixing and loading (handling of concentrate), appropriate RPE would be used due to the acidity of the formic acid fumes released.

(Semi-)quantitative assessment for oral, dermal and inhalation routes

Results table exposure to PT3 animal feet/transport vehicle disinfection						
Exposure scenario	Tier/PP E	Estimated inhalation uptake (mg/kg bw/d)	Estimated dermal uptake (mg/kg bw/d)	Estimated total uptake (mg/kg bw/d)	Local dermal exposure (conc., %)	Local inhalation exposure (mg/m³)
3a M&L 55%, for in-use dil 5%	1/none	6.1×10^{-3} (ConsExpo vapour)	0.0275	0.0336	55	3.5 (vapour)

	2/coveralls, boots, gloves; face protection	$6.1 \cdot 10^{-3}$ (ConsExpo vapour)	0.00275	0.0089	55	3.5 (vapour)
3b application 5%	1/none	$1.3 \cdot 10^{-2}$ (ConsExpo vapour)	Covered by M&L	$1.3 \cdot 10^{-2}$	5	1.0 (vapour)
	2/coveralls, boots, gloves; face protection	$1.3 \cdot 10^{-2}$ (ConsExpo vapour)	Covered by M&L	$1.3 \cdot 10^{-2}$	5	1.0 (vapour)
3c draining & disposal 5%	1/none	$1.3 \cdot 10^{-3}$ (ConsExpo vapour)	Covered by M&L	$1.3 \cdot 10^{-3}$	5	0.38 (vapour)
	2/coveralls, boots, gloves; face protection	$1.3 \cdot 10^{-3}$ (ConsExpo vapour)	Covered by M&L	$1.3 \cdot 10^{-3}$	5	0.38 (vapour)
PT3 animal feet disinfection totalled						
3 M&L, application 5%, post-application	1/none	$2.0 \cdot 10^{-2}$ (ConsExpo vapour)	0.0275	0.0475	55 (M&L) 5 ((post)appl)	3.5 (M&L) 1.0 (appl) 0.38 (post)
3 M&L, application 5%, post-application	2/coveralls, boots, gloves; face protection	$2.0 \cdot 10^{-2}$ (ConsExpo vapour)	0.00275	0.0228	55 (M&L) 5 ((post)appl)	3.5 (M&L) 1.0 (appl) 0.38 (post)

Qualitative local assessment for dermal route

As formic acid is corrosive at or above a 10% dilution, a qualitative risk characterisation is needed for local dermal exposure. This RC is triggered for those BP classified for local effects.

In BP where formic acid is present at concentrations that do not trigger classification of the product according to the CLP criteria, RC for local effects is not required.

The concentrate (55% FA) for PT2 professional use is classified as corrosive to the skin, cat. 1B. Some in-use dilutions (2-10% FA) are classified as skin and eye irritant cat. 2. These classifications trigger a qualitative local assessment for the dermal route. We refer to section 12.4.2 for relevant RMM end PPE and the conclusion on the acceptability of the risk.

SECONDARY EXPOSURE

8.4.4 Scenario 4: restocking of animal housing after disinfection

Secondary exposure may occur from a farm hand opening and restocking disinfected animal housings. Indirect inhalation exposure is possible for the farm hand.

During disinfection of animal housing by fogging the treated housing is sealed tightly. There is a resting stage before fresh litter is added to the housing. The housing is then ready for reuse or restocking. The resting period as recommended by the applicant is a minimum of 2 hours but this is often extended to 24-36 hours prior to restocking.

Farm hands opening and restocking the treated animal house are exposed to the formic acid-containing vapour by inhalation. Dermal contact is possible but unlikely due to the volatility of the active substance.

As a worst case approach, all the applied formic acid evaporates and remains in the housing unit, and no additional ventilation during the resting stage is assumed. Next the effect of ventilation is assessed.

Description of Scenario 4

Scenario: indirect exposure, farm hand, restocking of animal housing after disinfection

Frequency: up to 12 times/year

Duration of exposure: 1 h⁽¹⁾

Concentration of a.s.: 5.5% (ambient temperature fogging), 19% (thermal fogging)

Amount applied: 1L/100 m³ (ambient temperature fogging), 2.25 L/1000 m³ (thermal fogging)

Room size: 2810 m³, surface area 1850 m² ⁽²⁾

⁽¹⁾ Expert judgement.

⁽²⁾ ENV/JM/MONO(2006)4, Table 5.2, defaults for animal housings, standard poultry housing.

Calculations for Scenario 4

Tier 1 Exposure to vapour, all formic acid evaporated

There is potential for inhalation exposure as formic acid is volatile and would be expected to evaporate as the cleaned surface dries. A standard poultry shed has a volume of 2810 m³ and a surface area of 1850 m² (ENV/JM/MONO(2006)4). For inhalation, the worst-case situation would be to assume that all of the applied formic acid had evaporated and remained in the housing unit.

	Ambient temperature fogging	Thermal fogging
Surface or volume treated	2810 m ³	2810 m ³
Concentration	5.5%	19%
Application rate product	1L/100 m ³	2.25L/1000 m ³
Application rate a.s.	0.055 kg/100 m ³	0.43 kg/1000 m ³
Concentration of formic acid in air	550 mg/m ³	430 mg/m ³
internal dose on day of exposure	11.5 mg/kg/d	9.0 mg/kg/d

Worst-case, the farm hand would be exposed to a concentration of formic acid in air of 550 and 430 mg/m³ (for ambient temperature and thermal fogging, respectively) when opening a treated poultry shed.

It is assumed that the farm hand may be present in the housing unit for one hour for restocking, and an inhalation rate of 1.25 m³/h is used based on short-term exposure values for an adult (Recommendation no. 14 (2017) of the BPC Ad hoc WG on HE, Default human factor values for use in exposure assessments for biocidal products). Exposure is therefore equivalent to 11.5 and 9.0 mg/kg bw/d for ambient temperature and thermal fogging, respectively, for a 60 kg adult entering a treated animal housing.

This approach is considered very conservative. In practice, concentrations in air would be reduced due to ventilation once the shed is opened.

Exposure to vapour, ventilation time needed for safe re-entry

The ConsExpo model exposure to vapour, instantaneous release was used to assess the decrease of formic acid concentrations in air due to ventilation, and the time needed for safe re-entry. Two ventilation rates were considered: a conservative ventilation rate of 2/h, and a realistic ventilation rate of 10/h

(see appendix II graph II.4).

	Ambient temperature fogging	Thermal fogging
	T1 ventilation rate 2/h T2 Ventilation rate 10/h	
volume treated	2810 m ³	
Formic acid concentration in dilution	5.5%	19%

Application rate product	1L/100 m ³	2.25L/1000 m ³
Application rate a.s.	0.055 kg/100 m ³	0.43 kg/1000 m ³
Starting concentration of formic acid vapours in air	550 mg/m ³	430 mg/m ³
Required ventilation time to reach <6 mg/m ³ *	T1 136 min T2 28 min	T1 129 min T2 26 min
internal dose on day of exposure	0.125 mg/kg/d	

*AEC_{respiratory irritation} for formic acid

For animal housing treated by fogging, it would take just over 2h of ventilation for for the air concentration of formic acid to drop to levels below the AEC for respiratory irritation of 6 mg/m³ (fogging at ambient temperature starting from 550 mg/m³ FA: 136 min, thermal fogging starting from 430 mg/m³ FA: 129 min; see appendix II graph II.4) when a ventilation rate of 2/hour is considered. At a ventilation rate of 10/hour, 30 mins of ventilation would suffice to reduce the air concentration of formic acid appropriately (fogging at ambient temperature: 28 min, thermal fogging: 26 min; see appendix II graph II.4).

Assuming that the necessary ventilation phase to reach <6 mg/m³ is respected, and that the farm hand may be present in the housing unit for one hour for restocking, and when an inhalation rate of 1.25 m³/h is used (Recommendation no. 14 (2017) of the BPC Ad hoc WG on HE, Default human factor values for use in exposure assessments for biocidal products), total exposure would amount to 7.5 mg/d, equivalent to 0.125 mg/kg/d for a 60 kg adult.

Exposure to vapour, RMM approach

It could be argued that ConsExpo is not appropriate to calculate safe re-entry times after fogging. In that case, the following RMM is suggested:

After disinfection by fogging, the treated area can be re-entered only when an ambient air concentration of formic acid is ensured to be below 6 mg/m³.

(Semi-)quantitative assessment for oral, dermal and inhalation routes

Results table exposure to PT3 restocking of animal housing							
Exposure scenario	Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated oral uptake	Estimated total uptake	Local dermal exposure (conc., %)	Local inhalation exposure (mg/m ³)
Scenario 4 – ambient temperature fogging	1/ all FA evaporated	11.5 mg/kg/d	N.A.	N.A.	11.5 mg/kg/d	N.A.	550

	2/ventilation until <6 mg/m ³ AND 3/ RMM approach	0.125 mg/kg/d	N.A.	N.A.	0.125 mg/kg/d	N.A.	<6
Scenario 4 – thermal fogging	1/ all FA evaporated	9 mg/kg/d	N.A.	N.A.	9 mg/kg/d	N.A.	430
	2/ventilation until <6 mg/m ³ AND 3/ RMM approach	0.125 mg/kg/d	N.A.	N.A.	0.125 mg/kg/d	N.A.	<6

8.4.5 Summary tables: systemic and local exposure from professional uses

Summary table: PT3 systemic exposure from professional uses			
Tier/PPE	Estimated inhalation uptake	Estimated dermal uptake	Estimated total uptake
Scenario 1, fogging			
1/none	0.013 mg/kg bw/d	16.418 mg/kg bw/d	16.431 mg/kg bw/d
2/impermeable coveralls, boots, gloves and face protection	0.013 mg/kg bw/d	0.0247 mg/kg bw/d	0.0377 mg/kg bw/d
Scenario 2, footwear disinfection, 5% FA			
1/none	0.013 mg/kg bw/d	0.199 mg/kg bw/d	0.212 mg/kg bw/d
2/coated coveralls, boots, gloves and face protection	0.013 mg/kg bw/d	0.0199 mg/kg bw/d	0.0329 mg/kg bw/d
Scenario 3, animal feet/transport vehicle disinfection, 5% FA			
1/none	0.02 mg/kg bw/d	0.0275 mg/kg bw/d	0.0475 mg/kg bw/d
2/coveralls, boots, gloves and face protection	0.02 mg/kg bw/d	0.00275 mg/kg bw/d	0.0228 mg/kg bw/d

Scenario 4, restocking of animal housing after ambient temperature fogging			
1/ all FA evaporated	11.5 mg/kg bw/d	N.A.	11.5 mg/kg bw/d
2/ventilation until <6 mg/m ³ AND 3/ RMM approach	0.125 mg/kg/d	N.A.	0.125 mg/kg/d
Scenario 4, restocking of animal housing after thermal fogging			
1/ all FA evaporated	9 mg/kg/d	N.A.	9 mg/kg/d
2/ventilation until <6 mg/m ³ AND 3/ RMM approach	0.125 mg/kg/d	N.A.	0.125 mg/kg/d

Summary table: PT3 local exposure from professional uses		
Tier/PPE	Local inhalation exposure	Local dermal exposure
Scenario 1, fogging		
1/none	3.8 mg/m ³ (vapour)	55% (M&L) 5.5-19% (disposal)
2/impermeable coveralls, boots, gloves and face protection		
Scenario 2, footwear disinfection, 5% FA		
1/none	3.8 mg/m ³ (vapour)	55% (M&L) 5% (disposal)
2/impermeable coveralls, boots, gloves and face protection	3.8 mg/m ³ (vapour)	
Scenario 3, animal feet disinfection, 5% FA		
1/none	3.5 mg/m ³ (vapour, M&L) 1.0 mg/m ³ (vapour, application) 0.38 mg/m ³ (vapour, post-application)	55% (M&L) 5% ((post)application)
2/coveralls, boots, gloves and face protection	3.5 mg/m ³ (vapour, M&L) 1.0 mg/m ³ (vapour, application) 0.38 mg/m ³ (vapour, post-application)	
Scenario 4, restocking of animal housing after ambient temperature fogging		
1/ all FA evaporated	550 mg/m ³	N.A.

2/ventilation until <6 mg/m ³ AND 3/ RMM approach	<6 mg/m ³	N.A.
Scenario 4, restocking of animal housing after thermal fogging		
1/ all FA evaporated	430 mg/m ³	N.A.
2/ventilation until <6 mg/m ³ AND 3/ RMM approach	<6 mg/m ³	N.A.

8.4.6 Combined scenarios

Fogging is assumed to be performed by professional contractors; fogging scenarios combined with footwear or animal feet disinfection, or with restocking, will not be considered here.

For farmers, scenarios 2, 3 and 4 (disinfection of footwear in dipping troughs; animal feet disinfection; restocking of animal housing after disinfection) are not unlikely to be combined within a normal working day.

For local exposure, no addition of exposure levels is performed; only the highest exposure level in air is considered relevant.

Summary table: combined systemic exposure from professional uses			
Scenarios combined	Estimated inhalation uptake	Estimated dermal uptake	Estimated total uptake
Combined restocking of animal housing and exposure from footwear and animal feet disinfection, experienced farmers			
Scenarios 2,3,4 restocking after ambient temperature fogging, footwear, animal feet disinfection Tier 1	11.533 mg/kg bw/d	0.2265 mg/kg bw/d	11.76 mg/kg bw/d
Scenarios 2,3,4 restocking after ambient temperature fogging, footwear, animal feet disinfection Tier 2	0.158 mg/kg bw/d	0.02265 mg/kg bw/d	0.181 mg/kg bw/d
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection Tier 1	9.033 mg/kg bw/d	0.2265 mg/kg bw/d	9.26 mg/kg bw/d
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection Tier 2	0.158 mg/kg bw/d	0.02265 mg/kg bw/d	0.181 mg/kg bw/d

8.5 NON-PROFESSIONAL EXPOSURE

No non-professional use of the biocidal product is foreseen.

8.6 SECONDARY EXPOSURE OF THE GENERAL PUBLIC EXCLUDING DIETARY EXPOSURE

For the representative products in this CAR, secondary or indirect exposure of the general public should be avoided by implementation of appropriate RMM, considering the volatility and corrosive properties of the a.s., and the fact that PPE and RPE are not applicable for the general public. As access of the general public to treated areas needs to be excluded by RMM, the scenario for secondary exposure is not considered further.

8.7 DIETARY EXPOSURE

Livestock can be exposed to residues of biocidal products used for disinfection of animal houses. Possible routes of exposure are inhalation of residues from air, dermal contact through rubbing of skin on treated surfaces and oral exposure through licking of treated surfaces and by consumption of feed/water from treated troughs. It could be deemed necessary to assess whether these residues can enter the food chain. Walking through hoof baths will also contribute to dermal and oral exposure; here also, livestock exposure might lead to residues in food.

Formic acid occurs naturally in animals and most plants, and is an inherent ingredient in human food. It is an intermediate in normal metabolism. Its toxicokinetic properties and its metabolism have been investigated in rat, dog, monkey, pig and humans. Formic acid and formate salts are rapidly absorbed, converted and eliminated. There is no indication for accumulation of formate. The possibility for crossing of barriers such as exposure via breastmilk is low.

Formic acid, and two of the formate salts, are approved feed additives in the EU at concentrations up to 1.2% (pigs) and 1.0% in all other species including ruminants and poultry. Approved drinking water concentration for food producing animals is 0.4%. The EFSA Panel concluded that this use would not increase the human formic acid exposure through the consumption of products obtained from the treated animals (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03). The use and risks of feed and drinking water additives containing formic acid and formate salts did not indicate the need for any further studies.

For formic acid currently default MRLs of 0.01 mg/kg apply according to Art.18(1)(b) Reg 396/2005.

Due to its rapid turnover and unlikely accumulation, an estimation of exposure of humans to formic acid residues through diet as a consequence of animal house disinfection is not considered here.

It is proposed that assessment of dietary risk for humans and livestock be undertaken at biocidal product authorisation.

For a tentative approach to the 'Guidance on the BPR V III HH-Assessment & Evaluation, Section 6: Guidance On Estimating Livestock Exposure to Active Substances used in Biocidal Products, see Appendix II.

8.7.1 Information of non-biocidal use of the active substance

Summary table of other (non-biocidal) uses			
	Sector of use ¹	Intended use	Reference value(s) ²
1.	industry	Industrial manufacture of polymers, resins	
2.	industry/professional workers	Polymer processing	
3.	industry/professional workers	(Industrial) use as processing aid	
4.	industry/professional workers	Industrial use in laboratories	
5.	industry	Use as an intermediate	
6.	industry	Uses in coatings	
7.	Industry/professional workers	Use in cleaning agents	
8.	Animal nutrition	Feed hygiene agent	Maximum proposed dose ³ : pigs: 12000 mg/kg All other animal species 10000 mg formic acid equivalents/kg complete feed

¹ e.g. plant protection products, veterinary use, food or feed additives

² e.g. MRLs. Use footnotes for references.

³ (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03)

8.7.2 Estimating transfer of biocidal active substances into foods as a result of professional and/or industrial application(s)

At the time of writing, no consolidated guidance is available on estimating dietary risk from transfer of biocidal active substances into foods for professional uses. Therefore transfer of FA into foods as a result of professional applications will not be assessed.

8.7.3 Estimating transfer of biocidal active substances into foods as a result of non-professional use

Not applicable; the active substance is not intended for non-professional use.

8.8 EXPOSURE ASSOCIATED WITH PRODUCTION, FORMULATION AND DISPOSAL OF THE BIOCIDAL PRODUCT

Please refer to section 8.3 on industrial exposure; disposal of the biocidal product is mentioned for each scenario in section 8.4 on professional exposure.

9 ENVIRONMENTAL EXPOSURE ASSESSMENT

The representative product Protectol® FM 85 is intended to be used in a wide variety of products under PT2, 3, 4, 5 and 6. As a PT3 biocide, the product is intended to be used for the disinfection of means of transport, footwear, livestock hoofs and machinery.

For the intended PT3 scenarios, Protectol® FM 85 should not be used without first being diluted from a formulated concentrate of 45-55%. The applicant proposes formic acid concentrations in the ready-for-use solution obtained via dilution of 4.5%-5.5% (= 5.3% to 6.5% of Protectol® FM 85) for ambient temperature fogging, of 19% a.s. (= 22.4% of Protectol® FM 85) for thermal fogging, and of 5% a.s. (= 5.9% of Protectol® FM 85) for dipping for the disinfection of means of transport, footwear and animal feet.

The applicant proposes to include the following scenarios in the environmental exposure assessment:

- Disinfection of animal housing by means of fogging;
- Disinfection of means of transport, footwear and animal feet through dipping.

Fogging is not fully supported by the available efficacy data (please refer to section 7 'Efficacy'). Furthermore, disinfection of animal housing is typically performed by spraying. According to the Emission Scenario Document for Product Type 3 (2011), disinfection with aerosol by nebulizer or vaporizer is only carried out in exceptional cases, since it can only be carried out in sealable small housings. Nevertheless, the scenario 'Disinfection of animal housing by means of fogging' has been retained in the environmental exposure assessment of the product Protectol® FM 85.

The proposed disinfection of means of transport through dipping does not correspond to the harmonised scenario 'Disinfection of vehicles used for animal transport' available in the Emission Scenario Document for Product Type 3 (2011), which is a scenario for spraying disinfection at slaughterhouse sites. No product-specific information for the proposed use is provided by the applicant for the product Protectol® FM 85. Therefore, the use is not retained in the environmental exposure assessment. However, it is assumed that this use is covered by the scenario for disinfection of animal's feet available in the Emission Scenario Document for PT3.

Based on the applicant's use description, the validated efficacy data and the available harmonised environmental emission scenarios, the following uses will be evaluated:

Field of use envisaged	Users	Likely concentration at which a.s. will be used
Disinfection of footwear and animal feet through dipping	Professionals	Formulated concentrate of 45% - 55% formic acid (52.9% - 64.7% of Protectol® FM 85). Concentrate is further diluted to use concentration of 5% a.s. in the ready-for-use solution (= 5.9% of Protectol® FM 85), unspec. volume.

Field of use envisaged	Users	Likely concentration at which a.s. will be used
Disinfection of animal housing by means of fogging (ambient temperature or thermal fogging)	Professionals	Formulated concentrate 45-55% a.s.; diluted to: <ul style="list-style-type: none"> 4.5-5.5 % a.s. in the ready-for-use solution (= 5.3 % to 6.5 % of Protectol® FM 85); 1.0 L/100 m³ for ambient temperature fogging 19 % a.s. in the ready-for-use solution (= 22.4% of Protectol® FM 85); 2.25 L/1000 m³ for thermal fogging

General information	
Assessed PT	PT 3
Assessed scenarios	<u>Scenario 1</u> : Disinfection of footwear <u>Scenario 2</u> : Disinfection of animal's feet <u>Scenario 3</u> : Disinfection of animal housing by means of fogging
ESD(s) used	Emission Scenario Document for Product Type 3: Veterinary hygiene biocidal products (JRC Scientific and Technical Reports, 2011) Addendum to OECD Emission Scenario Document for Insecticides for Stables and Manure Storage Systems (Agreed at the Environment Working Group on November 26, 2015) Addendum to OECD Emission Scenario Document for Insecticides for Stables and Manure Storage Systems: Addition of calculation routines to incorporate degradation in manure (Agreed at the Environment Working Group V on November 24, 2016)
Approach	<u>Scenario 1</u> : Average consumption <u>Scenario 2</u> : Average consumption <u>Scenario 3</u> : Average consumption
Distribution in the environment	Calculated based on the ECHA Guidance on the Biocidal Products Regulation, Volume IV Environment – Assessment and Evaluation (Parts B + C)
Groundwater simulation	FOCUS PEARL v4.4.4. (see §13.7 'Aggregated exposure')
Confidential Annexes	No
Life cycle steps assessed	<u>Scenarios 1 to 3</u> :

	Production: No Formulation No Use: Yes Service life: No
Remarks	/

Biocidal product specific data

The applicant provided two addenda to the biocidal active substance registration dossier aiming at assessing the fate of formic acid in soil and manure in order to refine the exposure calculations. The addenda ('Formic acid: Fate and degradability – Soil and Manure' (August 20, 2019) and 'Formic acid: Degradability in Manure' (September 07, 2020)) give an overview of the data found in the public literature on degradability and fate of formic acid in soil and manure.

In addition to the mentioned addendum, also Doc IIIA robust study summaries of open literature data were submitted for the degradability and fate of formic acid in soil and manure. Reference is made to sections 4.1.1.3.5 and 4.1.1.3.6 of Part A of the present CAR.

The addenda and the evaluation by the eCA are included in Doc IIIB 10.2.

Following ENV WG-I-2022, a DT₅₀ value for soil of 1 day (12 °C; please refer to section 4.1.1.3.6) and a DT₅₀ value of ≤ 10.5 days (20 °C; please refer to section 4.1.1.3.5) are agreed. At the time of writing (April 2022), no agreed environmental relevant temperature exists for the manure. For this specific case, from a precautionary principle, it was agreed at ENV WG-I-2022 to reconvert the DT₅₀ value for manure to a temperature of 12 °C as a first tier.

9.1 EMISSION ESTIMATION

9.1.1 Scenario 1: Disinfection of footwear

Only the values which have been included as "Set values" in the emission scenario (ESD PT3 §2.4.1.4), default values which are under discussion or when it is possible to choose between different default values are stated in the table below. Detailed calculation sheets are provided in Appendix III.

Taking into account TABv2 entry ENV60, only nitrogen immission standards are considered.

For degradation in the manure storage tank, a DT50-value of 10.5 days is assumed. The amount of active ingredient in manure/slurry after the relevant number of biocide applications for the manure application to grassland (Qai_{grass}) is calculated as follow:

$$Qai_{grass} = \sum_{t=1}^{Tgr-int} Qai_{manure} * e^{-kdeg_{manure}*t}$$

with:

- t: number of days in the manure/slurry storage [d];
- Tgr-int: land application interval for grassland (=manure storage time) [d];
- Qai_{manure} : amount of active ingredient in manure/slurry after one application [kg];
- $kdeg_{manure}$: rate constant for degradation in manure [d^{-1}].

Similarly, the amount of active ingredient in manure/slurry after the relevant number of biocide applications for the manure application to arable land (Qai_{arab}) is calculated as follow:

$$Qai_{arab} = \sum_{t=1}^{Tar-int} Qai_{manure} * e^{-kdeg_{manure}*t}$$

with:

- t: number of days in the manure/slurry storage [d];
- Tar-int: land application interval for arable land (=manure storage time) [d];
- Qai_{manure} : amount of active ingredient in manure/slurry after one application [kg];
- $kdeg_{manure}$: rate constant for degradation in manure [d^{-1}].

For Tgr-int and Tar-int, the default values of respectively 53 days and 212 days are used.

Input parameters for calculating the local emission			
Input	Value	Unit	Remarks
Scenario 1: Disinfection of footwear			
Content of active ingredient in formulation	50	g/L	

Dilution factor (for preparation of the working solution from the formulation (product))	1	-	
Half-life for biodegradation in soil	1	d	at 12°C
Half-life for biodegradation in manure	19.9	d	at 12 °C

Emissions are calculated according to the ESD PT3 §2.4.1.4. The remaining solution in the footbath is either discharged to the waste water or to the manure (both scenarios are considered). Emission to air is considered negligible, taking into account the low surface area of the tub and that the solution is only stirred up a few times per day.

Resulting local emission to relevant environmental compartments			
	Soil (one year) [mg/kg wwt]		STP [kg/d]
	Grassland, degradation PIECgrs4-N_degr	Arable land PIECars-N	
1 Dairy cows	3.307E-01	9.811E-02	5.000E-01
2 Beef cattle	3.111E-01	9.230E-02	5.000E-01
3 Veal calves	5.881E+00	1.745E+00	5.000E-01
4 Sows, in individual pens	1.195E+00	3.545E-01	5.000E-01
5 Sows in groups	1.195E+00	3.545E-01	5.000E-01
6 Fattening pigs	9.207E-01	2.732E-01	5.000E-01
7 Laying hens in battery cages without treatment	2.642E-01	7.838E-02	5.000E-01
8 Laying hens in battery cages with aeration (belt drying)	2.948E-01	8.747E-02	5.000E-01
9 Laying hens in battery cages with forced drying (deep pit, high-rise)	2.948E-01	8.747E-02	5.000E-01
10 Laying hens in compact battery cages	2.948E-01	8.747E-02	5.000E-01
11 Laying hens in free range with litter floor (partly litter floor, partly slatted)	6.554E-01	1.944E-01	5.000E-01
12 Broilers in free range with litter floor	3.592E-01	1.066E-01	5.000E-01

13 Laying hens in free range with grating floor (aviary system)	3.277E-01	9.722E-02	5.000E-01
14 Parent broilers in free range with grating floor	5.373E-01	1.594E-01	5.000E-01
15 Parent broilers in rearing with grating floor	9.089E-01	2.697E-01	5.000E-01
16 Turkeys in free range with litter floor	2.325E-01	6.898E-02	5.000E-01
17 Ducks in free range with litter floor	4.090E-01	1.213E-01	5.000E-01
18 Geese in free range with litter floor	2.325E-01	6.898E-02	5.000E-01

9.1.2 Scenario 2: Disinfection of animal's feet

Only the values which have been included as "Set values" in the emission scenario (ESD PT3 §2.4.2.4), default values which are under discussion or when it is possible to choose between different default values are stated in the table below. Detailed calculation sheets are provided in Appendix III.

Taking into account TABv2 entry ENV60, only nitrogen immission standards are considered.

For degradation in the manure storage tank, a DT50-value of 10.5 days is assumed. The biocidal product is applied twice a week. For the calculations, it is supposed that the manure storage time for grassland (53 days) corresponds to 8 weeks. The amount of active ingredient in manure/slurry after the relevant number of biocide applications for the manure application to grassland ($Q_{ai_{grass}}$) is calculated as follow:

$$Q_{ai_{grass}} = \sum_{t=1}^{8 \text{ weeks}} 2 * Q_{ai_{manure}} * e^{-k_{deg_{manure}} * t * 7}$$

with:

- t: number of weeks in the manure/slurry storage [week];
- $Q_{ai_{manure}}$: amount of active ingredient in manure/slurry after one application [kg];
- $k_{deg_{manure}}$: rate constant for degradation in manure [d^{-1}].

Similarly, the amount of active ingredient in manure/slurry after the relevant number of biocide applications for the manure application to arable land ($Q_{ai_{arab}}$) is calculated as follow (assuming a manure storage time of 30 weeks (212 days)):

$$Q_{ai_{arab}} = \sum_{t=1 \text{ week}}^{30 \text{ weeks}} 2 * Q_{ai_{manure}} * e^{-k_{deg_{manure}} * t * 7}$$

with:

- t: number of weeks in the manure/slurry storage [week];
- $Q_{ai_{manure}}$: amount of active ingredient in manure/slurry after one application [kg];
- $k_{deg_{manure}}$: rate constant for degradation in manure [d^{-1}].

Input parameters for calculating the local emission			
Input	Value	Unit	Remarks
Scenario 2: Disinfection of animal's feet			
Content of active ingredient in formulation	50	g/L	
Dilution factor (for preparation of the working solution from the formulation (product))	1	-	
Half-life for biodegradation in soil	1	d	at 12°C
Half-life for biodegradation in manure	19.9	d	at 12 °C

Emissions are calculated according to the ESD PT3 §2.4.2.4. The remaining solution in the footbath is either discharged to the waste water or to the manure (both scenarios are considered). Emissions to air are considered negligible; the air is not an environmental compartment of concern.

Resulting local emission to relevant environmental compartments			
	Soil (one year) [mg/kg wwt]		STP [kg/d]
	Grassland, degradation PIECgrs4-N_degr	Arable land PIECars-N	
Dairy cows	5.254E+00	1.530E+00	6.075E+01

9.1.3 Scenario 3: Disinfection of animal housing by means of fogging

No harmonised scenario for disinfection of animal housing by means of fogging is available. The ESD for PT3 mentions (§2.1.1): *Disinfection with aerosol (droplet size < 10 µm) by nebulizer or vaporizer is only carried out in exceptional cases, since it can only be carried out in sealable small housings (Bodenschatz, 2006).*

For the present emission estimation of the product Protectol® FM 85, a veal calves stable type (with the same defaults as for the spraying scenario) is considered and this for two reasons⁹:

1. in spraying scenarios the veal calves stable type generally results in the worst-case emissions compared to the other stable types;
2. the veal calves stable has the smallest volume and thus can be considered as most representative for a fogging scenarios since this can only be carried in sealable small housings.

The emission scenario is based on the harmonised spraying scenario available in the ESD for PT3. Only new values, values which have been included as "Set values" in the emission scenario (ESD PT3 §2.1.4.1), default values which are under discussion or when it is possible to choose between different default values are stated in the table below. Detailed calculation sheets are provided in Appendix III.

Regarding degradation in manure, the recommendations agreed at the Environment Working Group V on November 24, 2016 are followed. For grassland, we are in the situation where $T_{bioc-int} \geq T_{gr-int}$ and $N_{app-manure_{gr}} = 1$, and therefore the 'average quantity' approach applies. For arable land, the calculation routine where $T_{ar-int} = T_{bioc-int}$ is followed, and thus the 'average quantity' approach also applies for this case. The calculation sheets provided with the addendum are used for the degradation calculations in the manure and are provided in Appendix III.

For ambient temperature fogging, 1.0 L of a diluted solution containing 5.5% active ingredient (or 55 g a.i./L) is used for a treated volume of 100 m³, which corresponds to 0.55 g a.i./m³. For thermal fogging, 2.25 L of a diluted solution containing 19% active ingredient (or 190 g a.i./L) is used for a treated volume of 1000 m³, which corresponds to 0.43 g a.i./m³. Application frequency and interval are the same for both methods, and therefore ambient temperature fogging is considered worst-case compared to thermal fogging.

The volume of a veal calves stable (590 m³) is taken from Table 5.2 of the ESD for PT18, Stables and Manure Storage Systems.

During the peer-review phase, one MS commented that there are several animal categories (i.e. 8, 11, 12, 16, 17, 18) where the housings can have a connection to the sewer system. It was agreed bilaterally to include emissions to the STP for turkeys (animal category 16), which can

⁹ Please note that Protectol® FM 85 is ██████████ and thus by definition has no real use case. Application of a real product in other situations or stable types will have to be dealt with at product authorisation stage based on product specific information.

be considered as the worst-case for this emission route. Only the emissions to the STP are considered for this stable type. According to the ESD for PT3, the fraction of active ingredient released to the STP is 0.2 for turkey stables.

Input parameters for calculating the local emission – manure route			
Input	Value	Unit	Remarks
Scenario 3: Disinfection of animal housing by means of fogging			
Type of housing/manure storage	Veal calves	-	worst-case
Type of application	Ambient temperature fogging	-	worst-case
Content of active ingredient in formulation	55	g/L	
Amount of (undiluted) product prescribed to be used per m ³	0.01	L/m ³	Application rate of 1.0 L/100 m ³
Dilution factor (for preparation of the working solution from the formulation (product))	1	-	
Stable volume	590	m ³	ESD PT18
Half-life for biodegradation in soil	1	d	at 12°C
Half-life for biodegradation in manure	19.9	d	at 12 °C

Emissions are calculated according to the ESD PT3 §2.1.4 and the addendum on the addition of calculation routines to incorporate degradation in manure. No emissions take place to the STP for this stable type. Emissions to air are considered negligible; the air is not an environmental compartment of concern.

Input parameters for calculating the local emission – STP route			
Input	Value	Unit	Remarks
Scenario 3: Disinfection of animal housing by means of fogging			
Type of housing/manure storage	Turkeys	-	worst-case

Type of application	Ambient temperature fogging	-	worst-case
Content of active ingredient in formulation	55	g/L	
Amount of (undiluted) product prescribed to be used per m ³	0.01	L/m ³	Application rate of 1.0 L/100 m ³
Dilution factor (for preparation of the working solution from the formulation (product))	1	-	
Stable volume	12500	m ³	ESD PT18
Fraction of a.i. released to waste water	0.2	-	ESD PT3
Half-life for biodegradation in soil	1	d	at 12°C
Half-life for biodegradation in manure	19.9	d	at 12°C (degradation in manure not applicable for this emission route)

Resulting local emission to relevant environmental compartments			
	Soil (one year) [mg/kg wwt]		STP [kg/d]
	Grassland, degradation PIECgrs4-N_degr	Arable land PIECars-N	
Veal calves	4.654E-02	2.451E-02	N/A
Turkeys	N/A	N/A	1.375

9.2 FATE AND DISTRIBUTION IN EXPOSED ENVIRONMENTAL COMPARTMENTS

Identification of relevant receiving compartments based on the exposure pathway									
	Fresh-water	Sediment	Sea-water	Seawater sediment	STP	Air	Soil	Ground-water	Other
Scenario 1 (footwear)	+	(-)	(+)	(-)	++	(-)	+	+	(-)
Scenario 2 (animal's feet)	+	(-)	(+)	(-)	++	(++)	+	+	(-)
Scenario 3 (animal housing, fogging)	+	(-)	(+)	(-)	++	(+)	+	+	(-)

++ *Compartment directly exposed*

- *Compartment not exposed*

+ *Compartment indirectly exposed*

() *Compartment potentially exposed [but unlikely to be a significant concern due to hazard data and / or scale of exposure]*

Input parameters (only set values) for calculating the fate and distribution in the environment			
Input	Value	Unit	Remarks
Molecular weight	46.03	g/mol	
Melting point	8	°C	
Boiling point	100.23	°C	
Vapour pressure (at 12 °C)	2400	Pa	
Water solubility (at 12 °C)	1.09x10 ⁶	mg/l	
Log10 Octanol/water partition coefficient	-2.10	---	(pH 7)

Organic carbon/water partition coefficient (Koc)	30	l/kg	(pH 7)
Henry's Law Constant (at 12 °C)	0.101	Pa/m ³ /mol	
Acid dissociation constant	3.7	---	Predominant species at a pH of 7 is formate, which is reflected in the pH dependent Koc.
Biodegradability	Ready biodegradable		
DT ₅₀ for degradation in soil (12 °C)	1	day	
DT ₅₀ for degradation in manure (12 °C)	19.9	days	

Fate and distribution in the sewage treatment plant (STP) is calculated using SimpleTreat 4.0. In accordance to the TAB v2 (ENV9), the concentration of suspended solids (C_{ss}) in the effluent is changed manually to 30 mg/L (0.030 kg/m³).

Calculated fate and distribution in the STP		
Compartment	Percentage [%]	Remarks
	All scenarios	
Air	0.04222	Calculated with SimpleTreat 4.0 ¹⁰
Water	7.991	
Sludge	0.27946	
Degraded in STP	91.69	

Notes to take into account when performing fate and distribution calculations at product authorisation stage:

¹⁰ In accordance with TAB entry ENV 9, the concentration of suspended solids (C_{ss}) in the effluent is changed manually to 30 mg/L (0.03 kg/m³).

-
- For the calculations of the PECsoil, an 'overall removal rate constant' should be considered, taking degradation in soil, leaching and volatilisation into account (Guidance on the BPR Vol IV, Part B+C, Equation 56 (v. 10/19)). In the CAR, only the rate constant for degradation in soil (k_{deg_soil}) is used (which result in worst-case PECsoil values).
 - The calculation for PEC surface water (via run-off) should also consider the sorption onto suspended matter, see TAB (v.2021) ENV 11. The PEC values presented in the CAR represent the worst-case.
 - In the calculation of the P(I)ECsoil_10years, $T_{gr-int_no_manure}$ should be 365d (instead of 206d) according to the revision of AHEE Recom WG V 2015 discussed at WG ENV I 2018. However, this change will not have a significant effect on the PEC values for a rapidly degrading substance as formic acid.
 - The calculation routines for degradation in manure should be performed following the 'quantity average' approach as presented in Addendum 'Addition of calculation routines to incorporate degradation in manure', section 4.3.

9.3 CALCULATED PEC VALUES

For scenarios 1 and 2, emissions to the STP and to the manure are considered independently. The maximum PEC values per compartment are presented in the table below. All presented PEC values for scenarios 1 result from the manure route, except the PEC values for the STP. For scenario 2, the PEC values for the STP and surface water result from the STP route, and the PEC values for soil and groundwater from the manure route. Taking into account TABv2 entry ENV60, only nitrogen immission standards are considered. Please refer to Appendix III for detailed results.

Summary table on calculated PEC values								
	PEC _{STP}	PEC _{water}	PEC _{sed} [*]	PEC _{seawater}	PEC _{seased}	PEC _{soil,twa} ^{**}	PEC _{GW} ^{***}	PEC _{air}
	[mg/L]	[mg/L]	[mg/kg _{dwt}]	[mg/L]	[mg/kg _{wwt}]	[mg/kg _{wwt}]	[µg/L]	[mg/m ³]
Scenario 1 (footwear)	2.00E-02	7.285E-03	see PEC _{water} [*]	n/a	n/a	2.828E-01	7.285E+01	n/a
Scenario 2 (animal's feet)	2.43E+00	2.43E-01	see PEC _{water} [*]	n/a	n/a	2.527E-01	6.508E+01	n/a
Scenario 3 (animal housing, fogging)	5.49E-02	5.49E-03	see PEC _{water} [*]	n/a	n/a	2.238E-03	2.757	n/a

* Since the PNEC sediment was calculated according to the equilibrium partitioning method, the risk assessment for freshwater covers that for the sediment.

** The PNEC_{soil} is derived by equilibrium partitioning from a PNEC_{aquatic} for chronic exposure which justifies the use the time weighted average PEC_{soil}. The PEC_{soil} presented in this table are the PEC_{grs_10_degr-N_twa}.

*** The values for PEC_{GW} presented in this table are the TIER1 porewater concentrations.

The calculated porewater concentrations (PEC_{GW}) exceed the threshold of 0.1 µg/L. Further refinement using FOCUS PEARL to model more realistic groundwater concentrations instead of porewater concentrations is presented in section 13.7 of this CAR (Aggregated exposure).

9.4 PRIMARY AND SECONDARY POISONING

9.4.1 Primary poisoning

Not relevant

9.4.2 Secondary poisoning

Formic acid is not expected to bioaccumulate based on the experimentally derived log Kow of -2.1 (23 °C, pH7) and the calculated BCF (see §4.1.3 above). Therefore, secondary poisoning of formic acid in either the aquatic or terrestrial food chain is considered not relevant.

10 ASSESSMENT OF EFFECTS ON HUMAN HEALTH FOR THE PRODUCT

10.1 PRODUCT(S)

The toxicological properties of the product may be derived from the properties of the active substance and other components of the product. Information on the toxicity of the active substance is presented in Part A, Section 3. There are no compounds of concern in the formulated product that adversely affect the conclusions of the risk assessment for the active substance in the product, therefore limited further assessment is needed.

10.2 DERMAL ABSORPTION

Since the biocidal product **Protectol® FM 85**, containing 85% formic acid with [REDACTED], and formic acid itself are classified as corrosive it is expected that the irritation potential would be sufficient to prevent use of the solution without taking precautions to prevent dermal exposure and so minimising the potential for absorption.

Furthermore, the corrosive nature of formic acid would also corrode the skin sample used in the test, thereby producing meaningless absorption results.

Severe metabolic acidosis resulting from dermal contact with formic acid from biocidal products as described in several case reports (see section 3.3.1 and 3.14), demonstrated rapid dermal absorption through the acid-burned skin.

Therefore, a dermal absorption study using the biocidal product **Protectol® FM 85** is scientifically unjustified.

Value(s) used in the Risk Assessment – Dermal absorption	
Value(s)*	In a first tier of risk assessment, a worst-case value for dermal absorption of 100% is used for external dermal exposure.
Justification for the selected value(s)	Severe metabolic acidosis resulting from dermal contact with formic acid from biocidal products as described in several case reports, demonstrated rapid dermal absorption through the acid-burned skin. Due to the corrosive properties the dermal absorption of formic acid was not tested. Dermal absorption is known to occur from incidental exposure to large quantities of concentrated formic acid which led to systemic toxicity (section 3.3.1 and 3.14).

Data waiving

Information requirement	Dermal absorption of the biocidal product Protectol® FM 85 containing 85% formic acid has not been investigated.
Justification	Due to the corrosive properties of the biocidal product and formic acid, no dermal absorption study is requested.

10.3 ACUTE TOXICITY

The acute toxic action profile of formic acid is predominantly determined by its inherent irritating/corrosive properties. The toxicity values after oral uptake and inhalation in rats suggest formic acid to be acutely harmful. The clinical signs give no evidence of specific systemic adverse effects.

The biocidal product, **Protectol® FM 85**, contains the active substance to 85% with [REDACTED]. The marketed products may be further diluted by downstream users to give less concentrated solutions, which are further diluted by the end users with water during the application or to give the ready-for-use solutions. The intended concentrations of the market products and the concentrations during the use as PT3 products are summarized as follows:

Product Type	Formic acid concentration [%]		Remarks
	Market product	Ready-for-use solution	
3	45-55%	4.5-5.55	Professional, Disinfection of animal housings, ambient temperature fogging
3	45-55%	19%	Professional, Disinfection of animal housings, thermal fogging
2	45-55%	5%	Professional, Disinfection of means of transport, footwear and animal feet through dipping

It is evident from the above that 55% formic acid concentrates must be considered as the worst case.

Acute effects are likely to be caused by formic acid as the major component of the product. The acute oral and inhalation toxicity of formic acid has been characterised as described in section 3.2 and is applicable to that of the biocidal product.

10.3.1 Overall conclusion on acute toxicity

Value used in the Risk Assessment – Acute toxicity

Value(s)	LD ₅₀ oral 730 mg/kg bw ¹¹ LC ₅₀ inhalation 7.4 mg/l
Justification for the selected value	Appropriate studies are available for determining the LD ₅₀ oral and LC ₅₀ inhalation of formic acid. See sections 3.2.1 and 3.2.3.
Classification for the product according to CLP and DSD	Acute toxicity, oral, cat. 4, H302 Acute toxicity, inhalation, cat. 3, H331 <i>Corrosive properties determine the toxicity of formic acid; additional labelling EUH071</i>

Data waiving	
Information requirement	Acute toxicity of Protectol® FM 85
Justification	Since both formic acid and the biocidal product are classified as corrosive, additional acute toxicity testing with the biocidal product is scientifically unjustified and is not in the interests of animal welfare.

10.4 CORROSION AND IRRITATION

No skin and eye irritation study reports on formic acid and the biocidal product, **Protectol® FM 85**, are available.

Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive (C, R 35) in the EU (12th ATP to Directive 67/548/EEC).

According to Directive EU CLP 1272/2008, Formic Acid is to be classified as skin corrosive 1A and with the following concentration limits:

Skin Corr. 1B; H314: 10% ≤ C < 90%

Skin Corr. 1A; H314: C ≥ 90%

Skin Irrit. 2; H315: 2% ≤ C < 10%

¹¹ Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

Eye Irrit. 2; H319: 2% ≤ C < 10%

In addition, the corrosive potential of formic acid and formulations containing formic acid has been reported on several occasions after accidental dermal exposure in humans and documented in case reports. For a more comprehensive discussion see section 3.3.1.

We propose additional labelling with EUH071, 'corrosive to the respiratory tract'. See section 3.3.3 for further details. This classification is transferred to **Protectol® FM 85**.

10.4.1 Skin corrosion and irritation

No data on the biocidal product are available.

10.4.2 Serious eye damage and eye irritation

No data on the biocidal product are available.

10.4.3 Respiratory tract irritation

No data on the biocidal product are available.

10.4.4 Overall conclusion on corrosion and irritation

Conclusion used in the Risk Assessment – Corrosion and irritation	
Value(s) or Conclusion(s)	Formic acid and Protectol® FM 85 are corrosive to skin Formic acid and Protectol® FM 85 are corrosive to the respiratory tract
Justification for the selected value/ conclusion	See justification below

Classification of the product according to CLP and DSD	Skin Corr. 1B; H314 EUH071
---	-------------------------------

Data waiving													
Information requirement	No skin and eye irritation study reports on formic acid and the biocidal product, <i>Protectol® FM 85</i> , are available.												
Justification	<p>Due to the inherent properties of formic acid (strong acid), the substance has been classified as corrosive (C, R 35) in the EU (12th ATP to Directive 67/548/EEC) with the following concentration limits:</p> <table> <tr> <td>C ≥ 90 %</td> <td>C, R35</td> <td>corresponds to</td> <td>Skin Corr. 1A; H314</td> </tr> <tr> <td>10 % ≤ C < 90 %</td> <td>C, R34</td> <td></td> <td>Skin Corr. 1B; H314</td> </tr> <tr> <td>2 % ≤ C < 10 %</td> <td>Xi, R36/38</td> <td></td> <td>Skin Irrit. 2; H315: 2% ≤ C < 10% Eye Irrit. 2; H319: 2% ≤ C < 10%</td> </tr> </table> <p>EUH071: the corrosive properties determine the toxicity of formic acid (CLP Regulation Annex II, point 1.2.6).</p>	C ≥ 90 %	C, R35	corresponds to	Skin Corr. 1A; H314	10 % ≤ C < 90 %	C, R34		Skin Corr. 1B; H314	2 % ≤ C < 10 %	Xi, R36/38		Skin Irrit. 2; H315: 2% ≤ C < 10% Eye Irrit. 2; H319: 2% ≤ C < 10%
C ≥ 90 %	C, R35	corresponds to	Skin Corr. 1A; H314										
10 % ≤ C < 90 %	C, R34		Skin Corr. 1B; H314										
2 % ≤ C < 10 %	Xi, R36/38		Skin Irrit. 2; H315: 2% ≤ C < 10% Eye Irrit. 2; H319: 2% ≤ C < 10%										

10.5 SENSITISATION

10.5.1 Skin sensitisation

There was no evidence of a sensitising potential for formic acid (technical, purity 85.3%) in guinea pigs using the method of Buehler according to the OECD test guideline 406 (see section 3.3.3). In addition, there is no data available (human data including market surveillance data, animal data, open literature) which may be indicative of the potential of formic acid to cause skin sensitisation and sensitisation by inhalation in humans.

The biocidal product, **Protectol® FM 85**, is comprised of 85% formic acid and water as only other ingredient. Skin sensitisation of the biocidal product would therefore likely to be caused by formic acid.

The biocidal product is not expected to be a sensitiser. Therefore, the request for a skin sensitisation study with the product would be scientifically unjustified and not in the interests of animal welfare.

Conclusion used in Risk Assessment – Skin sensitisation	
Value/conclusion	Formic acid and Protectol® FM 85 do not fulfill the criteria of the CLP regulation to be classified as a skin sensitiser
Justification for the value/conclusion	Skin sensitization (Buehler test) by formic acid (85.3%) has been assessed in an OECD 406 study (Buehler test). The results do not trigger a classification as skin sensitizer.
Classification of the product according to CLP and DSD	none

10.5.2 Respiratory sensitisation

No data on the biocidal product are available.

Conclusion used in the Risk Assessment – Respiratory sensitisation	
Value/conclusion	There is no indication that formic acid or Protectol® FM 85 would be respiratory sensitizers.

Justification for the value/conclusion	No data are available (human data e.g. market surveillance data, animal data, open literature) which may be indicative of the potential of formic acid to cause sensitisation by inhalation in humans. No respiratory sensitisation was seen with formic acid in two subchronic rat and mouse inhalation studies (see section 3.6.3, Thompson 1992). Hence, there is no indication that formic acid would be a respiratory sensitizer.
Classification of the product according to CLP and DSD	none

10.5.3 Overall conclusion on sensitisation

Conclusion used in the Risk Assessment – Sensitisation	
Conclusion(s)	Formic acid and Protectol® FM 85 are not skin sensitizers. There is no indication that formic acid or Protectol® FM 85 would be respiratory sensitizers.
Justification for the conclusion(s)	Classification as a sensitizer is not triggered by appropriate tests. Studies in guinea pigs (method of Buehler) showed that there is no evidence that formic acid has a potential to induce skin sensitisation. In addition, there are no data available (human data including market surveillance, animal studies, open literature) that may be indicative of the potential of formic acid to cause skin sensitisation and sensitisation by inhalation in humans.
Classification of the product according to CLP and DSD	none

Data waiving	
Information requirement	Skin sensitisation study on Protectol® FM 85
Justification	The biocidal product is not expected to be a sensitiser. Therefore, the request for a skin sensitisation study with the product would be scientifically unjustified and not in the interests of animal welfare.

10.6 OTHER

As far as known, there are no further inherent properties of the active substance and non-active substances (water) the classification of which has to be adopted to the biocidal product according to Regulation 1272/2008/EC.

11 ENVIRONMENTAL EFFECTS ASSESSMENT FOR THE PRODUCT

The ecotoxicological properties of the product may be derived from the properties of the active substance and other components of the product. Information on the ecotoxicity of the active substance is presented in Part A, Section 4.2. There are no compounds of concern in the formulated products that adversely affect the conclusions of the risk assessment for the active substance in the product , therefore no further assessment is needed.

11.1 ATMOSPHERE

No studies submitted.

11.2 STP

No studies submitted.

11.3 AQUATIC COMPARTMENT

No studies submitted.

11.4 TERRESTRIAL COMPARTMENT

No studies submitted.

11.5 PRIMARY AND SECONDARY POISONING

No studies submitted.

PART C : RISK CHARACTERISATION OF THE BIOCIDAL PRODUCT(S)

12 RISK CHARACTERISATION FOR HUMAN HEALTH

12.1 CRITICAL ENDPOINTS

The primary endpoint for formic acid is its corrosiveness. Formic acid is severely irritating and corrosive to the eyes, skin, and mucous membranes (gastrointestinal and respiratory tract) and may cause permanent damage. Due to the corrosivity of formic acid, local effects must be expected at all dose levels. Corrosive intoxication might mediate systemic injury as metabolic acidosis, intravascular hemolysis, and renal failure. Systemic adverse effects such as decrease in body weight gain (rat, mice), might be due to the inherent irritating potential. Formic acid is associated with optical nerve and photoreceptor toxicity which is observed in humans and monkeys following methanol intoxication.

Systemic toxicity of formic acid can be established by its salts, sodium formate and potassium diformate, and a closely related substance methanol, as these chemicals have a common breakdown product *in vivo*. Please see section 3.1 on Toxicokinetics for a justification of the read-across applied.

Reference values will be derived for formate and expressed as mg formate/kg bw/d. A conversion is not needed as the difference between formic acid and formate is limited to 1 H⁺ (MW of formate is 1 less than formic acid).

12.1.1 Systemic effects

Species	Route	Study duration	Test substance	Dose setting (mg/kg bw/d)	Critical effect	LO(A)EL and NO(A)EL (mg/kg bw/d)	References
Rat	Oral	Acute	Formic acid	501, 631, 794, 1000 gavage	Clinical signs and organ lesions indicated corrosive properties of the test substance - Local effect	LD50 = 730 mg/kg bw ¹²	BPD ID A6.1.1_01 FA_BPR_Ann_II 8 7 1 01 [REDACTED] 1985

¹² Final LD₅₀ will be set by RAC; it is the LD₅₀ value from the adopted RAC opinion that will need to be used in biocidal product authorisation.

					on of the gastro-intestinal tract		
N.R.	Derma l	Acute	Formic acid	-	-	No data, corrosivity	
Rat	Inhala tion	Acute	Formic acid	2.82, 6.60, 8.08, 10.6, 14.7 mg/L vapour	Clinical signs indicated corrosive properties of the test substance, evidenced by the occurrence of corneal opacity and corrosion of the dorsal nose - Local effect on the respiratory tract	LC50 = 7.4 mg/L	BPD ID A6.1.3_01 FA_BPR_Ann II 8 7 2 01 [REDACTED] 1980
Rat	Oral	Teratogen icity study	Sodium format e	0, 40, 160, 640 mg formate /kg bw/d	Systemic: no maternal systemic toxicity reached No evidence of teratogene tic or embryotoxi c effects	as formate: LOAELsyst emic > = 640 NOAELsyst emic = 640 (highest concentrat ion tested)	BPD ID A6.8.1_01 FA_BPR_Ann_II_8_1 0 3 01 [REDACTED] 2005
Rat	Oral	Subchroni c 90 day feeding study	Potassi um diforma te	0, 420, 840, 2100 mg formate /kg bw/d	Systemic: reduced bw gain Local: gastric irritation, hyperplasti c changes in the stomach	as formate: LOAELsyst emic = 2100 (highest concentrat ion tested) [REDACTED] LOAELlocal = 420 NOAELloca l < 420	BPD ID A6.4.1_01 FA_BPR_Ann_II_8_9 2_01 [REDACTED] 1998

Rat	Oral	Chronic 2-year feeding study	Potassium diformate	0, 35, 280, 1400 mg formate /kg bw/d	Systemic: reduced bw gain Local: gastric irritation, hyperplastic changes in the stomach and gastrointestinal tract	as formate: LOAEL _{systemic} = 1400 (highest concentration tested) LOAEL _{local} = 280 NOAEL _{local} = 35	BPD ID A6.5_01 FA_BPR_Ann_II_8_93_01 [REDACTED] 2002a
Rat	Oral	2-generation study	Sodium formate	0, 68, 203, 677 mg formate /kg bw/d	Systemic: decreased food consumption, decreased bw gain in F1 parental males No findings on reproduction and development	as formate: LOAEL _{systemic} = 670 (highest concentration tested) NOAEL _{systemic} = 200	BPD ID A6.8.2_01 [REDACTED] 2008b FA_BPR_Ann_II_8_10_2_01
Rat	Inhalation	Subchronic 90-day inhalation study	Formic acid	0, 15, 30, 61, 122, 244 mg/m ³ Vapour, whole body	Systemic: no evidence of systemic toxicity Local: nasal irritation, histopathological changes in nasal region	LOAEL _{systemic} > 244 mg/m ³ NOAEL _{systemic} = 244 mg/m ³ (highest dose tested) LOAEL _{local} = 61 mg/m ³ NOAEL _{local} = 30 mg/m ³	BPD ID A6.4.3_01 FA_BPR_Ann_II_8_92_03 Thompson, 1992
Mouse	Oral	Carcinogenicity study: 80-week feeding study	Potassium diformate	0, 35, 280, 1400 mg formate	Systemic: reduced bw gain	as formate: LOAEL _{systemic} = 1400	BPD ID A6.7_02. FA_BPR_Ann_II_8_11_2_01 [REDACTED] 2002b

				/kg bw/d	Local: gastric irritation, hyperplastic changes in the forestomach	(highest concentration tested) NOAEL _{systemic} = 280 LOAEL _{local} = 1400 (highest concentration tested) NOAEL _{local} = 280	
Mouse	Inhalation	Subchronic 90-day inhalation study	Formic acid	0, 15, 30, 61, 122, 244 mg/m ³	Systemic: decreased bw gain Local: nasal irritation, histopathological changes in nasal region	LOAEL _{systemic} = 244 mg/m ³ (highest dose tested) NOAEL _{systemic} = 122 mg/m ³ LOAEL _{local} = 122 mg/m ³ NOAEL _{local} = 61 mg/m ³	BPD ID A6.4.3_01 FA_BPR_Ann_II_8_9 _2_04 Thompson, 1992
Rabbit	Oral	Teratogenicity study	Sodium formate	0, 68, 203, 677 mg formate /kg bw/d	Systemic: no maternal systemic toxicity reached No evidence of terato- genetic or embryotoxic effects	as formate: as formate: LOAEL _{systemic} >= 670 NOAEL _{systemic} = 670 (highest concentration tested)	BPD ID A6.8.1_02 [REDACTED] 2008 FA_BPR_AnnII_8_10 _1_01
Pig	Oral	Subchronic 140-day feed study	Potassium diformate	0, 149, 359, 760 mg formate /(kg bw/d	No signs of maternal systemic toxicity or toxicity to reproduction or	as formate: LOAEL _{systemic} > 760	BPD ID A6.4_02 FA_BPR_Ann_II_8_9 _2_02 [REDACTED] 2004

					development at any dose level. Local: gastric effects - forestomach gastritis and erosion/ulcer	NOAELsystemic = 760 (highest concentration tested) LOAELlocal = 149 NOAELlocal < 149	
Pig	Oral	Subchronic > 300-day feed study	Potassium diformate	0, 98, 301 mg formate /kg bw/d	No signs of maternal systemic toxicity or toxicity to reproduction or development at any dose level.	as formate: LOAELsystemic, local > 300 NOAELsystemic, local = 300 (highest concentration tested)	BPD ID A6.5_02 FA_BPR_Ann_II_8_9_4_0_JNS [REDACTED] 2003

12.1.2 Local effects

Route	Effect	Study	Classification	Hazard category ¹
Dermal	corrosive	n.a.	Skin corr 1A	Very high
Respiratory	corrosive	[REDACTED] (1980) BPD ID A6.1.3_01 FA_BPR_Ann_II_8_7_2_01	EUH071	
		BPD ID A6.4.3_01 FA_BPR_Ann_II_8_9_2_03 Thompson, 1992 (see 12.1.1)		
		BPD ID A6.4.3_01 FA_BPR_Ann_II_8_9_2_04 Thompson, 1992 (see 12.1.1)		
oral	irritating to the gastrointestinal tract (mouth, oesophagus, forestomach)	[REDACTED] (1998), BPD ID A6.4.1_01; FA_BPR_Ann_II_8_9_2_01 [REDACTED] (2002a). BPD ID A6.5_01; FA_BPR_Ann_II_8_9_3_01		

		<p>██████████ (2002b), BPD ID A6.7_02, FA_BPR_Ann_II_8_11_2_01</p> <p>██████████ (2004), BPD ID A6.4.1_02; FA_BPR_Ann_II_8_9_2_02</p> <p>High concentration intake – case reports, a.o. Westphal et al (2001), BPD ID A6.12.2_01, FA_BPR_Ann_II_8_12_2_01</p>		
--	--	---	--	--

¹ According to the guidance "Risk characterisation for local effects including sensitisation" – reference to be updated when the guidance is integrated into ECHA guidance.

12.1.3 Absorption

Route	Study	Test substance	Concentration of test substance	Applicability (concentration ranges)	Value
Oral	None, corrosive	/	/	/	Rapid, no quantitative data Assumed 100%
Dermal	None, corrosive	/	/	/	Assumed 100%
Inhalation	None, corrosive	/	/	/	Assumed 100%

12.2 REFERENCE VALUES

12.2.1 Uncertainties and assessment factors

AELshort-term		
Uncertainty	AF	Justification
Interspecies variability	10	Default AF in the absence of substance-specific data
Intraspecies variability	10	Default AF in the absence of substance-specific data
Route to route extrapolation	1	No indication for route-specific differences in systemic toxicity
Time duration extrapolation	1	no additional extrapolation factor for duration is considered for the calculation of the acute AEL from the repeated 90-day oral toxicity study

NOAEL to LOAEL extrapolation	/	
Dose response	/	
Severity of key health effects	/	reduced bw gain at 2100 mg formate/kg bw/d
Overall AF	100	(n.a.)

AELmedium-term		
Uncertainty	AF	Justification
Interspecies variability	10	Default AF in the absence of substance-specific data
Intraspecies variability	10	Default AF in the absence of substance-specific data
Route to route extrapolation	1	No indication for route-specific differences in systemic toxicity
Time duration extrapolation	1	Study duration subchronic
NOAEL to LOAEL extrapolation	/	
Dose response	/	
Severity of key health effects	/	reduced bw gain at 2100 mg formate/kg bw/d
Overall AF	100	(n.a.)

AELlong-term		
Uncertainty	AF	Justification
Interspecies variability	10	Default AF in the absence of substance-specific data
Intraspecies variability	10	Default AF in the absence of substance-specific data
Route to route extrapolation	1	No indication for route-specific differences in systemic toxicity
Time duration extrapolation	1	Study duration chronic
NOAEL to LOAEL extrapolation	/	
Dose response	/	
Severity of key health effects	/	reduced bw gain at 1400 mg formate/kg bw/d
Overall AF	100	(n.a.)

AEC respiratory tract irritation		
Uncertainty	AF	Justification
Interspecies variability	1	local effects, no toxicokinetic default assessment and toxicodynamic default assessment factor needed because of the similarity in local effects among rodents and humans: effects on the respiratory and olfactory epithelium, squamous metaplasia and degeneration, there is no evidence that humans should be more sensitive than rodents
Intraspecies variability	10	Default AF in the absence of substance-specific data
Route to route extrapolation	1	Subchronic inhalation studies
Time duration extrapolation	/	
NOAEL to LOAEL extrapolation	/	
Dose response	/	
Severity of key health effects	/	Local effects: squamous metaplasia and degeneration of the respiratory and olfactory epithelia
Overall AF	10	(n.a.)

12.2.2 AEL setting

Due to its inherent properties (acidic pH, corrosive substance, volatile) it is most likely that formic acid will induce local effects at a lower dose than systemic effects.

Therefore, it seems to be reasonable to do the risk characterisation starting from systemic AELs and local AECs.

In addition, other international AEL are available:

ADI (residues in food, feed) = 3 mg/kg bw/d (EU SANCO D3/AS D, 2005; JECFA, 2003)¹³

Occupational Exposure Limit: EU WEL, MAK/TLV = 5 ppm or 9.5 mg/m³ (8-hour TWA)); IOELV = 5 ppm or 9 mg/m³ (Commission directive 2006/15/EC).

An ArfD was not derived and not required.

The available data (medium term exposure) does not permit to characterize a significant systemic effect in the context of the reduction in body weight, in fact there is no obvious link between the irritant effects (NOAEL: 840 mg/kg) induced by the substance at a high dose, i.e. 2100 mg/kg (LOAEL), and the individual weight loss (decreased food consumption in males but not in females). Finally, in the recovery period, body weight development in males and females was comparable between the high dose and control groups. Therefore the derivation of ArfD value does not seem relevant.

Systemic AEL

¹³ No detailed information can be provided on how the ADI was derived. Despite this, the ADI can be taken up in the CAR because it is in line with the derived AEL_{long-term}.

BPC-43-2022-06B

Systemic toxicity is secondary to local irritant effects. The critical systemic endpoint of formate in the toxicological studies was identified as reduced body weight gain. The NOAELs have been derived from the studies in the most sensitive species showing these effects: the rat and mouse. It is suggested to consider this systemic effect in the risk assessment.

Additional note:

Next to the NOAEL_{systemic} used to derive AELs as reported below, the following NOAEL and LOAEL for local effects are also available:

Species	Route	Study duration	Test substance	Dose setting (mg/kg bw/d)	Critical effect	LO(A)EL and NO(A)EL (mg/kg bw/d)	References
Rat	Oral	Subchronic 90 day feeding study	Potassium diformate	0, 420, 840, 2100 mg formate/kg bw/d	Local: gastric irritation, hyperplastic changes in the stomach	as formate: LOAEL _{local} = 420 NOAEL _{local} < 420	BPD ID A6.4.1_01 FA_BPR_Ann_II_8_9_2_01 [REDACTED] 1998
Rat	Oral	Chronic 2-year feeding study	Potassium diformate	0, 35, 280, 1400 mg formate/kg bw/d	Local: gastric irritation, hyperplastic changes in the stomach and gastrointestinal tract	as formate: LOAEL _{local} = 280 NOAEL _{local} = 35	BPD ID A6.5_01 FA_BPR_Ann_II_8_9_3_01 [REDACTED] 2002a
Mouse	Oral	Carcinogenicity study: 80-week feeding study	Potassium diformate	0, 35, 280, 1400 mg formate/kg bw/d	Local: gastric irritation, hyperplastic changes in the forestomach	as formate: LOAEL _{local} = 1400 (highest concentration tested) NOAEL _{local} = 280	BPD ID A6.7_02. FA_BPR_Ann_II_8_11_2_01 [REDACTED] 2002b
Pig	Oral	Subchronic 140-day feed study	Potassium diformate	0, 149, 359, 760 mg formate/(kg bw/d)	Local: gastric effects - forestomach gastritis and erosion/ulcer	as formate: LOAEL _{local} = 149 NOAEL _{local} < 149	BPD ID A6.4.1_02 FA_BPR_Ann_II_8_9_2_02 [REDACTED] 2004
Pig	Oral	Subchronic > 300-day feed study	Potassium diformate	0, 98, 301 mg formate/kg bw/d	No signs of maternal systemic toxicity or toxicity to reproduction or development at any dose level.	as formate: LOAEL, local > 300 NOAEL, local = 300 (highest concentration tested)	BPD ID A6.5_02 FA_BPR_Ann_II_8_9_4_0_JNS [REDACTED] 2003

For setting of appropriate Reference Values for oral exposures, it is necessary to differentiate between systemic toxicity (decrease in bw) and local effects (irritation on the gastrointestinal tract); Reference Values should be set based on the most sensitive endpoint in the most sensitive species. The most sensitive endpoint is the irritation on the gastrointestinal tract.

However, in the case of Formic Acid, due to its corrosivity, local effects must be expected at all dose levels, and a qualitative RC would be the appropriate approach, assuming that the

effects leading to classification will also occur in repeated exposure and at lower concentrations/area doses, and the effects will be managed by means of CLP, RMM's and PPE.

Derived reference values based on systemic effects are lower than those based on local effects, considering the applied assessment factors. Hence the local effects are covered by the reference values for systemic effects; we will apply the AEL of systemic effects for the quantitative risk assessment.

Acute and Medium-term AEL

Although human exposure is mainly dermal and by inhalation, the PODs are based on oral studies.

The teratogenicity study performed with sodium formate in the rat cannot be used to derive a systemic NOAEL as no maternal systemic toxicity was reached. No other short-term toxicity studies are available.

A medium-term 90-day oral toxicity study performed with potassium diformate in the rat revealed a NOAEL oral, 90-days, rat = 840 mg formate/kg bw/d (based on decreased bw gain at 2100 mg formate/kg bw/d).

Two medium-term 90-day inhalation studies performed with formic acid itself in the rat and mouse are available. In the rat, no systemic effects were observed up to the highest concentration tested 244 mg/m³. In the mouse, a NOAEC of 122 mg/m³ was determined based on the reduced bw gain observed at 244 mg/m³. When taking into account: Minute Volume mouse = 0.041 L/min, BW mouse = 0.030 kg, inhalation = 360 min, then 122 mg/m³ corresponds with a systemic dose of ~60 mg/kg bw/d. However, the RMS is convinced that the systemic NOAELs derived from the inhalation studies are not suitable for the determination of systemic AEL's. The systemic effects seen in the mouse study were most probably secondary to the local effects of respiratory irritation induced by formic acid exposure (NOAEL_{local} = 64 mg/m³, based on histopathological changes in the nasal region). In these studies formic acid itself and not the salts were used. In the oral studies the less corrosive formate salts were used to reveal systemic effects not secondary to the corrosive effects.

In conclusion, for the derivation of the acute and medium-term AEL, the NOAEL of the oral 90-day study in the rat performed with potassium formate was used.

POD acute and medium-term: NOAEL formate, oral, 90-day feeding study, potassium diformate, rat = 840 mg formate/kg bw/d

Oral absorption: 100%

AF: 10 x 10 (no additional extrapolation factor for duration is considered for the calculation of the acute AEL from the repeated 90-day oral toxicity study)

Acute and Medium-term AEL _{systemic} = 8.4 mg formate/kg bw/d
--

Long-term AEL

Long-term toxicity studies are available for the rat and mouse.

The 2-year rat study and 80-week mouse study performed with potassium diformate both revealed a NOAEL oral, long-term = 280 mg formate/kg bw/d (based on decreased bw gain at 1400 mg formate/kg bw/d)

POD: NOAEL formate, oral, 2-year feeding study, potassium diformate, rat = 280 mg formate/kg bw/d

Oral absorption: 100%

AF: 10 x 10

Long-term AEL_{systemic} = 2.8 mg formate/kg bw/d rounded to 3 mg formate/kg bw/d¹⁴

This value corresponds to the ADI.

A NOAEL_{systemic} of 200 mg/kg bw/d is also available (Two-Generation Reproduction Toxicity Study, Rat, oral, feed). However, it can be justified not to derive the AEL_{long term} from this study.

Comparing the results of the 2-generation study (██████████ 2008b) and the combined chronic toxicity and carcinogenicity study, the results of both studies suggest that formate and its salts exhibit only very minor systemic effects. In both studies animals of the high dose show reduced body weights, body weight gains and food consumption. Unfortunately, the selected doses differed slightly in both studies. The mid dose of the chronic study corresponded to 280 mg/kg bw/d and the mid dose of the 2-generation study corresponded to 203 mg/kg bw/d.

	2-generation study		Chronic study	
	Formate [mg/kg bw/d]	Decrease in BW gain [%]	Formate [mg/kg bw/d]	Decrease in BW gain [%]
Low dose	68	-	35	-
Mid dose	203	-	280	-
High dose	677	m: 8.8	1400	m: 27, f: 19

The 2-generation study is a feeding study using sodium formate as test material. No systemic effects including effects on body weight were observed in the first parental generation. However, mean body weights of the high-dose parental F1 males (1000 mg/kg bw/d) of the 2-generation study were statistically significantly decreased during study weeks 9-15 (up to 5.7%). The mean body weight gain of the high-dose F1 males was statistically significantly decreased on several occasions during the study (up to 33.6%). If calculated for the entire treatment period (weeks 0-15) the high-dose F1 males gained about 8.8% less weight than the control males.

It has to be noted that the route of administration was orally via feed. As well as the body weight gain, the food consumption of male animals of the high dose was also reduced in a similar manner (in average about 9% decreased). Thus, the decrease in body weight gain and the reduced food consumption in high dose parental F1 males correlate and could be indicative

¹⁴ We refer to TAB entry TOX-4 as the impact of rounding is less than 10%. Please note that for this CAR, the risk characterization has been performed with the non-rounded 2.8 mg formate/kg bw/d value. The decision for rounding the AEL long-term was taken at HH WG I-2022; however it was decided that there was no need to alter the risk characterization of the CAR. For product approval, the rounded 3 mg formate/kg bw/d value should be used.

of a palatability problem of the highest dose (acidity of the test substance) since the decrease in body weight gain was not seen in the presence of normal food consumption.

The combined chronic toxicity/carcinogenicity study in the rat (██████████ 2002a) was performed via oral administration using potassium formate (1:2) as test material.

In this study, lower body weight and body weight gain than for the controls was seen in the high dose animals together with a minor decrease in food consumption. However, the variation in food consumption was of insufficient magnitude to account for the lower body weight gain. The average decrease in body weight gain accounted in males 27% and in females 19% in this study (104 weeks).

Comparing the results of both studies shows that systemically available formate exhibits only very minor toxicological effects at high doses. Systemic effects other than decreased food consumption, body weight and body weight gain were not observed.

The effects on body weight gain found in the parental F1 animals of the 2-generation toxicity study were, when calculated over the entire treatment period, lower than 10% and could be correlated with the decreased food consumption which may be a hint of palatability problems of the high dose group. Additionally, it should be noted that this minor effect was limited to parental F1 males and was not observed in other generations (e.g. P0).

The effects observed in the chronic feeding study with potassium formate (1:2) were, although also only minor, more pronounced, and not limited to males.

Hence, the minor difference in the established NO(A)ELs of both studies can only be attributed to the minor difference in dose setting. The mid dose, which was the highest dose showing no systemic effects corresponded to 280 mg/kg bw/d formate in the chronic and to 203 mg/kg bw/d formate in the 2-generation study.

In conclusion, the use of the systemic NO(A)EL of 280 mg/kg bw/d formate for derivation of the AEL_{long-term} is justified by the longer treatment period of the chronic study, the more pronounced systemic effects observed in the chronic study and the minor or negligible difference of established NO(A)ELs which can be attributed to the slightly different dose setting of both studies.

Local AECs

Formic acid is classified as corrosive. Formic acid is severely irritating and corrosive to the eyes, skin, and mucous membranes (gastrointestinal and respiratory tract) and may cause permanent damage. Due to the corrosivity of formic acid, local effects must be expected at all dose levels.

Inhalation AEC respiratory tract irritation

A quantitative risk characterisation can be performed as repeated dose 13-week inhalation studies are available performed with formic acid in the rat and mouse. An external reference value (AEC) has been derived for the local effect of respiratory tract irritation:

POD: NOAEC formic acid, inhalation, 13 weeks, rat/mice = 60 mg/m³

AF: 10 x 1

- default assessment factor intraspecies: 10;
- interspecies: assessment factor of 1, please see justification below.

- formic acid causes mainly local effects,

- no toxicokinetic default assessment and toxicodynamic default assessment factor are considered to be required because of the similarity in local effects among rodents and humans: effects on the respiratory and olfactory epithelium, squamous metaplasia and degeneration, there is no evidence that humans should be more sensitive than rodents)

Since there are currently no validated animal tests that deal specifically with respiratory tract irritation, an interspecies assessment factor of >1 could be called for in order to cover this additional uncertainty. However, during HH WGI2022 it was decided that an interspecies AF of 1 is acceptable and that a total assessment factor of 10 is sufficient, mainly due to the fact that FA is likely to be a case of direct/pH-driven chemical action on tissue/cell membranes.

The effect of FA is highly likely a simple destruction of membranes due to the physico-chemical properties (e.g. pH) of the chemical concerned as opposed to a mechanism involving local metabolism (e. g. reactive metabolite). If tissue metabolism is involved, which could lead to the formation of different metabolites at different rates in different species, interspecies dynamic differences on how these metabolites interact with specific targets should be considered.

However, Formic acid is a volatile and strongly corrosive organic acid which is in mammals rapidly metabolized to CO₂ and H₂O. It can be concluded that no toxicologically significant or reactive metabolites are formed and that local irritation due to corrosivity is the most sensitive response and leading health effect. Thus, the mechanism of respiratory irritation is direct pH-reactivity and no further kinetic considerations apply. Furthermore, in terms of toxicodynamic, it can be assumed that rats and humans will respond to the insult in the same way since no significant differences in buffer capacity of cells in respiratory tract against strong acids are expected.

For the following reasons an additional safety factor seems not to be necessary:

- NOAEC derived from validated and reliable subchronic inhalation studies in two species (rat, mice)
- Mechanism of respiratory tract irritation is direct pH-reactivity
- Rodents are obligate nasal breathers with a more complex nasal passage and therefore the upper respiratory tract may be more sensitive than in humans

AEC respiratory tract irritation = 6 mg/m ³
--

(EU workplace exposure limit = 5ppm (9.5 mg/m³), 8-hour time weighted average; IOELV = 5 ppm or 9 mg/m³ (Commission directive 2006/15/EC))

Dermal AEC

Repeated dose dermal studies are not available, and consequently the basis for setting an AEC is lacking.

Therefore a qualitative RC will be performed assuming that the effects leading to classification will also occur in repeated exposure and at lower concentrations/area doses, and the effects will be managed by means of CLP, RMM's and PPE.

AECdermal <2% formic acid: does not need classification

Oral AEC

No oral AEC will be derived because all repeated dose oral studies were performed with the salts, potassium diformate or sodium formate, because of their less irritating potency.

It is known from published human data (Malorny, 1969b; DocIII A6.2-07; section 3.1), that immediately after the drinking of 2 g formic acid as 0.4% aqueous solution transient gastric irritation was observed.

12.2.3 Reference values to be used in Risk Characterisation

Reference	Study	NOAEL (LOAEL)	AF	Correction for oral absorption	Value
AEL _{short-term}	Subchronic 90 day feeding study, rat	as formate: 840 mg/kg bw/d (2100 mg/kg bw/d)	100	1	8.4 mg/kg bw/d
AEL _{medium-term}	Subchronic 90 day feeding study, rat	as formate: 840 mg/kg bw/d (2100 mg/kg bw/d)	100	1	8.4 mg/kg bw/d
AEL _{long-term}	Chronic 2-year feeding study, rat	as formate: 280 mg/kg bw/d (1400 mg/kg bw/d)	100	1	2.8 mg/kg bw/d rounded to 3 mg/kg bw/d ¹⁵
ARfD	not required				
ADI	EU SANCO D3/AS D, 2005; JECFA, 2003				3 mg/kg bw/d
Occupational exposure limit		EU WEL, MAK/TLV (8-hour TWA) IOELV (Commission Directive 2006/15/EC)			5 ppm or 9.6 mg/m ³ 5 ppm or 9 mg/m ³
AEC _{respiratory tract irritation}	Subchronic 13w	Rat: 30 mg/m ³	10	n.a.	6 mg/m ³

¹⁵ We refer to TAB entry TOX-4 as the impact of rounding is less than 10%. Please note that for this CAR, the risk characterization has been performed with the non-rounded 2.8 mg formate/kg bw/d value. The decision for rounding the AEL long-term was taken at HH WG I-2022; however it was decided that there was no need to alter the risk characterization of the CAR. For product approval, the rounded 3 mg formate/kg bw/d value should be used.

	inhalation study, rat/mice	(61 mg/m ³) Mice: 61 mg/m ³ (122 mg/m ³) Overall NOAEC formic acid, inhalation, 13 weeks, rat/mice = 60 mg/m ³			
--	----------------------------	---	--	--	--

12.2.4 Maximum residue limits or equivalent

MRLs or other relevant reference values	Reference	Relevant commodities	Value
default MRL	Art.18(1)(b) Reg 396/2005	all	0.01 mg/kg

12.3 INDUSTRIAL USES

This section has not been evaluated by the CA-BE because the production/formulation process of the active substance is outside the scope of the Biocidal Products Regulation (EU) No 528/2012. As such, exposure estimates for industrial workers during these stages have not been calculated as they are already addressed by other legislation.

Formic acid is severely irritating and corrosive to the eyes, skin, and mucous membranes (gastrointestinal and respiratory tract) and may cause permanent damage. The effect must be managed by means of classification (CLP), Risk Management Measures (RMM's), and Personal Protective Equipment (PPE). The production processes are technically controlled. Workers in industry should be fully trained and protected.

The industry worker exposure during production, filling and mixing processes is routinely determined. The results of 138 measurements made during 2001-2006 indicate that the formic acid concentrations in the air at the workplace did not exceed the threshold limit value of 9.5 mg/m³ (5 ppm; AOEL) at any of the workplaces which cover all types of operations at the production plant.

Conclusion: There is no concern for industrial workers in the production and formulation of the active substance and the biocidal product.

12.4 PROFESSIONAL USES

The biocidal product, Protectol® FM 85, available for professional contractors and experienced farm workers is formulated as a concentrated product containing up to 55% formic acid to be further diluted to the recommended use concentration of 5% for dipping troughs, or to the recommended use concentrations of 4.5 to 19% for fogging. Professionals use products on a prolonged basis, mixing and loading, fogging. Professional contractors are exposed daily to loading fogging machines; experienced farm workers are exposed daily to dipping troughs, and up to 12 times a year for restocking treated animal housings.

12.4.1 Systemic effects

Task/ Scenario	Tier/PPE	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/ AEL (%)	Acceptable (yes/no)
Scenario 1, fogging	1/none	280	2.8	16.431	587	no
	2/impermeable coveralls, boots, gloves and face protection	280	2.8	0.0377	1.3	yes
Scenario 2, footwear disinfection, 5%	1/none	280	2.8	0.212	7.6	yes
	2/coated coveralls, boots, gloves and face protection	280	2.8	0.0329	1.2	yes
Scenario 3, animal feet/transport vehicle disinfection, 5%	1/none	280	2.8	0.0475	1.7	yes
	2/coveralls, boots, gloves and face protection	280	2.8	0.0228	0.8	yes
Scenario 4, restocking of animal housing after ambient temperature fogging	1/ all FA evaporated	280	2.8	11.5	411	no

	2/ventilation until <6 mg/m ³ AND 3/ RMM approach	280	2.8	0.125	4.5	yes
Scenario 4, restocking of animal housing after thermal fogging	1/ all FA evaporated	280	2.8	9	321	no
	2/ventilation until <6 mg/m ³ AND 3/ RMM approach	280	2.8	0.125	4.5	yes

12.4.1.1 **COMBINED SCENARIOS**

Possible scenario combinations were calculated for experienced farmers. Scenarios 2, 3 and 4 (restocking after fogging, footwear & animal feet disinfection) are not unlikely to be combined within a normal working day.

Scenarios combined	Tier/PPE	Systemic NOAEL mg/kg bw/d	AEL mg/kg bw/d	Estimated uptake mg/kg bw/d	Estimated uptake/AEL (%)	Acceptable (yes/no)
Combined restocking of animal housing after fogging and exposure from footwear and animal feet disinfection, experienced farmers						
Scenarios 2,3,4 restocking after ambient temperature fogging, footwear, animal feet disinfection	1/ none	280	2.8	11.76	420	no
	2/ scen 2 & 3: coated coveralls, boots, gloves and face protection; scen 4: <6mg/m ³ FA	280	2.8	0.181	6.5	yes
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection	1/ none	280	2.8	9.26	331	no
Scenarios 2,3,4 restocking after thermal fogging, footwear, animal feet disinfection	2/ scen 2 & 3: coated coveralls, boots, gloves and	280	2.8	0.181	6.5	yes

	face protection; scen 4: <6mg/m ³ FA					
--	--	--	--	--	--	--

12.4.2 Local effects

As a local AEC for respiratory tract irritation is available, a quantitative risk characterisation can be performed.

Task/ Scenario	Tier/PPE	NOAEC mg/m ³	AEC mg/m ³	Estimated inhalation exposure mg/m ³	Estimated exposure / AEC (%)	Acceptable (yes/no)
Scenario 1, fogging	1/none	60	6			
				3.8 (vapour)	63.3	yes
	2/impermeable coveralls, boots, gloves and face protection	60	6			
				3.8 (vapour)	63.3	yes
Scenario 2, footwear disinfection, 5%	1/none	60	6	3.8 (vapour)	63.3	yes
	2/impermeable coveralls, boots, gloves and face protection	60	6	3.8 (vapour)	63.3	yes
Scenario 3, animal feet/transport vehicle disinfection, 5%	1/none	60	6	3.5 (vapour, M&L)	58.3	yes
				1.0 (vapour,	16.7	yes

				applicatio n)		
				0.38 (vapour, post- applicatio n)	6.3	yes
	2/coveralls, boots, gloves and face protection	60	6	3.5 (vapour, M&L)	58.3	yes
				1.0 (vapour, applicatio n)	16.7	yes
				0.38 (vapour, post- applicatio n)	6.3	yes
Scenario 4, restocking of animal housing after ambient temperature fogging	1all FA evaporated	60	6	550	9167	no
	2&3/ <6 mg/m ³ FA	60	6	<6	<100	yes
Scenario 4, restocking of animal housing after thermal fogging	1all FA evaporated	60	6	430	7167	no
	2&3/ <6 mg/m ³ FA	60	6	<6	<100	yes

As formic acid is corrosive at or above a 10% dilution, a qualitative risk characterisation is needed for local dermal and inhalation exposure. This RC is triggered for those BP classified for local effects. In BP where formic acid is present at concentrations that do not trigger classification of the product according to the CLP criteria, RC for local effects is not required.

For use in PT3, the following concentrations are either marketed or made by dilution of a concentrate:

concentration	PT	task	Classification with regard to corrosivity	Hazard category	Exposure foreseen
concentrate					
55%	3.1, 3.2	Mixing and loading For fogging, footwear disinfection, animal feet/vehicle transport disinfection	Skin corr 1B EUH071	high	Yes, accidental, skin, eye, RT
In-use dilution					
19%	3.1	Thermal fogging	Skin corr 1B EUH071	high	No dermal & RT exposure
Up to 5.5%	3.1	Ambient temperature fogging	Skin irrit 2 Eye irrit 2	Low	No dermal & RT exposure
5%	3.2	Footwear disinfection Draining & disposal	Skin irrit 2 Eye irrit 2	Low	Yes, skin eye (accidental)
5%	3.2	Animal feet/transport vehicle disinfection Draining & disposal	Skin irrit 2 Eye irrit 2	low	Yes, skin, RT eye (accidental)

Professional user – concentrate – M&L										
Hazard			Exposure							Risk
Hazard Category	Effects in terms of C&L	Additional relevant hazard information	PT	Who is exposed?	Tasks, uses, processes	Potential exposure route	Frequency and duration of potential exposure	Potential degree of exposure	Relevant RMM&PPE	Conclusion on risk
High	Skin corr. 1B (H314)	pH _{85%} formic acid = -1.6 pH _{5%} formic acid to be determined @ product level EUH071	3	Scenario 1: Professional users Scenario 2&3: experienced farmers	M&L: Dilution of the theoretical product in fogging device/foot bath/hoof bath	Skin Eye RT	Scenario 1-2: 10 minutes per day, Scenario 3: 5 minutes per day Scenario 1: Daily for professionals Scenario 2&3: daily for experienced farmers	Scenario 1-3: 55% Splashes, hand to eye transfer vapour	<u>Product integrated RMM Labelling</u> Labelling according to CLP Instructions for use and storage Labelling for general safety and hygiene measures (see below) <u>Formulation</u> Product formulation which reduces e.g. splashes <u>Packaging</u> Packaging reducing risk for eye exposure by splashes <u>Trained personnel</u> Trained workers/experienced farmers Containment as appropriate Regular cleaning of equipment and work area Avoidance of contact with contaminated tools and objects Training for staff on good practice	ACCEPTABLE +engineering controls +low frequency +short duration +professionals using PPE +professionals following instructions for use +good standard of personal hygiene +professional bystander is expected to use the same set of PPE as the professional user

									<p>Good standard of personal hygiene</p> <p><u>PPE</u></p> <p>Respiratory protection: In case sufficient ventilation cannot be guaranteed</p> <p>Suitable respiratory protection for lower concentrations or short-term effect: Gas filter for acid inorganic gases/vapours such as SO₂, HCl (e.g. EN 14387 Type E). Gas filter for gases/vapours of inorganic compounds (e.g. EN 14387 Type B) Combination filter for gases/vapours of organic, inorganic, acid inorganic and alkaline compounds (e.g. EN 14387 Type ABEK).</p> <p>Suitable respiratory protection for higher concentrations or long-term effect: Self-contained breathing apparatus.</p> <p>The professional bystander needs to observe the same set of PPE as the worker.</p> <p>Hand protection: chemical-resistant gloves</p> <p>Chemical resistant protective gloves (EN 374) Suitable materials also with prolonged, direct contact (Recommended: Protective index 6, corresponding ></p>
--	--	--	--	--	--	--	--	--	--

								<p>480 minutes of permeation time according to EN 374): chloroprene rubber (CR) - 0.5 mm coating thickness butyl rubber (butyl) - 0.7 mm coating thickness fluoroelastomer (FKM) - 0.7 mm coating thickness Polyethylene-Laminate (PE laminate) - ca. 0.1 mm coating thickness Suitable materials for short-term contact (recommended: At least protective index 2, corresponding > 30 minutes of permeation time according to EN 374) polyvinylchloride (PVC) - 0.7 mm coating thickness natural rubber/natural latex (NR) - 0.5 mm coating thickness</p> <p>Eye protection: Tightly fitting safety goggles (cage goggles) (e.g. EN 166) and face shield</p> <p>see respiratory protection</p> <p>Skin and body protection: coveralls, boots Body protection must be chosen depending on activity and possible exposure, e.g. apron, protecting boots, chemical-protection suit (according to EN 14605 in case of splashes or EN ISO</p>
--	--	--	--	--	--	--	--	---

									13982 in case of dust).	
									<u>General safety and hygiene measures</u> Avoid contact with skin, eyes and clothing. Wash hands before breaks and immediately after handling the product. Remove and wash contaminated clothing and gloves, including the inside, before re-use	

Professional user – dilution – draining/disposal										
Hazard			Exposure						Risk	
Hazard Category	Effects in terms of C&L	Additional relevant hazard information	PT	Who is exposed?	Tasks, uses, processes	Potential exposure route	Frequency and duration of potential exposure	Potential degree of exposure	Relevant RMM&PPE	Conclusion on risk
Low	5% formic acid: Skin irrit 2 (H315) Eye irrit 2 (H319)	pH to be determined (product evaluation)	3	experienced farmers	Draining and disposal of trough content	Skin, eye	10 min per day Daily	5% FA Splashes, hand to eye transfer vapour	<u>RMM</u> Trained workers respecting general safety and hygiene measures (see below) Good standard of general ventilation Avoidance of contact with contaminated tools and objects Training for staff on good practice Good standard of personal hygiene <u>PPE</u> Hand protection: chemical-resistant gloves are recommended	ACCEPTABLE +engineering controls +reversible effect +professionals following instructions for use +experience expected +professional bystander is expected to use the same set of PPE as the professional user

									<p>Details: see M&L of concentrate above</p> <p>Skin and body protection: coveralls, boots, face/eye protection</p> <p><u>General safety and hygiene measures</u></p> <p>Avoid contact with skin, eyes and clothing. Wash hands before breaks and immediately after handling the product. Remove and wash contaminated clothing and gloves, including the inside, before re-use</p>	
--	--	--	--	--	--	--	--	--	---	--

12.4.3 Conclusion

PT3.1: Fogging animal housing by contractors

Exposure for fogging animal housing was assessed for contractors. The assessment includes exposure at re-entry for professional bystanders. Systemic exposure was determined for the dermal and inhalation route. A quantitative assessment was done for inhalation of vapour. Where relevant, a qualitative assessment was included for local dermal and inhalation exposure.

Because during fogging activities nobody is inside the building, it is considered that only the mixing and loading phase needs to be taken into consideration for risk characterisation.

During the manual dilution of the concentrated biocidal product into the ready-to-use solution, professionals without PPE may be exposed to spills of the concentrate. Due to the corrosive properties of the concentrate, professionals must use PPE (impermeable coveralls, gloves, boots and face protection) to prevent exposure to skin and eyes, and this should be advised on the label. The use by trained professionals, the short duration of the exposure during mixing & loading, the suggested RMM for skin and eye, and general safety and hygiene measures should make the risk for local dermal exposure acceptable.

Sufficient ventilation should be advised on the label for the mixing and loading phase. In case of spillage of the concentrated product (55 % formic acid), the onset of odour and irritant symptoms associated with formic acid exposure would be expected shortly after the exposure begins. Fortunately formic acid has good warning properties. Nevertheless, in situations where sufficient ventilation cannot be guaranteed, RPE (as defined in the SDS of Protectol[®] FM 85) are advised due to the acidity of the concentrate.

For the in-use dilution (up to 19%), local dermal effects are excluded as no workers are expected to be present in the area to be treated.

Total systemic exposure is mostly determined by the dermal exposure during mixing and loading. The use of appropriate PPE (tier 2: coveralls, gloves, boots, face protection) reduces total systemic exposure considerably. Taking these PPE into account, risk of adverse effects is minimised. Thus, with the assumption that the obligatory PPE are used, the total internal dose is below the long-term AEL for formic acid.

Even though formic acid is a volatile substance (vapour pressure >0.01 Pa at 20°C), for the mixing and loading phase of the fogging scenario, acceptable exposure levels can be demonstrated for workers exposed to formic acid vapours even at tier 1. In situations where sufficient ventilation cannot be guaranteed, RPE will be required.

Inhalation exposure at re-entry is acceptable only when ventilation measures are in place to reduce the FA concentration in air to below 6 mg/m³. The ventilation time required to reduce the concentration in air to this level depends on the ventilation rate (2/h: fogging at ambient temperature: 136 min, thermal fogging: 129 min; 10/h: fogging at ambient temperature: 28 min, thermal fogging: 26 min). Therefore re-entry must only occur after appropriate ventilation of the treated area; appropriate

RMM to attain an acceptable FA concentration in air should be included on the label (ventilation time or measurement of FA concentration before re-entry).

Conclusion: There is no concern for professionals using the biocidal product during fogging of animal housing PT3.1, when appropriate RMM are applied.

RMM:

M&L: impermeable coveralls, boots, gloves and face protection

Appropriate RPE when ventilation is insufficient

Application: not relevant

Post-application: impermeable coveralls, boots, gloves and face protection

Re-entry : ventilation should be sufficient to reduce the concentration of FA in air to levels below the AEC_{respiratory tract irritation} (ventilation time or measurement of FA concentration before re-entry).

PT3.2: Dipping troughs by agricultural workers: footwear disinfection, animal feet/transport vehicle disinfection

Exposures for disinfection of footwear, animal feet and transport vehicles in dipping troughs were assessed for agricultural workers. Systemic exposure was determined for the dermal and inhalation route. A quantitative assessment was done for inhalation of vapour. Where relevant, a qualitative assessment was included for local dermal and inhalation exposure.

During the manual dilution of the concentrated biocidal product in the dipping trough (footwear disinfection, animal feet disinfection) to the ready-to-use solution, Professionals without PPE may be exposed to spills of the concentrate. Due to the corrosive properties of the concentrate, professionals must use PPE (coveralls, gloves, boots and face protection) to prevent exposure to skin and eyes, and this should be advised on the label. The use by trained professionals, the short duration of the exposure during mixing & loading, the suggested RMM for skin and eye, and general safety and hygiene measures should make the risk for local dermal exposure acceptable.

Dermal exposure to the in-use dilution of 5% triggers a local RA as the in-use dilution is classified as skin/eye irritant. Care should be taken when coming into contact with the content of the trough, and RPE do apply: coveralls, boots, gloves and face protection.

Good ventilation should be advised on the label for the mixing and loading phase. Exposure to vapour during mixing and loading for these dipping tasks is below the AEC for respiratory tract irritation, even without the use of RPE. However, in situations where sufficient ventilation cannot be guaranteed, RPE will be required. In case of spillage of the concentrated product (55% formic acid), the onset of odour and irritant symptoms associated with formic acid exposure would be expected shortly after the exposure begins. Fortunately, formic acid has good warning properties. No risk is identified for inhalation of the in-use dilution (5%).

Total systemic exposure is less than the AEL and therefore indicates no concern for the agricultural workers preparing, using and emptying troughs for boots, vehicles and livestock PT3.2., using a concentrate of 55% FA and an in-use dilution of 5% FA.

Conclusion: There is no concern for professionals using the biocidal product preparing, using and emptying dipping troughs PT3.2, when appropriate RMM are applied.

RMM:

M&L: coveralls, boots, gloves and face protection

Appropriate RPE when ventilation is insufficient

Application: not relevant

Post-application: coveralls, boots, gloves and face protection (5% FA in-use dilutions)

Combined scenarios

Possible scenario combinations were calculated for experienced farmers.

Agricultural workers: restocking of treated animal housings, footwear disinfection and animal feet disinfection

For farmers, scenarios 2, 3 and 4 (disinfection of footwear in dipping troughs; animal feet disinfection; restocking of animal housing after disinfection) are not unlikely to be combined within a normal working day. The use of PPE as described for the respective scenarios will lead to acceptable systemic and local dermal exposure when these tasks are performed by the same agricultural worker on the same day. For inhalation of formic acid vapours, no addition of exposure levels is performed; only the highest exposure level in air is considered relevant, and the same exposure-reducing measures should be followed as those for single restocking and dipping scenarios. Appropriate RPE should be used when ventilation is insufficient. RMS BE considers the exposure of agricultural workers during combined restocking and use of disinfection baths acceptable when an in-use dilution of 5% formic acid is applied.

Inhalation exposure at re-entry of treated animal housings is acceptable only when ventilation measures are in place to reduce the FA concentration in air to below 6 mg/m³.

Conclusion:

There is no concern for professionals during combined use of the biocidal product for restocking of treated animal housings PT3.1, and for preparing and using dipping troughs PT3.2, when appropriate PPE are applied.

General remark:

The main issue identified is the high vapour pressure of formic acid and the resulting inhalation of formic acid vapours.

However, eCA BE is convinced that this should not be the decisive factor in identifying safe uses for Formic acid, and that these concerns should be dealt with at product authorization level. When required, possible refinements that can be suggested include improved assessment factors for ventilation, the use of air measurements and identification of acceptable RMM per type of application.

12.5 NON-PROFESSIONAL USERS

N.A.; the PT3 products presented here are not intended for use by non-professionals.

12.6 SECONDARY (INDIRECT) EXPOSURE AS A RESULT OF USE

12.6.1 Systemic effects

Secondary exposure of professionals:

See 12.4 professional uses – restocking of treated animal housing after (ambient temperature/thermal) fogging.

12.6.1.1 COMBINED SCENARIOS

Secondary exposure of professionals:

See 12.4 professional uses – restocking of treated animal housing after (ambient temperature/thermal) fogging.

12.6.2 Local effects

Secondary exposure of professionals:

Re-entry after fogging: ventilation should be sufficient to reduce the concentration of FA in air to levels below the AEC_{respiratory tract irritation} (ventilation time or measurement of FA concentration before re-entry).

12.6.3 Conclusion

Secondary exposure of professionals:

There is no concern for indirect exposure as a result of the use of the biocidal product to disinfect animal housing PT3, on the condition that entry only occurs after appropriate ventilation of the treated area.

Secondary exposure of the general public:

For the representative products in this CAR, indirect exposure of the general public should be avoided by implementation of appropriate RMM, considering the volatility and corrosive properties of the a.s., and the fact that PPE and RPE are not applicable for the general public.

The following RMM is proposed:

Use is not authorized in areas where public can be received/present.

As access of the general public to treated areas needs to be excluded by RMM, the RA for secondary exposure of the general public has not been considered further.

12.7 INDIRECT EXPOSURE VIA FOOD

Due to its rapid turnover and unlikely accumulation, an estimation of exposure of humans to formic acid residues through diet as a consequence of animal house disinfection was not considered.

It is proposed that assessment of dietary risk for humans and livestock be undertaken at biocidal product authorisation.

We refer to Appendix II for a tentative approach to the draft 'Guidance on the BPR V III HH-Assessment & Evaluation, Section 6: Guidance on Estimating Livestock Exposure to Active Substances used in Biocidal Products'.

If ventilation is considered as a risk mitigation measure for workers restocking animal housings, then animals would be exposed to max 6 mg a.s./m³ (AEC for respiratory tract irritation). This would lead to values for inhalation exposure between 0.51 and 1.38 mg formic acid/kg bw/d, exceeding the trigger value of 0.004 mg/a.s./kg bw/d, calling for refinement to obtain a safe use.

However, if we consider the maximum percentage of Formic Acid which is considered safe in animal feed (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03), then exposure to Formic Acid in feed would greatly surpass the amount exposed to via treated animal housing (max exposure to formic acid via feed and water: between 785 and 1286 mg formic acid/kg bw/d, depending on the animal species). See also section 8.7.

Moreover, toxicokinetic studies referred to in section 3.1 show that formic acid is fully metabolised in the animal leaving no residues in the animal products. In addition, formic acid is used as a food preservative and has been granted GRAS status (Generally Regarded As Safe) by the US FDA regulatory authority¹⁶.

16

<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?fr=186.1316&SearchTerm=formic%20acid>

12.8 PRODUCTION / FORMULATION OF ACTIVE SUBSTANCE

In accordance with the Commission Document agreed at the 22nd CA meeting in September 2006, detailed information on exposure associated with the manufacturing process is not required for biocidal product risk assessment.

13 RISK CHARACTERISATION FOR THE ENVIRONMENT

The risks to the environment resulting from the use of formic acid as a PT3 biocide are summarised in the paragraphs below.

The product, Protectol® FM 85, is intended to be used in a wide variety of products all intended as disinfectants under PT3. The uses assessed here is the use as a disinfectant for footwear and animal's feet.

Direct emissions are to the manure/slurry storage followed by indirect emissions to soil, groundwater and surface water, and to the STP followed by indirect emissions to surface water, soil and groundwater.

The air compartment is potentially exposed in scenarios 2 (animal's feet) and 3 (animal housing, fogging), but unlikely to be a significant concern due to hazard data and scale of exposure, and therefore not further considered.

13.1 ATMOSPHERE

The vapour pressure of 42.71 hPa (20 °C; ██████████; BPD ID A3_01) and the Henry's Law Constant of 0.16 Pa.m³/mol (20 °C; ECT Oekotoxicologie GmbH; BPD ID A3_11) indicate low to moderate potential for volatilization and evaporation from water and wet surfaces.

Conclusion:

The atmosphere is not considered a compartment of concern.

13.2 SEWAGE TREATMENT PLANT (STP)

Summary table on calculated PEC/PNEC values	
	PEC/PNEC _{STP}
Scenario 1 (footwear)	< 4.00x10 ⁻⁴
Scenario 2 (animal's feet)	< 4.85x10 ⁻²
Scenario 3 (animal housing, fogging)	< 1.10x10 ⁻³

Conclusion:

The PEC/PNEC_{stp} are all below 1.

13.3 AQUATIC COMPARTMENT

Summary table on calculated PEC/PNEC values	
	PEC/PNEC _{water}
Scenario 1 (footwear)	≤ 3.64x10 ⁻³

Scenario 2 (animal's feet)	$\leq 1.22 \times 10^{-1}$
Scenario 3 (animal housing, fogging)	$\leq 2.75 \times 10^{-3}$

Conclusion:

All PEC/PNEC_{water} are below 1.

13.4 TERRESTRIAL COMPARTMENT

Calculated PEC/PNEC values	
	PEC/PNEC _{soil}
Scenario 1 (footwear)	≤ 0.219
Scenario 2 (animal's feet)	≤ 0.196
Scenario 3 (animal housing, fogging)	$\leq 1.73 \times 10^{-3}$

All PEC/PNEC ratios for soil are below 1.

Conclusion:

The risk for the soil compartment is considered acceptable.

13.5 GROUNDWATER

The PEC_{groundwater} values are compared to the allowed maximum concentration of 0.1 µg/L (98/8/EC, Annex VI, art. 82).

Calculated PEC values (TIER 1*)	
	PEC _{groundwater} (µg/L)
Scenario 1 (footwear)	7.29×10^1
Scenario 2 (animal's feet)	6.51×10^1
Scenario 3 (animal housing, fogging)	2.76
* Porewater concentration	

The calculated porewater concentrations (as an estimate for the groundwater concentration) are above the threshold value of 0.1 µg/L. To obtain more realistic groundwater concentrations, the modelling software FOCUS PEARL v.4.4.4. is used to simulate leaching to groundwater.

The FOCUS PEARL calculations are presented in §13.7 (Aggregated exposure) and show that the modelled groundwater concentrations are well below the threshold value of 0.1 µg/L.

Conclusion:

The risks for the groundwater compartment are considered acceptable.

13.6 PRIMARY AND SECONDARY POISONING

13.6.1 Primary poisoning

Not considered relevant.

13.6.2 Secondary poisoning

Conclusion:

Not considered relevant.

13.7 AGGREGATED EXPOSURE (COMBINED FOR RELEVANT EMISSION SOURCES)

Formic acid is intended to be used as an active substance for biocidal products in a wide variety of product types: PT2, PT3, PT4, PT5 and PT6. An aggregated exposure assessment is conducted by summing up all release streams for which an overlap in time and space is to be expected.

Two main release routes separated in time and/or space are considered:

1. STP route: PT2, PT4, PT6;
2. Manure route: PT3 (Footwear, Animal feet), PT5.

For the STP route, it is considered that the emissions of PT2, PT4 and PT6 are redirected to the same STP. For the manure route, it is considered that the representative products for the respective PT3 and PT5 uses are used on the same farm. In the following paragraphs, the aggregated exposure of those two main release routes is elaborated. Releases to the air compartment are considered not relevant.

Note that in section §9 for the PT3 scenarios (Footwear and Animal feet), emissions were doubled: the same emission was considered to be directed to both the manure storage and the STP (total fraction emitted >1). To avoid double emissions, it is for the aggregated exposure calculation considered that all emissions from disinfection of footwear and animal feet are directed to the manure storage.

Disinfection of animal housing by means of fogging (PT3, scenario 3) is not included in the aggregated exposure exercise for the following reasons:

- According to the ESD for PT3, fogging is only carried out in exceptional cases.
- The aggregated exposure exercise considers a dairy cows farm (see below, §13.7.2.1) whereas the fogging scenario considers a veal calves farm. No overlap in time and space is thus to be expected.
- The emissions of the fogging scenario are 1 or 2 orders of magnitude lower as compared to the other 'manure' scenarios.

13.7.1 STP route

Not relevant for PT3.

13.7.2 Manure route

13.7.2.1 EMISSION ESTIMATION

It is considered that the representative products for PT3 (Footwear, Animal feet) and PT5 (Animal drinking water) are applied on the same farm and that the emission streams are collected in the same manure/slurry storage pit. For PT5 only scenario 1a (disinfection of animal drinking water at a use concentration of the representative biocidal product of 0.2 %) is considered since this concentration is in line with the EFSA recommendations.

The sum of emissions for the aggregated exposure of the manure route is based on the concentration of the biocide (active ingredient) in soil (mg/kg) in the case of an immission standard for nitrogen and land application on grassland (PIECgrs-N_i1,i2,i3,i4) and arable land (PIECars-N_i1,i2,i3,i4) respectively.

The PT3 scenario 'Disinfection of animal feet' is only elaborated for dairy cows. Therefore, a dairy cows farm is considered to calculate the aggregated emissions. The emissions are summarised in the table below.

Summary of aggregated emissions for the manure route			
	PIECgrs4-N_degr [mg/kg_wwt]	PIECarab-N [mg/kg_wwt]	Remarks
PT3 – Footwear	0.33	0.10	dairy cows
PT3 – Animal feet	5.25	1.53	dairy cows
PT5 – Animal drinking water (scenario 1a – 0.2% b.p.)	3.66	1.09	dairy cows
SUM	9.24	2.72	aggregated emission

13.7.2.2 FATE AND DISTRIBUTION IN EXPOSED ENVIRONMENTAL COMPARTMENTS

Identification of relevant receiving compartments based on the exposure pathway									
	Fresh-water	Sediment	Sea-water	Seawater sediment	STP	Air	Soil	Ground-water	Other
Manure route	+	(-)	-	-	-	(+)	++	+	(-)

++ *Compartment directly exposed*

- *Compartment not exposed*

+ *Compartment indirectly exposed*

() *Compartment potentially exposed [but unlikely to be a significant concern due to hazard data and / or scale of exposure]*

Input parameters (only set values) for calculating the fate and distribution in the environment			
Input	Value	Unit	Remarks
Molecular weight	46.03	g/mol	
Melting point	8	°C	
Boiling point	100.23	°C	
Vapour pressure (at 12 °C)	2400	Pa	
Water solubility (at 12 °C)	1.09x10 ⁶	mg/l	
Log10 Octanol/water partition coefficient	-2.10	---	(pH 7)
Organic carbon/water partition coefficient (Koc)	30	l/kg	(pH 7)
Henry's Law Constant (at 12 °C)	0.101	Pa/m ³ /mol	
Acid dissociation constant	3.7	---	Predominant species at a pH of 7 is formate, which is reflected in the pH dependent Koc.
Biodegradability	Ready biodegradable		
DT50 for degradation in soil (12 °C)	1	day	

13.7.2.3 **CALCULATED AGGREGATED Σ PEC VALUES**

The maximum Σ PEC values per compartment are presented in the table below.

Summary table on calculated ΣPEC values					
	Σ PEC _{STP}	Σ PEC _{water}	Σ PEC _{sed} ¹	Σ PEC _{soil,twa} ²	Σ PEC _{GW}
	[mg/L]	[mg/L]	[mg/kg _{wwt}]	[mg/kg _{wwt}]	[µg/L]
Manure route	N/A	1.145E-02	see Σ PEC _{water} ¹	4.444E-01	1.145E+02

1 Since the PNEC sediment was calculated according to the equilibrium partitioning method, the risk assessment for freshwater covers that for the sediment.

2 Initial concentration after sludge application considering the average time for the terrestrial ecosystem. The PNEC_{soil} is derived by equilibrium partitioning from a PNEC_{aquatic} for chronic exposure.

13.7.2.3.1 Manure route: refinement of the exposure calculation

The calculation of the groundwater concentration is further refined using FOCUS PEARL v.4.4.4. to simulate leaching to groundwater, taking into consideration the specific parameters and formulas indicated according to the TAB v2.

In the table below, the FOCUS PEARL input parameters for Formic Acid are summarised.

PEARL input parameters for substance Formic Acid			
Parameter	Value	Unit	Remarks
<u>GENERAL</u>			
Molecular weight	46.03	g/mol	
Vapour Pressure	2400	Pa	at 12°C
Water solubility	1.00x10 ⁶	mg/l	maximum allowed value
<u>Freundlich sorption</u>			
Kom	17.4	L/kg	pH 7, 20°C (Kom = Koc/1.724)
Freundlich sorption exponent (1/n)	1	[-]	TAB v2, ENV 22 (conservative value)
<u>Transformation</u>			
Half-life	1	d	
Molar activation energy	54	kJ/mol	TAB v2, ENV 23
<u>Crop</u>			
Coefficient for uptake by plant	0	[-]	TAB v2, ENV 23

Simulation was run for both grassland (alfalfa) and arable land (maize) (cfr. TAB v2, ENV 165).

In the case of alfalfa, the scenario considers 4 manure/slurry applications per year on fixed dates 1st of March, 23rd of April, 15th of June and 7th of August (considering 53 days between application) and 5 cm incorporation depth. In the case of maize, one manure/slurry application per year 20 days before crop emergence and 20 cm incorporation depth is considered.

The application rate of the active substance *Appl_rate* [kg/ha] at one specific application date as necessary input parameter in FOCUS groundwater models is calculated on basis of the aggregated predicted initial environmental concentrations (Σ PIEC).

The application rate for the grassland scenario is calculated by:

$$Appl_rate_{grass} = \Sigma PIEC_{grs} \times RHO_{soil_wet} \times DEPTH_{grassland} \times 10^{-2} = \Sigma PIEC_{grs} \times 0.85$$

With:

$Appl_rate_{grs}$ = concentration of active ingredient in grassland soil after 1 manure slurry application based on the nitrogen immission standard for grassland [kg/ha]

Σ PIEC_{grs} = aggregated concentration of the active ingredient in grassland soil after 1 manure/slurry application based on the nitrogen immission standard for grassland [mg/kg] according to OECD ESD PT 18 No.14 (2006)

RHO_{soil_wet} = wet bulk soil density = 1,700 kg/m³

$DEPTH_{grassland}$ = mixing depth with soil for grassland = 0.05 m

The application rate for the arable land scenario is calculated by:

$$Appl_rate_{ar_maize} = \Sigma PIEC_{ars} \times RHO_{soil_wet} \times DEPTH_{arableland} \times 10^{-2} = \Sigma PIEC_{ars} \times 3.4$$

Appl_rate_{ar_maize} = initial concentration of the active substance in soil of arable land after 1 manure/slurry application based on the nitrogen immission standard for arable land [kg/ha]

$\Sigma PIEC_{ars}$ = initial aggregated concentration of the active substance in soil of arable land after 1 manure/slurry application based on the nitrogen immission standard for arable land [mg/kg] according to OECD ESD PT 18 No.14 (2006) and to the Addendum (Nov.2015)

RHO_{soil_wet} = wet bulk soil density = 1,700 kg/m³

$DEPTH_{arableland}$ = mixing depth with soil for arable land = 0.2 m

PEARL input parameters for Application Schemes				
Parameter	Value		Unit	Remarks
	Grassland	Arable Land		
Crop	Alfalfa	Maize	[-]	
Application type	incorporation	incorporation	[-]	
Date(s)	01 March, 23 April, 15 June, 07 August	20 days before emergence		TAB v2, ENV 165
Mixing depth	0.05	0.2	m	
$\Sigma PIEC$	9.24	2.72	mg/kg _{wwt}	
Dosage (Appl_rate)	7.85	9.25	kg/ha	

PEARL was then run for the nine available locations for each application scheme. Repeat interval for years was set to 1. The resulting groundwater concentrations closest to the 80th percentile are presented below.

PEARL groundwater assessment [µg/L]		
Location	Grassland	Arable Land
Chateaudun	0.000000	0.000000
Hamburg	0.000000	0.000000
Jokioinen	0.000000	N/A
Kremsmuenster	0.000000	0.000000
Okehampton	0.000000	0.000000
Piacenza	0.000004	0.000000
Porto	0.000000	0.000000
Sevilla	0.000000	0.000000
Thiva	0.000000	0.000000

All modelled groundwater concentrations are below the threshold value of 0.1 µg/L.

13.7.2.4 **AGGREGATED RISK CHARACTERISATION**

The calculated aggregated Σ PEC/PNEC values for the manure route are summarised in the table below.

Summary table on calculated aggregated ΣPEC/PNEC values for the manure route					
	Σ PEC/PNEC _{STP}	Σ PEC/PNEC _{water}	Σ PEC/PNEC _{sed} ¹	Σ PEC/PNEC _{soil,twa}	Σ PEC _{GW} ²
Manure route	N/A	5.73E-03	see Σ PEC/PNEC _{water} ¹	3.45E-01	1.15E+02
<p>1 Since the PNEC sediment was calculated according to the equilibrium partitioning method, the risk assessment for freshwater covers that for the sediment.</p> <p>2 Tier 1 porewater concentration</p>					

The groundwater concentrations are below the threshold of 0.1 µg/L after refinement of the PEC calculations using FOCUS PEARL.

13.8 SUMMARY OF THE RISK ASSESSMENT FOR THE ENVIRONMENT

Summary table on environmental risk assessment					
	STP	Fresh water	Sediment	Soil	Groundwater
Scenario 1 (Footwear)	acceptable	acceptable	acceptable	acceptable	acceptable (FOCUS PEARL)
Scenario 2 (Animal's feet)	acceptable	acceptable	acceptable	acceptable	acceptable (FOCUS PEARL)
Scenario 3 (Animal housing, fogging)	acceptable	acceptable	acceptable	acceptable	acceptable (FOCUS PEARL)
Aggregated exposure (manure route)	N/A	acceptable	acceptable	acceptable	acceptable (FOCUS PEARL)

Conclusion:

The risks for the environment from the intended uses of the representative product for PT3 are acceptable.

The risks for the environment from the aggregated exposure of biocidal products containing formic acid are acceptable.

14 RISK CHARACTERISATION FOR THE PHYSICO-CHEMICAL PROPERTIES

Formic acid is thermally stable up to 350 °C at which combustion starts. It is a flammable liquid (flash point in closed cup: 49.5 °C) with a high auto-ignition temperature of 528 °C. Thermal breakdown and combustion products are carbon monoxide and water/hydrogen. Pure formic acid is not corrosive to metals, while FA85% is corrosive to steel, but not corrosive to aluminium (UN test 37.4 C1). Formic acid is not explosive and has no oxidizing properties.

The biocidal product Protectol® FM 85 contains to 85% of the active substance formic acid and [REDACTED]. Physical-chemical properties are expected to be similar to the active substance. It is not a flammable liquid (flash point in closed cup: 73.5°C). Protectol® FM 85 is stable in terms of ambient storage conditions. As an acidic product, Protectol® FM 85 is in general compatible with other acid and neutral pH solutions. Contact with strongly alkali solutions should be avoided as neutralization of Protectol® FM 85 (as is the case for many concentrated acids) with alkalis may result in a vigorous reaction. Protectol® FM 85 containing formic acid may have a reducing effect and therefore compatibility with strong oxidizers such as phosphorus pentoxide should be evaluated carefully. As with many concentrated acids contact of Protectol® FM 85 with powdered metals and inorganic catalysts should be avoided. FA 85% is corrosive to steel, but not corrosive to aluminium. Protectol® FM 85 is corrosive and as such can be incompatible with some metals and other materials of construction (BPD IDs A3_02, A3_03, A3_05, A3_06, B3_02, B3_05).

15 MEASURES TO PROTECT MAN, ANIMALS AND THE ENVIRONMENT

Professional users need to be trained and instructed on the proper use of the formic acid, its handling, storage, disposal, the selection and use of protective equipment, and First Aid measures. Safety Data Sheets (SDS) should be supplied.

Consumer products should be labelled with the same or similar information. The labels should transfer the information contained in the SDS into the consumer's language, taking into account the concentration of formic acid.

Human exposure:

Formic acid is corrosive for skin and eye at concentrations from 10% onwards. Concentrations from 2% onwards are skin and eye irritants. Personal protection should be applied, as recommended by classification and labelling, and as established through the risk assessment. See the relevant sections in the CAR for details.

Due to the high volatility and corrosiveness of formic acid, care should be taken when there is potential for exposure via the inhalation route for professionals and professional bystanders. For the professional user and bystander appropriate RPE are required when handling high formic acid concentrations in conditions of insufficient ventilation. When using lower in-use concentrations, the end user should apply ventilation-related risk mitigation measures to protect himself and possible bystanders. See the relevant sections in the CAR for details. Ventilation-related RMM should be defined at product authorization level, especially if the risk assessment cannot be refined in other ways (e.g. by performing actual measurements of FA concentrations in air).

If an unacceptable risk is identified for the general public, indirect exposure should be avoided by implementation of appropriate RMM, considering the volatility and corrosive properties of the a.s., and the fact that PPE and RPE are not applicable for the general public.

If an unacceptable risk is identified for animal health, indirect exposure should be avoided by implementation of appropriate RMM.

At product level, the risk assessment should take into account the in-use dilutions for which efficacy is supported by sufficient testing. Effects of other parameters on the risk assessment, such as the necessary contact time and drying time of the mixture, should also be taken into account.

Exposure through the dietary route and livestock exposure: due to its rapid turnover and unlikely accumulation, an estimation of exposure of humans to formic acid residues through diet, and exposure of livestock to FA were not considered. However, as for most of the applications presented in the CAR a rinsing step is not foreseen, it is proposed that the need for assessment of dietary risk for humans and livestock be evaluated at biocidal product authorisation.

Environmental precautions:

Do not empty into drains.

PART D : APPENDICES

APPENDIX I: LIST OF ENDPOINTS

Chapter 1: Identity, Physical and Chemical Properties, Classification and Labelling

Active substance (ISO Name)

Formic Acid

Product-type

2, 3, 4, 5, 6

Identity

Chemical name (IUPAC)

Formic Acid

Chemical name (CA)

Formic Acid

CAS No

64-18-6

EC No

200-579-1

Other substance No.

Minimum purity of the active substance as manufactured (g/kg or g/l)

99%

Identity of relevant impurities and additives (substances of concern) in the active substance as manufactured (g/kg)

This information is contained in the BASF PT3 Confidential Annex

Molecular formula

CH₂O₂

Molecular mass

46.025

Structural formula

HCOOH

Physical and chemical properties

Melting point (state purity)

8°C

Boiling point (state purity)

100.23

Thermal stability / Temperature of decomposition

350°C

Appearance (state purity)

Liquid (20°C)

Relative density (state purity)

D₄²⁰ = 1.2195

Surface tension (state temperature and concentration of the test solution)

At 20 °C: 71.5 mN/m

Vapour pressure (in Pa, state temperature)	At 20 °C: 42.71 hPa At 25 °C: 54.96 hPa At 50 °C: 170.7 hPa
Henry's law constant (Pa m ³ mol ⁻¹)	At 20 °C: 0.16 Pa.m ³ /mol
Solubility in water (g/l or mg/l, state temperature)	pH 5 at ___ °C: pH 9 at ___ °C: pH [X] at ___ °C: Temperature dependence was not investigated due to complete miscibility.
Solubility in organic solvents (in g/l or mg/l, state temperature)	Miscible at ratios: 1:9, 1:1 and 9:1 Miscible at 20 and 30 °C Corresponding to: > 850 g/L N,N-dimethylformamide > 929 g/L 1,4-dioxane > 1190 g/L Dichloromethane
Stability in organic solvents used in biocidal products including relevant breakdown products	Waived, since no organic solvent is used in the biocidal product.
Partition coefficient (log P _{ow}) (state temperature)	pH 5 at 20 °C: -1.9 pH 9 at 20 °C: -2.1 pH [X] at 20 °C: -2.3
Dissociation constant	At 20 °C: pK _a = 3.70
UV/VIS absorption (max.) (if absorption > 290 nm state ε at wavelength)	n.a.
Flammability or flash point	49.5°C
Explosive properties	The substance is not explosive.
Oxidising properties	The substance is not an oxidising liquid
Auto-ignition or relative self-ignition temperature	528°C
Classification and proposed labelling with regard to physical hazards	H290 H226

with regard to human health hazards

H302
H331
H314
H318
EUH071

with regard to environmental hazards

Chapter 2: Methods of Analysis

Analytical methods for the active substance

Technical active substance (principle of method)

Titration with sodium hydroxide
Confirmatory method: GC-MS chromatography

Impurities in technical active substance (principle of method)

Determination of Water by Karl-Fischer titration

Analytical methods for residues

Soil (principle of method and LOQ)

UV absorption after stoichiometric , enzyme-catalyzed reduction of NAD⁺ to NADH by formic acid

Formic acid (formate) is quantitatively oxidized to bicarbonate by nicotinamide adenine dinucleotide (NAD) in the presence of formate dehydrogenase (FDH).

$$\text{Formate} + \text{NAD}^+ + \text{H}_2\text{O} \xrightarrow{\text{FDH}} \text{bicarbonate} + \text{NADH} + \text{H}^+$$

The amount of NADH formed is stoichiometric to the amount of formic acid. The increase in NADH is measured by means of its light absorbance at 334, 340 or 365 nm

10 mg/kg

Air (principle of method and LOQ)

Ion chromatography; LOQ = 0.1 µg

Water (principle of method and LOQ)

UV absorption after enzymatic reaction; LOQ = 0.2 mg/L in drinking water; LOQ = 0.2 mg/L in surface water

Body fluids and tissues (principle of method and LOQ)

0.2 mg/L

Food/feed of plant origin (principle of method and LOQ for methods for monitoring purposes)

UV absorption after enzymatic reaction; LOQ = 0.2 mg/L

Food/feed of animal origin (principle of method and LOQ for methods for monitoring purposes)

UV absorption after enzymatic reaction; LOQ = 0.2 mg/L

Chapter 3: Impact on Human Health

Absorption, distribution, metabolism and excretion in mammals

Rate and extent of oral absorption:	Rapid, no quantitative data Assumed 100%
Rate and extent of dermal absorption*:	Corrosive Assumed 100%
Rate and extent of inhalation absorption:	Corrosive Assumed 100%
Distribution:	Significant, no quantitative data
Potential for accumulation:	no indication of accumulation
Rate and extent of excretion:	Rapid elimination via exhalation of CO ₂ ; low urinary excretion of formic acid
Toxicologically significant metabolite(s)	none

* the dermal absorption value is applicable for the active substance and might not be usable in product authorization

Acute toxicity

Rat LD ₅₀ oral	730 mg/kg bw ¹⁷ Classification as Acute tox cat. 4 (oral) is warranted; H302.
Rat LD ₅₀ dermal	No data for Formic Acid Sodium formate: LD ₅₀ >2000 mg/kg bw
Rat LC ₅₀ inhalation	7.4 mg/l Classification as Acute tox cat. 3 (inhalation) is warranted; H331.

Skin corrosion/irritation

Formic Acid is classified as Skin Corr 1A, H314 (harmonised classification)
Formic acid solutions ≥ 2% are considered skin irritants

¹⁷ RAC agreed in June 2022 on the classification and labelling for formic acid according to Regulation (EC) No 1272/2008 : H302 (& H331) duly confirmed. LD₅₀ values from the adopted RAC opinion that will need to be used in biocidal product authorisation.

Eye irritation

Formic Acid is classified as Skin Corr 1A, H314 (harmonised classification), covering also eye damage/irritation effects
Formic acid solutions $\geq 2\%$ are considered eye irritants

Respiratory tract irritation

Classification as EUH071 'corrosive to the respiratory tract' is warranted as the substance is classified for inhalation toxicity with corrosivity as the mechanism of toxicity.

Skin sensitisation (test method used and result)

No classification for skin sensitization warranted (Buehler test: no sensitising properties shown)

Respiratory sensitisation (test method used and result)

There is no indication that formic acid would be a respiratory sensitizer.

Repeated dose toxicity**Short term**

Species / target / critical effect

No data available on short-term toxicity
Covered by subchronic toxicity studies

Relevant oral NOAEL / LOAEL

No oral repeated dose study available

Relevant dermal NOAEL / LOAEL

No dermal repeated dose study available

Relevant inhalation NOAEL / LOAEL

No inhalation repeated dose study available

Subchronic

Species/ target / critical effect

Rat, pig (oral), rat, mouse (inhal)
local: histological changes in stomach (rat, pig) and upper respiratory tract (rat, mouse)
syst: decreased body weight gain (rat, oral & mouse, inhalation)

Relevant oral NOAEL / LOAEL

As formate:
Rat LOAEL_{syst} 2100 mg/kg bw/d
NOAEL_{syst} 840 mg/kg bw/d
LOAEL_{local} 420 mg/kg bw/d
NOAEL_{local} <420 mg/kg bw/d
Pig LOAEL_{syst} >760 mg/kg bw/d
NOAEL_{syst} 760 mg/kg bw/d
LOAEL_{local} 149 mg/kg bw/d
NOAEL_{local} <149 mg/kg bw/d

Relevant dermal NOAEL / LOAEL

No dermal repeated dose study available

Relevant inhalation NOAEL / LOAEL

Rat	LOAEC _{syst} not achieved NOAEC _{syst} 244 mg/m ³ LOAEC _{local} 61 mg/m ³ NOAEC _{local} 30 mg/m ³
Mouse	LOAEC _{syst} 244 mg/m ³ NOAEC _{syst} 122 mg/m ³ LOAEC _{local} 122 mg/m ³ NOAEC _{local} 61 mg/m ³
overall NOAEC _{local} 60 mg/m ³ (histopathological changes in nasal region of rats and mice at 122 mg/m ³)	

Long term

Species/ target / critical effect

Rat, pig (oral)
local: histological changes in stomach & GI (rat)
syst: decreased body weight gain (rat)

Relevant oral NOAEL / LOAEL

As formate:	
Rat	LOAEL _{syst} 1400 mg/kg bw/d NOAEL _{syst} 280 mg/kg bw/d LOAEL _{local} 280 mg/kg bw/d NOAEL _{local} 35 mg/kg bw/d
Pig	NOAEL _{syst} 301 mg/kg bw/d

Relevant dermal NOAEL / LOAEL

No dermal repeated dose study available

Relevant inhalation NOAEL / LOAEL

No inhalation repeated dose study available

Genotoxicity

Formic acid gave negative results in the <i>in vitro</i> gene mutation study in bacteria, the <i>in vitro</i> cytogenicity study in mammalian cells, and <i>in vitro</i> gene mutation assay in mammalian cells. Chromosome aberrations were observed; it was concluded that formic acid is not itself clastogenic but that the acidic conditions of the medium were responsible for the chromosome aberrations. No <i>in vivo</i> genotoxicity studies are warranted. Formic acid has no genotoxic potential.

Carcinogenicity

Species/type of tumour	Rat, mouse: no evidence of a tumorigenic effect in the stomach or any other tissue was found. Mouse: a higher incidence of primary lung tumours, bronchiolo-alveolar adenomas and carcinomas was not of toxicological relevance.
Relevant NOAEL/LOAEL	As formate: Rat LOAEL _{local} 280 mg/kg bw/d NOAEL _{local} 35 mg/kg bw/d LOAEL _{syst} 1400 mg/kg bw/d NOAEL _{syst} 280 mg/kg bw/d Mouse LOAEL _{local/syst} 1400 mg/kg bw/d NOAEL _{local/syst} 280 mg/kg bw/d

Reproductive toxicity**Developmental toxicity**

Species/ Developmental target / critical effect	Rat, rabbit Formate: no developmental toxicity and teratogenicity observed
Relevant maternal NOAEL	As formate: Rat NOAEL 640 mg/kg bw/d Rabbit NOAEL 670 mg/kg bw/d
Relevant developmental NOAEL	As formate: Rat NOAEL 640 mg/kg bw/d Rabbit NOAEL 670 mg/kg bw/d

Fertility

Species/critical effect	Rat Formate: no adverse effects on fertility observed
Relevant parental NOAEL	As formate: NOAEL 200 mg/kg bw/d
Relevant offspring NOAEL	As formate: NOAEL 670 mg/kg bw/d
Relevant fertility NOAEL	As formate: NOAEL 670 mg/kg bw/d

Neurotoxicity

Species/ target/critical effect

Formic acid is associated with optical nerve and photoreceptor toxicity at high doses. However, adverse effects on the optical nerve and photoreceptors are considered to be an exclusive sequel of acute methanol intoxication in primates. Classification of formic acid as neurotoxic is not warranted.

Developmental Neurotoxicity

Species/ target/critical effect

No evidence of a neurotoxic effect is found in developmental toxicity studies.

Immunotoxicity

Species/ target/critical effect

No immunotoxicity studies available
There is no evidence from skin sensitisation, repeated dose or reproduction toxicity studies, that formic acid may have immunotoxic properties.

Developmental Immunotoxicity

Species/ target/critical effect

No developmental immunotoxicity studies available

Other toxicological studies

None available

Medical/human data

Human data are available from health records from industry and from clinical case reports (accidental or suicidal).

Oral exposure

Due to the corrosivity of formic acid, local effects must be expected at all dose levels. The amount ingested and the concentration determine the grade and the location of the effects. Therefore, the observations range from moderate burns around the mouth to severe corrosion of the gastro-intestinal tract with destruction of the esophagus, perforation of the stomach, and corrosion of the small intestine together with massive bleeding and systemic toxicity (Systemic toxicity observed after ingestion of 30 g formic acid or more).

Accidental and suicidal oral exposure records report reversible burns of the oesophagus after ingestion of small quantities (up to 10g). Consumption of between 5 and 30 g of formic acid led to minor superficial oropharyngeal burns or more severe symptoms including abdominal pain, vomiting, dyspnea and dysphagia, hematemesis and pneumonitis, and esophageal strictures. Doses up to 45 g formic acid were survived by most patients. The majority of patients died after doses between 45 – 200 g formic acid. Reported symptoms at high doses were corrosion of the gastro-intestinal tract, metabolic acidosis, haemolysis, loss of blood pressure, massive bleeding, hepatic and renal failure, and death.

Dermal exposure

Due to the corrosivity of concentrated formic acid, local effects must be expected following contact to the skin and to the eyes. Local burns heal only slowly. Tissue destruction of the skin may result in scarring. Systemic effects may result after contact of concentrated formic acid to extended areas of the body surface. Occupational and accidental dermal exposure records report skin corrosion and metabolic acidosis.

Inhalation

Systemic effects are unlikely to occur. Workplace measurements showed mean values and 95% percentiles far below the threshold limit of 5 ppm or 9.5 mg/m³. Uptake of formic acid at this threshold exposure concentration equals approx. 0.5% of the metabolic rate observed in non-human primates. Therefore, an effect on the blood pH is unlikely. Formic acid inhalation concentrations from 30 ppm onwards are regarded as being immediately dangerous to life and health.

One accidental inhalation exposure record reported reversible Pulmonary dysfunction in the form of Reactive Airway Dysfunction Syndrome. Suicidal inhalation exposure records (mixing of formic acid with concentrated sulphuric acid to form carbon monoxide) report death due to CO intoxication alongside corrosion/irritation of skin, trachea, lungs, stomach due to formic acid fumes.

Summary

	Value	Study	Safety factor
AEL _{short-term}	8.4 mg/kg bw/d	Subchronic 90 day feeding study, rat	100
AEL _{medium-term}	8.4 mg/kg bw/d	Subchronic 90 day feeding study, rat	100

AEL _{long-term}	2.8 mg/kg bw/d rounded to 3 mg/kg bw/d ¹⁸	Chronic 2-year feeding study, rat	100
ADI ¹⁹	3 mg/kg bw/d	EU SANCO D3/AS D, 2005; JECFA, 2003	
ARfD	not required		
Occupational exposure limit	5 ppm or 9.5 mg/m ³ 5 ppm or 9 mg/m ³	EU WEL, MAK/TLV (8-hour TWA) IOELV (Commission Directive 2006/15/EC)	
AEC _{resp tract irrit}	6 mg/m ³	Subchronic 13w inhalation study, rat/mice	10

MRLs

Relevant commodities

default MRL acc to Art.18(1)(b) Reg 396/2005

Reference value for groundwater

According to BPR Annex VI, point 68

N/A

Dermal absorptionStudy (*in vitro/vivo*), species tested

None, corrosive substance

Formulation (formulation type and including concentration(s) tested, vehicle)

N.A.

Dermal absorption values used in risk assessment

100%

Acceptable exposure scenarios (including method of calculation)

Formulation of biocidal product

Not evaluated

Intended uses

Professional uses: Animal house disinfection by fogging, Disinfection of footwear in dipping troughs, Animal feet disinfection and disinfection of transport vehicle tyres

Industrial users

Not evaluated

¹⁸ We refer to TAB entry TOX-4 as the impact of rounding is less than 10%. Please note that for this CAR, the risk characterization has been performed with the non-rounded 2.8 mg formate/kg bw/d value. The decision for rounding the AEL long-term was taken at HH WG I-2022; however it was decided that there was no need to alter the risk characterization of the CAR. For product approval, the rounded 3 mg formate/kg bw/d value should be used.

¹⁹ If residues in food or feed.

Professional users

Animal house disinfection by fogging:

Mixing & loading:

PPE: chemical-resistant gloves, eye/face protection, coveralls, boots;

M&L: appropriate RPE when ventilation is insufficient

RMM: sufficient ventilation

Models used:

Dermal exposure: EUROPOEM II database, liquid manual loading/pouring

Inhalation of vapours: ConsexpoWeb, evaporation, area of release constant

Re-entry: Reuse/restocking of disinfected animal housing:

RMM: sufficient ventilation

Models used:

ConsexpoWeb, exposure to vapour, instantaneous release

OR

RMM approach

Disinfection of footwear in dipping troughs:

Mixing & loading, post-application:

PPE: chemical-resistant gloves, eye/face protection, coveralls, boots;

M&L: appropriate RPE when ventilation is insufficient

RMM: sufficient ventilation

Models used:

Dermal exposure: EUROPOEM II database, Manual loading/pouring volumes up to 20L

Inhalation of vapours: ConsexpoWeb, evaporation, area of release constant

Animal feet disinfection, disinfection of transport vehicle tyres:

Mixing & loading, post-application:

PPE: chemical-resistant gloves, eye/face protection, coveralls, boots;

M&L: appropriate RPE when ventilation is insufficient

RMM: sufficient ventilation

	Models used: Dermal exposure: Mixing & loading Model 4 for M&L and (post)application Inhalation of vapours: ConsexpoWeb, evaporation, area of release constant
Non professional users	Not relevant
General public	Not relevant
Exposure via residue in food	Estimation of dietary exposure of humans to formic acid residues as a consequence of the representative uses was not considered. It is proposed that assessment of dietary risk for humans and livestock be undertaken at biocidal product authorisation.

Chapter 4: Fate and Behaviour in the Environment

Route and rate of degradation in water

Hydrolysis of active substance and relevant metabolites (DT₅₀) (state pH and temperature)

DT₅₀ > 1 year (pH 4, 7 and 9; 49.9±0.5 °C)
DT₅₀ > 20.7 years (pH 7; 12 °C)

Photolytic / photo-oxidative degradation of active substance and resulting relevant metabolites

- Direct photolysis: not expected
- Photo-oxidation with OH-radicals in water: DT₅₀ HCOO⁻ = 35 years

Readily biodegradable (yes/no)

Yes

Inherent biodegradable (yes/no)

-

Biodegradation in freshwater

-

Biodegradation in seawater

-

Non-extractable residues

-

Distribution in water / sediment systems (active substance)

-

Distribution in water / sediment systems (metabolites)

-

Route and rate of degradation in soil

Mineralization (aerobic)

-

Laboratory studies (range or median, with number of measurements, with regression coefficient)

-

DT_{50lab} (20°C, aerobic):

-

DT_{90lab} (20°C, aerobic):

-

BPC-43-2022-06B

DT _{50lab} (10°C, aerobic):	-
DT _{50lab} (20°C, anaerobic):	-
degradation in the saturated zone:	-
Field studies (state location, range or median with number of measurements)	Open literature data suggest DT ₅₀ -values in the range of 1 day for biodegradation of formic acid in soil, even at low temperatures.
DT _{50f} :	1 day (12 °C)
DT _{90f} :	-
Anaerobic degradation	Indication that anaerobic degradation may be possible.
Soil photolysis	-
Non-extractable residues	-
Relevant metabolites - name and/or code, % of applied a.i. (range and maximum)	-
Soil accumulation and plateau concentration	Not relevant due to rapid degradation in soil.

Biodegradation during manure storage

Biodegradation during manure storage	DT ₅₀ ≤ 10.5 days (20 °C) DT ₅₀ ≤ 19.9 days (12 °C)
--------------------------------------	--

Adsorption/desorption

K _a , K _d K _{aoc} , K _{doc} pH dependence (yes / no) (if yes type of dependence)	The K _{oc} for formic acid is pH dependent, with an increasing K _{oc} at increasing pH levels. For risk assessment purposes at a pH of 7, a K _{oc} value of 30 L/kg (log K _{oc} of 1.48) is used.
--	--

Fate and behaviour in air

Direct photolysis in air	-
Quantum yield of direct photolysis	-
Photo-oxidative degradation in air	Latitude: - Season: - DT ₅₀ = 855.7 hours
Volatilization	-

Reference value for groundwater

According to BPR Annex VI, point 68	0.1 µg/L
-------------------------------------	----------

Monitoring data, if available

Soil (indicate location and type of study)	-
--	---

Surface water (indicate location and type of study)	-
Ground water (indicate location and type of study)	-
Air (indicate location and type of study)	-

Chapter 5: Effects on Non-target Species

Toxicity data for aquatic species (most sensitive species of each group) : FRESHWATER

Species	Time-scale	Endpoint	Toxicity
<u>Fish</u>			
<i>Oncorhynchus mykiss</i>	96 h	LC ₅₀	3500 mg/L
<u>Invertebrates</u>			
<i>Daphnia magna</i>	48 h	EC ₅₀	540 mg/L
<i>Daphnia magna</i>	21 d	NOEC	100 mg/L
<u>Algae</u>			
<i>Desmodesmus subspicatus</i>	72 h	E _r C ₅₀	> 1000 mg/L
	72 h	NOE _r C	1000 mg/L
<u>Microorganisms</u>			
Activated sludge	3 h	EC ₅₀	> 500 mg/L

Toxicity data for aquatic species (most sensitive species of each group) : SEAWATER

Species	Time-scale	Endpoint	Toxicity
<u>Fish</u>			
<i>Scophthalmus maximus</i>	96 h	LC ₅₀	1720 mg/L
<u>Invertebrates</u>			
<i>Acartia tonsa</i>	48 h	EC ₅₀	531 mg/L
<u>Algae</u>			
<i>Skeletonema costatum</i>	72 h	E _r C ₅₀	> 1000 mg/L

Effects on earthworms or other soil non-target organisms

Acute toxicity to	-
Reproductive toxicity to	-

Effects on soil micro-organisms

Nitrogen mineralization	-
Carbon mineralization	-

Effects on terrestrial vertebrates

Acute toxicity to mammals	NOAEL _{mammal, oral_chr} = 280 mg/kg _{bw.day}
Acute toxicity to birds	-
Dietary toxicity to birds	-
Reproductive toxicity to birds	-

Effects on honeybees

Acute oral toxicity	-
Acute contact toxicity	-

Effects on other beneficial arthropods

Acute oral toxicity	-
Acute contact toxicity	-
Acute toxicity to	-

Bioconcentration

Bioconcentration factor (BCF)	<ul style="list-style-type: none"> • Estimated BCF_{fish} = 0.00327 L/kg_{wwt} • Estimated BCF_{earthworm} = 0.84 L/kg_{wwt}
Depration time (DT ₅₀)	-
Depration time (DT ₉₀)	-
Level of metabolites (%) in organisms accounting for > 10 % of residues	-

Chapter 6: Other End Points

APPENDIX II: HUMAN EXPOSURE CALCULATIONS

Scenario 1a, fogging, mixing and loading, exposure to vapour, 55% formic acid concentrate - ConsExpo

Substance

Name formic acid

Molecular weight 46g/mol

K_{ow} -2.110Log

Product

Name concentrate 55%

Weight fraction substance 55%

Population

Name worker

Body weight 60kg

Frequency 1per day

Description pt3 M&L in preparation of fogging task/animal housing disinfection

Inhalation

Exposure model Exposure to vapour - Evaporation

Exposure duration 10minute

Product amount 3000g

Weight fraction substance 55%

Room volume 24m³

Ventilation rate 10per hour

Inhalation rate 1.25m³/hr

Application temperature 20°C

Vapour pressure 4.27E+03Pa

Molecular weight 46g/mol

Mass transfer coefficient 10m/hr

Release area mode Constant

Release area 100cm²

Emission duration 5minute

Product in pure form No


Molecular weight matrix 18g/mol

Absorption model Fixed fraction

Absorption fraction 100%

Results

Mean event concentration	3.8mg/m ³
Peak concentration (TWA 15 min)	3.8mg/m ³
Mean concentration on day of exposure	2.6 × 10 ⁻² mg/m ³
Year average concentration	2.6 × 10 ⁻² mg/m ³
External event dose	1.3 × 10 ⁻² mg/kg bw
External dose on day of exposure	1.3 × 10 ⁻² mg/kg bw
Internal event dose	1.3 × 10 ⁻² mg/kg bw
Internal dose on day of exposure	1.3 × 10 ⁻² mg/kg bw/day
Internal year average dose	1.3 × 10 ⁻² mg/kg bw/day

Graph II.1 Formic Acid air concentration during M&L in preparation of fogging task

M&L for fogging
animal housing.xlsx

Scenario 2a, footwear disinfection, mixing and loading, exposure to vapour, 55% formic acid concentrate - ConsExpo

Substance

Name formic acid

Molecular weight 46g/mol

K_{ow} -2.110Log

Product

Name concentrate 55%

Weight fraction substance	55%
Population	
Name	worker
Body weight	60kg
Frequency	104per year
Description	pt3 mixing & loading footbath exposure to vapour

Inhalation


Exposure model	Exposure to vapour - Evaporation
Exposure duration	10minute
Product amount	1090g
Weight fraction substance	55%
Room volume	24m ³
Ventilation rate	10per hour
Inhalation rate	1.25m ³ /hr
Application temperature	20°C
Vapour pressure	4.27E+03Pa
Molecular weight	46g/mol
Mass transfer coefficient	10m/hr
Release area mode	Constant
Release area	100cm ²
Emission duration	5minute
Product in pure form	No
Molecular weight matrix	18g/mol
Absorption model	Fixed fraction
Absorption fraction	100%

Results

Mean event concentration	3.8 mg/m ³
Peak concentration (TWA 15 min)	3.8 mg/m ³
Mean concentration on day of exposure	2.6 × 10 ⁻² mg/m ³
Year average concentration	7.5 × 10 ⁻³ mg/m ³
External event dose	1.3 × 10 ⁻² mg/kg bw
External dose on day of exposure	1.3 × 10 ⁻² mg/kg bw
Internal event dose	1.3 × 10 ⁻² mg/kg bw
Internal dose on day of exposure	1.3 × 10 ⁻² mg/kg bw/day
Internal year average dose	3.8 × 10 ⁻³ mg/kg bw/day

Graph II.2 Formic Acid air concentration during M&L for footwear disinfection

	55% stock solution to be diluted
--	---



Scenario 3a, animal feet disinfection, mixing and loading, exposure to vapour, 55% formic acid concentrate - ConsExpo

Substance	
Name	formic acid
Molecular weight	46g/mol
Kow	-2.110Log
Product	
Name	55% concentrate
Weight fraction substance	55%
Population	
Name	worker
Body weight	60kg

Frequency	1per day
Description	pt3 hoof disinfection vapour inhalation during M&L

Exposure model	Exposure to vapour - Evaporation
Exposure duration	5minute
Product amount	110000g
Weight fraction substance	55%

Room volume	24m ³
Ventilation rate	10per hour
Inhalation rate	1.25m ³ /hr
Application temperature	20°C
Vapour pressure	4.27E+03Pa
Molecular weight	46g/mol
Mass transfer coefficient	10 m/hr
Release area mode	Constant
Release area	100cm ²
Emission duration	5minute
Product in pure form	No
Molecular weight matrix	18g/mol
Absorption model	Fixed fraction
Absorption fraction	100%

Results

Mean event concentration	3.5 mg/m ³
Peak concentration (TWA 15 min)	3.5 mg/m ³
Mean concentration on day of exposure	1.2 × 10 ⁻² mg/m ³
Year average concentration	1.2 × 10 ⁻² mg/m ³
External event dose	6.1 × 10 ⁻³ mg/kg bw
External dose on day of exposure	6.1 × 10 ⁻³ mg/kg bw
Internal event dose	6.1 × 10 ⁻³ mg/kg bw
Internal dose on day of exposure	6.1 × 10 ⁻³ mg/kg bw/day
Internal year average dose	6.1 × 10 ⁻³ mg/kg bw/day

Scenario 3b, animal feet disinfection, application, exposure to vapour, 5% formic acid dilute - ConsExpo

Substance	
Name	formic acid
Molecular weight	46g/mol
K _{ow}	-2.110Log
Product	
Name	5% dilution
Weight fraction substance	5%
Population	
Name	worker
Body weight	60kg
Frequency	1per day
Description	pt3 hoof disinfection exposure to vapour during application

Exposure model	Exposure to vapour - Evaporation
Exposure duration	35minute
Product amount	1000000g
Weight fraction substance	5%
Room volume	9630m ³
Ventilation rate	2per hour
Inhalation rate	1.25m ³ /hr
Application temperature	20°C
Vapour pressure	4.27E+03Pa
Molecular weight	46g/mol
Mass transfer coefficient	10 m/hr
Release area mode	Constant
Release area	3m ²
Emission duration	35minute
Product in pure form	No
Molecular weight matrix	18g/mol
Absorption model	Fixed fraction
Absorption fraction	1

Results

	In-use dilution 1.1% FA
Mean event concentration	1.0 mg/m ³
Peak concentration (TWA 15 min)	1.5 mg/m ³
Mean concentration on day of exposure	2.5 × 10 ⁻² mg/m ³
Year average concentration	2.5 × 10 ⁻² mg/m ³
External event dose	1.3 × 10 ⁻² mg/kg bw
External dose on day of exposure	1.3 × 10 ⁻² mg/kg bw
Internal event dose	1.3 × 10 ⁻² mg/kg bw
Internal dose on day of exposure	1.3 × 10 ⁻² mg/kg bw/day
Internal year average dose	1.3 × 10 ⁻² mg/kg bw/day

Scenario 3c, animal feet disinfection, post-application, exposure to vapour, 5% formic acid dilute - ConsExpo

Substance	
Name	formic acid
Molecular weight	46g/mol
K _{ow}	-2.110Log
Product	
Name	5% dilution
Weight fraction substance	5%
Population	
Name	worker

Body weight 60kg

Frequency 1per day

Description pt3 hoof disinfection exposure to vapour during application

Exposure model	Exposure to vapour - Evaporation
Exposure duration	10minute
Product amount	1000000g
Weight fraction substance	1.1%-3%
Room volume	9630m ³
Ventilation rate	2per hour
Inhalation rate	1.25m ³ /hr
Application temperature	20°C
Vapour pressure	4.27E+03Pa
Molecular weight	46g/mol
Mass transfer coefficient	10 m/hr
Release area mode	Constant
Release area	3m ²
Emission duration	10minute
Product in pure form	No
Molecular weight matrix	18g/mol
Absorption model	Fixed fraction
Absorption fraction	1


Results

	In-use dilution 5% FA
Mean event concentration	3.8 × 10 ⁻¹ mg/m ³
Peak concentration (TWA 15 min)	3.8 × 10 ⁻¹ mg/m ³
Mean concentration on day of exposure	2.6 × 10 ⁻³ mg/m ³
Year average concentration	2.6 × 10 ⁻³ mg/m ³
External event dose	1.3 × 10 ⁻³ mg/kg bw
External dose on day of exposure	1.3 × 10 ⁻³ mg/kg bw
Internal event dose	1.3 × 10 ⁻³ mg/kg bw
Internal dose on day of exposure	1.3 × 10 ⁻³ mg/kg bw/day
Internal year average dose	1.3 × 10 ⁻³ mg/kg bw/day

Graph II.3 Formic Acid air concentration during preparation and use of animal feet bath

55 % stock solution – M&L

5 % in-use solution - application



Scenario 4, exposure during restocking of an area fogged with formic acid solution, ConsExpo

Substance

Name

FA

Molecular weight

46g/mol

K_{ow}

-2.110Log

Product

Name

5,5% dil/19% dil

Weight fraction substance

5,5% / 19%

Population

Name

farm hand

Body weight 60kg

Scenario *ventilation instantaneous release*


Frequency 12 per year


Inhalation

Exposure model	Exposure to vapour - Instantaneous release
Exposure duration	180minute
Product amount	28000g / 6400g
Weight fraction substance	5.5% /19%
Room volume	2810m ³
Ventilation rate	2 per hour / 10 per hour
Inhalation rate	1.25m ³ /hr
Limit concentration to saturated air concentration	Yes
Application temperature	20°C
Vapour pressure	4.27E+03Pa
Molecular weight	46g/mol
Absorption model	Fixed fraction
Absorption fraction	1

Results

Graph II.4 Formic Acid air concentration – re-entry for restocking fogged animal housing





PT3 restocking fog5.5
ventilation time at ven



PT3 restocking fog5.5
ventilation time at ven



PT3 restocking fog19
ventilation time at ven



PT3 restocking fog19
ventilation time at ven

Tentative approach to 'Guidance on the BPR V III HH-Assessment & Evaluation, Section 6: Guidance On Estimating Livestock Exposure to Active Substances used in Biocidal Products'

eCA BE proposes that the assessment of dietary risk for humans and livestock resulting from use of formic acid in PT3 biocidal products be undertaken at biocidal product authorisation. However, a preliminary estimate of this dietary risk is presented here.

Exposure as a consequence of fogging treatment of animal housing:

Considering the volatility of Formic Acid and required waiting times after treatment of animal housing, eCA BE proposes to limit this risk assessment to the inhalation route of exposure.

Saturated vapour concentration of formic acid:

$$\text{SVC} = (\text{vapour pressure} \times \text{MW}) / (\text{gas constant} \times \text{temperature in } ^\circ\text{K})$$

$$= (4271 \text{ Pa} \times 46 \text{ g/mol}) / (8.31451 \text{ J/K mol} \times 293^\circ\text{K})$$

$$= 80.65 \text{ g a.s./m}^3 = 8.065 \times 10^4 \text{ mg a.s./m}^3$$

The high vapour pressure of Formic Acid implies that the trigger value of 0.004 mg/a.s./kg bw/d will be greatly exceeded for inhalation exposure alone, and this for

all animal species considered in the draft Guidance. A Tier II refinement would then be needed. Looking at the alveolar ventilation rate to body weight ratio, the highest and lowest inhalation exposure will be calculated for slaughter goat and turkey, respectively:

Animal species	Bw (kg)	Alveolar ventilation rate (m ³ /d)	Inhalation exposure (mg a.s./kg bw/d)
Slaughter goat	13	3	18612
turkey	7	0.6	6913

If we consider as a refinement that ventilation is used as a risk mitigation measure for workers restocking animal housings, then animals would be exposed to max 6 mg a.s./m³ (AEC for respiratory tract irritation). This would lead to the following values for inhalation exposure:

Animal species	Bw (kg)	Alveolar ventilation rate (m ³ /d)	Inhalation exposure (mg a.s./kg bw/d)
Slaughter goat	13	3	1.38
turkey	7	0.6	0.51

Still, the trigger value of 0.004 mg/a.s./kg bw/d is exceeded for inhalation exposure alone.

However, if we consider the maximum percentage of Formic Acid which is considered safe in animal feed and water (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03), then exposure to Formic Acid via feed/drinking water would greatly surpass the amount exposed to via treated animal housing. It could be disputed whether the default trigger value for a Tier II assessment is applicable here.

Animal species	Bw (kg)	Feed (kg/d)	Max F.A. in feed (mg/d)	Water (kg/d)	Max F.A. in water (mg/d)	Total FA intake (mg/kg bw/d)

Slaughter goat	13	0.5	5000	1.3	5200	785
turkey	7	0.5	5000	1	4000	1286

Exposure to formic acid in hoof disinfectant baths:

Oral exposure: not relevant; cattle do not lick their hooves.

Inhalation exposure: negligible; exposure is transient.

Dermal exposure:

Daily passes through the tub = 2

Exposed skin/hoof area = 1590 cm²

Layer of product absorbed: 0.01 cm

Body weight: 650 kg

Product amount in contact with one hoof/skin:

$$0.01 \text{ cm} \times 1590 \text{ cm}^2 = 15.9 \text{ cm}^3 = 0.0159 \text{ L}$$

5% FA in-use dilution:

If 1L diluted product contains 50000 mg a.s. (5%), then 0.0159 L diluted product contains 795 mg a.s.

Assuming each hoof steps into the hoof bath once at each pass through the bath, then the amount of a.s. each animal comes into contact with during one pass equals 4 x 795 mg a.s. = 3180 mg a.s.

$$3180 \text{ mg a.s.} \times 2 \text{ daily passes} / 650 \text{ kg} = 9.78 \text{ mg a.s./kg bw/d}$$

Here also the trigger value is exceeded.

However, if we consider the maximum percentage of Formic Acid which is considered safe in animal feed and water (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03), then exposure to Formic Acid via feed/drinking water would greatly surpass the

amount exposed to via hoof baths. It could be disputed whether the default trigger value for a Tier II assessment is applicable here.

In fact, the main concern for animal health will be the concentration in air of Formic Acid and its classification as a substance corrosive to the respiratory tract.

Animal species	Bw (kg)	Feed (kg/d)	Max F.A. in feed (mg/d)	Water (kg/d)	Max F.A. in water (mg/d)	Total FA intake (mg/kg bw/d)
Dairy cow	650	25	250000	115	460000	1092

To conclude, in multiple scenarios where animals are exposed to Formic acid, the trigger value for a Tier 2 assessment will be greatly exceeded, even when considering inhalation exposure alone. It is proposed that assessment of dietary risk for humans and livestock be refined at biocidal product authorisation.

However, this assessment should take into consideration that Formic acid is an approved feed additive in the EU. Moreover, this use is not expected to increase the human formic acid exposure through the consumption of products obtained from the treated animals (EFSA, 2009, FA_BPR_Ann_II_8_16_01; EFSA, 2014; FA_BPR_Ann_II_8_16_02; EFSA, 2015, FA_BPR_Ann_II_8_16_03), even though exposure of animals to FA as a feed and drinking water additive is considerably higher than exposure to FA as a PT3 biocidal product.

RMM should be put in place in order to avoid exposure to Formic acid concentrations in air which exceed the AEC for respiratory irritation.

APPENDIX III: ENVIRONMENTAL EMISSION (AND EXPOSURE) CALCULATIONS

This appendix contains the following documents :

- Emission and PEC estimation scenario 'Footwear' : manure route;
- Emission and PEC estimation scenario 'Animal feet' : manure route;
- Aquatic PEC calculations scenario 'Footwear' : STP-route;
- Aquatic PEC calculations scenario 'Animal feet' : STP-route;
- Emission and PEC estimation scenario 'Animal housing, fogging': manure route ;
- Calculation routines for degradation in manure (grassland and arable land) for scenario 'Animal housing, fogging';
- Aquatic PEC calculations scenario 'Animal housing, fogging': STP-route;
- PEC calculations aggregated exposure : manure route.

BPC-43-2022-06B

Emission_PEC_PT3_
Footwear_Final_rev/Emission_PEC_PT3_
Animal feet_Final_rePEC_aq_Footwear_r
evApril2022.xlsxPEC_aq_Animal-feet
_revApril2022.xlsxEmission_PEC_PT3_
Animal_housing_FinDegradation_manu
re_routines_grasslarDegradation_manu
re_routines_arable_lPEC_aq_Animal-hou
sing_revApril2022.xl:PEC_aggregated_re
vApril2022.xlsm

APPENDIX IV: LIST OF TERMS AND ABBREVIATIONS

Not relevant

APPENDIX V: OVERALL REFERENCE LIST

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	1994	Annex II.1 - 8.12.3 / BPD ID A6.12.3_01a	Werksärztlicher Dienst, Department of Occupational Medicine, Unveröffentlichte Mitteilung. BASF, Internal information, -. non-GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
██████████	2002	Annex II.1 - 8.12.3 / BPD ID A6.12.3_01b	Werksärztlicher Dienst, Department of Occupational Medicine, Unveröffentlichte Mitteilung. BASF, Internal information, -. non-GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
Altaweel MM et al.	2009	Annex II.1 - 8.8 / FA_BPR_Ann_II_8_8_11	Ocular and Systemic Safety Evaluation of Calcium Formate as a Dietary Supplement. JOURNAL OF OCULAR PHARMACOLOGY AND THERAPEUTICS Volume 25, Number 3, 223-230, -. - / Published	No	Public
Altiparmak UE	2013	Annex II.1 - 8.13.2 / FA_BPR_Ann_II_8_13_5_01	Toxic optic neuropathies. Curr Opin Ophthalmol, 24:534-539, -. - / Published	No	Public
Andreae, M. O. & Merlet, P.	2001	CAR (ED) / -	Emission of trace gases and aerosols from biomass burning. Global Biogeochem. Cy. 15, 955-966, -. - / Published	No	Public
Anonymous	1990	Annex II.1 - 8.12.8 / BPD ID A6.12.8_01b	NIOSH Pocket Guide to Hazardous Chemicals. U.S. Department of Health and Human Services. Washington, D.C., USA, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Anonymous	2007	Annex II.1 - 5.1, 5.2, 5.3 / BPD ID A4.1_01	UV test for the determination of Formic Acid in foodstuffs and other materials. R-Biopharm, Cat. No. 10 979732 035. - / Published	No	Public
Anonymous	2019	Annex II.1 - 10.1 / 190910 FA_Addendum_Water_final sent to BE 2019-09-10	Formic acid: Degradation kinetics in water, Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. FATF, September10_2019. non-GLP / Unpublished	Yes	FATF
Anonymous	2019	Annex II.1 - 10.1, 10.2 / FA_Addendum_Soil_Deg_2019-08-20	Formic acid: Fate and degradability, Soil and Manure, Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. FATF, August20_2019. non-GLP / Unpublished	Yes	FATF
Anonymous	2020	Annex II.1 - 10.1 / FA_Addendum_Manure_Deg_2020-09-07	Formic Acid: Degradability in Manure; Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. FATF, September07_2020. Non-GLP / Unpublished	Yes	FATF
Anonymous	2020	Annex II.1 - 8.9.2 / 20200904_BASF_FA_Inhalation MAK	Compilation on public information on the MAK value of formic acid; FATF, September04_2020. non-GLP / Unpublished	Yes	FATF
Anonymous	2021	Annex II.1 - 8 / 20210117_FA_BASF_ToxicityEndpoints	Formic acid: Toxicity Endpoints (LC ₅₀ acute inhalation, NOAEC local effects in 90-days rat; Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. FATF, January17_2021. Non-GLP / Unpublished	Yes	FATF
Anonymous	2021	Addendum: use of public data as key data / 20210225 FA_Justification_Public data as key info_deg soil manure	Formic acid: Use of information from public literature as key studies: Degradation in soil, Degradation in manure; Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. BASF SE and Kemira OYJ, February25_2021. Non-GLP / Unpublished	Yes	BASF SE, Kemira OYJ

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Anonymus	2021	Addendum: Parameter justification / 20210117_FA_BASF_Justification HHRA Parameters	Formic Acid: Human Health Risk Assessment, Justifications for parameter adaptations; Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. BASF SE, January17_2021. Non-GLP / Unpublished	Yes	BASF SE
Anonymus	2021	Addendum: Parameter justification / 20210630_FA_BASF_Justification_parti al vapour pressure	Formic Acid: Human Health Risk Assessment, Justifications for partial vapour pressure; Addendum to the biocidal active substance registration dossier of formic acid according to biocidal products regulation (EU) No 528/2012. BASF SE, June 30_2021. Non-GLP / Unpublished	Yes	BASF SE
Atkinson R	1989	Annex II.1 - 10.3.2 / BPD ID A7.3.2_01	Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. J. Phys. Chem. Ref. Data, Monograph No. 1, -. - / Published	No	Public
Bakovic M et al.	2015	Annex II.1 - 8.12.2 / FA_BPR_Ann_II_8_12_2_11.pdf	Suicidal chemistry: combined intoxication with carbon monoxide and formic acid. Int J Legal Med; published online; DOI 10.1007/s00414-015-1208-0, -. - / Published	No	Public
██████████	2014	Annex III.1 - 3.9 / FA_BPR_ID_3_9	Dichte und Viskosität von 75 % Ameisensäure in Wasser (Density and viscosity of formic acid 75% in water). BASF SE Process Research & Chemical Engineering, 2014.209.1. non-GLP / Unpublished	Yes	BASF SE
Baziramakenga R and Simard RR	1998	Annex II.1 - 10.1 / -	Low molecular weight aliphatic acid contents of composted manures. J. Environ. Qual. 27, 557-561., -. - / Published	No	Public
██████████	2007	Annex II.1 - 3.11, 4.6, 4.17.1; [BASF: III-B 3.4] , Annex III.1 - 4.17.1 / BPD ID A3_02	Evaluation of physical and chemical properties according to Directive 67/548/EC Annex V. BASF AG, GCT/S-L511. Laboratory study code SIK-Nr. 07/1018. GLP / Unpublished	Yes	FATF
Boeniger MF	1987	Annex II.1 - 8.8 / BPD ID A6.2_09	Formate in urine as a biological indicator of formaldehyde exposure: a review . Am. Ind. Hyg. Assoc. J. 48(11), 900-908, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Bouchard M, Brunet RC, Droz P-O, Carrier G	2001	Annex II.1 - 8.8 / BPD ID A6.2_03	A biologically based dynamic model for predicting the disposition of methanol and its metabolites in animals and humans . Toxicol. Sci. 64, 169-184, -. - / Published	No	Public
██████████	2007	Annex III.1 - 3.6.2 / BPD ID B3_05	Protectol FM 85 (85% Formic acid) - Compatibility with other products. BASF plc - Biocides Development, --. non-GLP / Unpublished	Yes	BASF SE
Buxton GV, Greenstock CL, Helman WP, Ross AB	1988	Annex II.1 - 10.1.1.1.b / BPD ID A7.1.1.1.2_01	Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxy radicals (.OH/.O-) in aqueous solution. J. Phys. Chem Data 17(2), 513-882, -. - / Published	No	Public
Chameides, W. L. & Davis, D. D.	1983	Annex II.1 - 10.1 / -	Aqueous-phase source of formic acid in clouds. Nature 304, 427-429, -. - / Published	No	Public
Chan TC, Williams SR, and Clark RF	1995	Annex II.1 - 8.12.2 / BPD ID A6.12.2_09	Formic acid skin burns resulting in systemic toxicity.. Annals of Emerg. Medicine 26, 383-386, -. - / Published	No	Public
Chou WL, Speece RE, Siddiqi RH	1979	Annex II.1 - 10.1.3.1.b, Annex II.1 - 10.1.5 / BPD ID A.7.1.2.1.2_01	Acclimation and degradation of petrochemical wastewater components by methane fermentation. Biotechnol. Bioeng. Symp 8. 391-414, -. - / Published	No	Public
Clay KL, Murphy RC, Watkins D	1975	Annex II.1 - 8.8 / BPD ID A6.2_11	Experimental methanol toxicity in the primate: Analysis of metabolic acidosis. Toxicol. Appl. Pharmacol.34, 49-61, -. - / Published	No	Public
██████████	1994	Annex II.1 - 9.1.3.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.3_04	The growth inhibition to Skeletonema costatum of potassium formate liquor. Binnie Environmental Ltd. , ENV340/109410.OUL. GLP / Unpublished	Yes	FATF
██████████	1994	Annex II.1 - 9.1.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex III.1- 9.1 / BPD ID A7.4.1.2_05	The toxicity to Acartia tonsa of potassium formate liquor. Binnie Environmental Ltd. , ENV341/109410.OUL. GLP / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Dalus D et al.	2013	Annex II.1 - 8.12.8 / FA_BPR_Ann_II_8_12_8_03.pdf	FORMIC ACID POISONING IN A TERTIARY CARE CENTER IN SOUTH INDIA: A 2-YEAR RETROSPECTIVE ANALYSIS OF CLINICAL PROFILE AND PREDICTORS OF MORTALITY. The Journal of Emergency Medicine, Vol. 44, No. 2, pp. 373-380, -. - / Published	No	Public
██████████	1998	Annex II.1 - 8.3_03 / BPD ID A6.1.5_02/ FA_BPR_Ann_II_8_3_03	Formi-LHS - Skin sensitisation Study in the Guinea Pig. ██████████, Report No. 1516/22-1032, January 1998 / unpublished.	Yes	BASF (LoA Kemira)
██████████	2007	Annex II.1 - 3.2, 3.4, 3.5, 3.7, 3.1.2, 3.1.3, 3.6, 3.8, 3.13, 3.15, 3.16, 9.1.2, Annex III.1 - 3.1.1, 3.1.2, 3.4.2.2, 3.8, 3.9 / BPD ID A3_01	Spectroscopic characterization and determination of physico-chemical properties of "Formic acid". BASF AG, GKA Competence Center Analytics, 07L00084. GLP / Unpublished	Yes	FATF
██████████	2018	Annex II.1 - 3.2 / 20181112_07L00084 Amendment01 Final Report BPD_ID_A3_01	1st Amendment to final report 'Spectroscopic characterization and determination of physico-chemical properties of "Formic acid"'. BASF SE, November12_2018, GKA Competence Center Analytics, Ludwigshafen Study No. 07L00084. GLP / Unpublished	Yes	FATF
██████████	1992	Annex II.1 - 9.1.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.2_03	The acute toxicity of potassium formate to Daphnia magna. Huntington Research Centre Ltd. (HRC) (sponsored by KSEPL, Rijswijk, NL), SLL 237(f)/920574. GLP / Unpublished	Yes	FATF
██████████	1992a	Annex II.1 - 10.1.1.2.a, Annex II.1 - 10.1.1.2.b, Annex II.1 - 10.1.3.1.a, Annex II.1 - 10.1.3.2.a, Annex II.1 - 10.1.3.2.b, Annex II.1 - 10.1.5, Annex II.1 - 10.2.1, Annex II.1 - 10.2.8, Annex II.1 - 10.2.4, Annex II.1 - 10.2.6, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex II.1 - 9.6 / BPD ID A7.1.1.2.1_03	Assessment of ready biodegradability of potassium formate (Closed Bottle Test). Huntington Research Centre Ltd. (HRC) (sponsored by KSEPL, Rijswijk, NL), SLL 237(a)/920737. GLP / Unpublished	Yes	FATF
██████████	1992b	Annex II.1 - 9.1.3.1, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3 / BPD ID A7.4.1.3_03	Potassium formate - the algistatic activity . Huntington Research Centre Ltd. (HRC) (sponsored by KSEPL, Rijswijk, NL), SLL 237(f)/920647. GLP / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	1992c	Annex II.1 - 9.1.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3 / BPD ID A7.4.1.2_04	The acute toxicity of potassium formate to brown shrimp (<i>Crangon crangon</i>). Huntington Research Centre Ltd. (HRC) (sponsored by KSEPL, Rijswijk, NL), SLL 217(d)/911712. GLP / Unpublished	Yes	FATF
██████████	1992d	Annex II.1 - 9.1.1, Annex II.1 - 9.1.6, Annex II.1 - 9.1.6.1, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3 / BPD ID A7.4.1.1_04	The acute toxicity of potassium formate to juvenile turbot (<i>Scophthalmus maximus</i>).. ██████████ SLL 217(h)/920037. GLP / Unpublished	Yes	FATF
██████████	1992e	Annex II.1 - 9.1.1, Annex II.1 - 9.1.6, Annex II.1 - 9.1.6.1, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex II.1 - 10.1.3.3 / BPD ID A7.4.1.1_03	The acute toxicity of potassium formate to rainbow trout (<i>Oncorhynchus mykiss</i>). ██████████ SLL 217(I)/911691. GLP / Unpublished	Yes	FATF
██████████	1994	Annex II.1 - 10.1.3.3 / BPD ID A7.1.1.2.3_01	The biodegradability in seawater of potassium formate liquor. Binnie Environmental Ltd. (sponsored by OSCA UK Ltd.), ENV342/109410.OUL. GLP / Unpublished	Yes	FATF
██████████	2002	Annex II.1 - 3.2, Annex II.1 - 3.4, Annex II.1 - 3.5, Annex II.1 - 3.7, Annex II.1 - 3.9, Annex II.1 - 3.10, Annex II.1 - 10.1.1.1.a, Annex II.1 - 10.1.4, Annex II.1 - 10.2.4, Annex II.1 - 10.2.6, Annex II.1 - 9.1.7, Annex II.1 - 9.1.7, Annex II.1 - 9.6 / BPD ID A7.1.1.1.1_01	Physico-chemical properties of "Ameisensäure". BASF AG, GKA Analytik, 02L00109. GLP / Unpublished	Yes	FATF
ECT Oekotoxikologie GmbH	2015	Annex II.1 - 3.7.1 / BPD ID A3_11	Henry's Law Constant calculated from water solubility and vapour pressure. ECT Oekotoxikologie GmbH, Flörsheim, Germany, -. non-GLP / Unpublished	Yes	FATF
Eells JT, Henry MM, Lewandowski MF, Seme MT and Murray TG	2000	Annex II.1 - 8.7 / BPD ID A6.10_01	Development and characterization of a rodent model of methanol of methanol-induced retinal and optical nerve toxicity. Neuro Tox 21, 321-330, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
EFSA	2009	Annex II.1 - 8.16, Annex III.1 - 7 / FA_BPR_Ann_II_8_16_01	Scientific Opinion on the safety and efficacy of Formi™ LHS (potassium diformate) as a feed additive for sows. EFSA Journal 2009; 7 (9): 1315, --. non-GLP / Published	No	Public
EFSA	2014	Annex II.1 - 8.16, Annex III.1 - 7 / FA_BPR_Ann_II_8_16_02	Scientific Opinion on the safety and efficacy of formic acid when used as a technological additive for all animal species. EFSA Journal 2014; 12 (10): 3827, --. non-GLP / Published	No	Public
EFSA	2015	Annex II.1 - 8.16 / FA_BPR_Ann_II_8_16_03	Scientific Opinion on the safety and efficacy of formic acid, ammonium formate and sodium formate as feed hygiene agents for all animal species. EFSA Journal 13 (5): 4113, -. - / Published	No	Public
██████████	2016	Annex II.1 - 9.1.5 / FA_BPR_Ann_II_9_1_5_01	A study on the respiration inhibition of activated sludge according to OECD Guideline for testing of chemicals No. 209. ECT Oekotoxikologie GmbH, Flörsheim/Main, Germany, 16EM1XA. GLP / Unpublished	Yes	FATF
██████████	2006	Annex II.1 - 8.12.1 / BPD ID A6.12_01	Workplace exposure of Formic acid. BASF AG, -. non-GLP / Unpublished	Yes	FATF
██████████	2002	Annex II.1 - 8.5.3 / BPD ID A6.6.3_01	In vitro gene mutation test with formic acid in CHO cells (HPRT locus assay) . BASF AG, Project No. 50M0102/024017, 27 June 2002. GLP / Unpublished	Yes	FATF
European Commission	2005	Annex II.1 - 8.16.1 / BPD ID A6.15.4_01a	Provisional list of monomers and additives notified to European commission as substances which may be used in the manufacture of plastics or coatings intended to come into contact with foodstuffs. European Commission, Synoptic Docum. (2005.07.25). - / Published	No	Public
Exner M, Herrmann H, Zellner R	1994	Annex II.1 - 10.1.1.1.b / BPD ID A7.1.1.1.2_03	Rate constants for the reactions of the NO ₃ radical with HCOOH/HCOO ⁻ and CH ₃ COOH/CH ₃ COO ⁻ in aqueous solution between 278 and 328 K. J. Atmos Chem. 18, 359 - 378, -. - / Published	No	Public
██████████	2014	Annex III.1 - 4.6 / BPD ID B3.4_01	Prüfbericht: Flammpunkt nach DIN EN ISO 2719 (Study report: Flash point according to DIN EN ISO 2719). BASF SE, SIK 14/1849. non-GLP / Unpublished	Yes	BASF SE

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Franco A, Fu W, Trapp S.	2009	Annex II.1 - 10.1.2 / 100000_Franco, A_Environ. Toxic_2009	Influence of the soil pH on the sorption of ionizable chemicals: modeling advances. Environ Toxicol Chem 28: 458-464, -. - / Published	No	Public
Gabriel R., Schäfer L., Gerlach C., Rausch T. & Kesselmeier, J.	1999	CAR (ED) / -	Factors controlling the emissions of volatile organic acids from leaves of Quercus ilex L. (Holm oak). Atmos. Environ. 33, 1347-1355, -. - / Published	No	Public
Galloway, J. N., Likens, G. E., Keene, W. C. & Miller, J. M.	1982	CAR (ED) / -	The composition of precipitation in remote areas of the world. J. Geophys. Res. 87, 8771-8786, -. - / Published	No	Public
██████████	2007	Annex II.1 - 8.7.3 / BPD ID A6.1.2_01	Natriumformiat (Sodium formate). Acute dermal toxicity study in rats. ██████████, 11A0123/031083. GLP / Unpublished	Yes	FATF
██████████	2002	Annex II.1 - 8.3 / -	Formic acid - Buehler test in Guinea pigs. ██████████, 32H0102/022005. non-GLP / Unpublished	Yes	FATF
Glanville H, Rousk J, Golyshin P, and Jones DL	2012	Annex II.1 - 10.2 / -	Mineralization of low molecular weight carbon substrates in soil solution under laboratory and field conditions. Soil Biology & Biochemistry 48, 88-95., -. - / Published	No	Public
██████████	1998	Annex II.1 - 8.8 / BPD ID A6.2_10	Formi LHS. Pharmacokinetics after oral dosing in pigs. ██████████, report No. 25280. GLP / Unpublished	Yes	FATF
██████████	2006	Annex II.1 - 4.1, 4.13, Annex III.1 - 4.1, 4.13 / BPD ID A3_03	Expert judgement on oxidising and explosive properties of formic acid. BASF AG, GCT/S-L511. non-GLP / Unpublished	Yes	FATF
Greim H	2003	Annex II.1 - 8.12.8 / BPD ID A6.12.8_01a	Formic Acid. Occupational Toxicants Vol. 19, 169-180, -. - / Published	No	Public
Hama T, Handa N	1981	Annex II.1 - 10.1 / -	[English title not available]. Rikusiugaku Zasshi 42: 8-19, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Hanzlik RP et al.	2005	Annex II.1 - 8.8 / FA_BPR_Ann_II_8_8_10.pdf	ABSORPTION AND ELIMINATION OF FORMATE FOLLOWING ORAL ADMINISTRATION OF CALCIUM FORMATE IN FEMALE HUMAN SUBJECTS. DMD 33:282-286, -. - / Published	No	Public
██████████	2016	Annex III.1 - 6.7 / BPR-6.7-06	PH measurements of solutions of Protectol FM 85 in hard water; report date: 05 Apr 2016. BASF Grenzach GmbH, Germany, BIO15-014-EX. non-GLP / Unpublished	Yes	BASF SE
██████████	2004	Annex II.1 - 8.9.1, _8.9.2, _8.9.3, ED-Assessment / BPD ID A6.4.1_02	Formi LHS: Target species safety study in the farrowing pig. ██████████, 1516/034-D6154. GLP / Unpublished	Yes	FATF
Hellstén PP, Kivimäki AL, Miettinen IT, Mäkinen RP, Salminen JM, Nystém TH	2005a	Annex II.1 - 10.1 / -	Degradation of potassium formate in the unsaturated zone of a sandy aquifer.. Journal of Environmental Quality 34(5), 1665-1671., -. - / Published	No	Public
Hellstén PP, Salminen JM, Jørgensen KS, Nystén TH	2005b	Annex II.1 - 10.1 / -	Use of potassium formate in road winter deicing can reduce groundwater deterioration. Environ Sci Technol 39, 5095-5100, -. - / Published	No	Public
██████████	2016a	Annex III.1 - 3.2 / KT_BPR_Ann3_5	Determination of the acidity/alkalinity of Formic acid 85% (also known under the tradename Fennopur MH85) according to CIPAC, MT 191. Laus GmbH, Kirrweiler, Germany, 16011907G975. GLP / Unpublished	Yes	Kemira / Taminco (LoA: BASF SE)
██████████	2016b	Annex III.1 - 3.4.2.3 / KT_BPR_Ann3_8	Determination of the corrosion of metals by Formic acid 85% (also known under the tradename Fennopur MH85) following method 37.4 C.1 of the UN Handbook. Laus GmbH, Kirrweiler, Germany, 16011907G979. GLP / Unpublished	Yes	Kemira / Taminco (LoA: BASF SE)
██████████	2016c	Annex III.1 - 3.2 / KT_BPR_Ann3_12	Determination of the pH-value of Formic acid 85% (also known under the tradename Fennopur MH85) according to CIPAC, MT 75. Laus GmbH, Kirrweiler, Germany, 16011907G907. GLP / Unpublished	Yes	Kemira / Taminco (LoA: BASF SE)

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	1997	Annex II.1 - 8.8 / BPD ID A6.2_01	The chemical behavior of Potassium Diformate in water solutions. Comparison with Formic Acid.. Hydro Research Centre Porsgrunn, Norway, 97B_AO5.SAM. non-GLP / Unpublished	Yes	FATF
HSDB	2006	Annex II.1 - 10.1.1.1.b, Annex II.1 - 10.3.2, ED-Assessment / BPD ID A7.1.1.1.2_04	Database extract. TOXNET, -. - / Published	No	Public
Iannotti EL, Porter JH, Fischer JR, and Sievers M	1997	Annex II.1 - 10.2 / -	Changes in swine manure during anaerobic digestion. In: Developments in industrial microbiology. Vol. 20: Proceedings of the 35th general meeting of the Society for Industrial Microbiology held at Houston , Texas: August 14-18, 1978. Arlington, VA, USA. Chapter 49, pp. 519-529, -. - / Published	No	Public
Jager T	1998	Annex II.1 - 9.6 / FA_BPR_Ann_II_9_6	Mechanistic approach for estimating bioconcentration of organic chemicals in earthworms (Oligochaeta). Environmental Toxicology and Chemistry 17(10), 2080-2090. Cited in ECHA (2017). Guidance on Biocidal Products Regulation: Volume IV Environment - Assessment and Evaluation (Parts B+C). DoI 10.2823/033935.	No	Public
██████████	1988	Annex II.1 - 9.1.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.2_01	Determination of the acute toxicity of formic acid to the waterflea Daphnia magna Straus. BASF AG, Department of Ecology, 1/0290/2/88-0290/88. non-GLP / Unpublished	Yes	FATF
JECFA	2003	Annex II.1 - 8.16.1 / BPD ID A6.15.4_01b	Formic acid. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives. JECFA, -. - / Published	No	Public
Jefferys DB and Wiseman HM	1980	Annex II.1 - 8.12.2 / BPD ID A6.12.2_05	Formic acid poisoning. Postgrad. Med. J. 56, 761-763, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	2000	Annex II.1 - 10.1.1.2.a, Annex II.1 - 10.1.1.2.b, Annex II.1 - 10.1.3.1.a, Annex II.1 - 10.1.3.2.a, Annex II.1 - 10.1.3.2.b, Annex II.1 - 10.1.5, Annex II.1 - 10.2.1, Annex II.1 - 10.2.8, Annex II.1 - 10.2.4, Annex II.1 - 10.2.6, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex II.1 - 9.6, Annex III.1- 10 / BPD ID A7.1.1.2.1_04	Biodegradability of potassium formate in water tested with OECD 310D (Closed Bottle Test). Norwegian Institute for Water Research (NIVA) (sponsored by Norsk Hydro ASA, Porsgrunn, Norway), B387/1. non-GLP / Unpublished	Yes	FATF
██████████	2022	Annex II.1 - 8.5.1_02 / BPD ID A6.6.1_02 / FA_BPR_Ann_II_8_5_1_02	Salmonella typhimurium / Escherichia coli reverse mutation assay. BASF SE. 40M0247/14M172	Yes	FATF
Kavet R and Naus KM	1990	Annex II.1 - 8.8 / BPD ID A6.2_12	The toxicity of inhaled methanol vapors . Crit. Rev. Toxicol. 21, 21-50, -. - / Published	No	Public
Kawamura, K., Ng., L. L. & Kaplan, I. R.	1985	CAR (ED) / -	Determination of organic acids (C1-C10) in the atmosphere, motor exhausts and engine oils.. Environ. Sci. Tech. 19, 1082-1086, -. - / Published	No	Public
██████████	2013	Annex II.1 - 5.1, _5.2, _5.3 / BPD ID A4.1_03	Validation of an enzymatic method for the determination of formic acid.. Institute Dr. Appelt, Mannheim, Germany, No. 001. non-GLP / Unpublished	Yes	FATF
██████████	1989	Annex II.1 - 9.1.1, Annex II.1 - 9.1.6, Annex II.1 - 9.1.6.1, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3 , Annex III.1- 9.1 / BPD ID A7.4.1.1_01	Report on the study of the acute toxicity to golden orfe (Leuciscus idus L., golden variety) (in German ██████████, 10F0218/885243. non-GLP / Unpublished	Yes	FATF
██████████	2003	Annex II.1 - 8.9.4, ED-Assessment / BPD ID A6.5_02	Effect of pre-mating administration of Formi LHS on ovulation/fertility of breeding sows. ██████████, Project No. 818 545M (F-446). non-GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
██████████	2017	Annex II.1 - 4.16 / KT_BPR_Ann2_13	Determination of the corrosion of metals by Formic acid 99% following method 37.4 C.1 of the UN Handbook. Laus GmbH, Kirrweiler, Germany, 16092902G979. - / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Lamarre et al.	2013	CAR (ED) / -	Formate: essential metabolite, a biomarker or more?. Clin Chem Lab Med 51(3):571-578, -. - / Published	No	Public
██████████	2017	Annex II.1 - 5.1 / FA_BPR_Ann_II_5_1_Analytics_methods_active_substance.pdf	Formic acid. Validation of Analytical Methods for the Determination of the Active Substance and Water. Noack Laboratorien GmbH, Sarstedt. Germany, 16091BE/CMV 177788. GLP / Unpublished	Yes	BASF SE / Taminco BV
██████████	2017	Annex II.1 - 5.1 / Formic acid. Validation of Analytical Methods for the Determination of the Active Substance and Water.	Noack Laboratorien GmbH, Sarstedt. Germany. FA_BPR_Ann_II_5_1_Analytics_methods_active_substance.pdf, 16091BE/CMV 177788. GLP / Unpublished	Yes	BASF SE, Taminco bvba
Lin PT and Dunn WA	2014	Annex II.1 - 8.12.2 / FA_BPR_Ann_II_8_12_2_12.pdf	Suicidal carbon monoxide poisoning by combining formic acid and sulfuric acid with a confined space. J Forensic Sci, January 2014, Vol. 59, No. 1, -. - / Published	No	Public
Lissner H, Wehrer M, Jartun M, Totsche KU	2014	Addendum: Fate and degradability: Soil and Manure / -	Degradation of deicing chemicals affects the natural redox system in airfield soils.. Environ Sci Pollut Res 21, 9036-9053. , -. - / Published	No	Public
Makar AB, Tephly TR, Sahin G, Osweiler G	1990	Annex II.1 - 8.8 / BPD ID A6.2_08	Formate metabolism in young swine. Toxicol. Appl. Pharmacol. 105, 315-320, -. - / Published	No	Public
Malizia E, Reale C, Pietropaoli P, and De Ritis GC	1977	Annex II.1 - 8.12.2, Annex II.1 - 8.12.2 / BPD ID A6.12.2_07a	Formic acid intoxications. Acta Pharm. Toxi.,S41342-347, -. - / Published	No	Public
Malorny G	1969a	Annex II.1 - 8.8, Annex II.1 - 8.7, ED-Assessment / BPD ID A6.2_06	Die akute und chronische Toxizität der Ameisensäure und ihrer Formiate. Z. Ernährungs-wiss. 9, 332-339, -. - / Published	No	Public
Malorny G	1969b	Annex II.1 - 8.8, Annex II.1 - 8.13.2 / BPD ID A6.2_07	Stoffwechselfersuche mit Natrium-formiat und Ameisensäure beim Menschen. Z. Ernährungs-wiss. 9, 340-348, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Martin-Amat G, McMartin, KE, Hayreh SS, Hayreh MS, Tephly TR	1978	Annex II.1 - 8.8, Annex II.1 - 8.13.2 / BPD ID A6.2_05	Methanol poisoning: Ocular toxicity produced by formate. . Toxicol. Appl. Pharmacol., 45, 201-208, -. - / Published	No	Public
██████████	2006	Annex II.1 - 10.3.2, Annex II.1 - 10.3.1 / BPD ID A7.3.1_01	Formic acid, EPI Suite v.3.12 calculations. BASF AG, Department of Product Safety, -. non-GLP / Unpublished	Yes	FATF
██████████	2016	Annex III.1 - 3.4.1.3 / BPR ID 3.4.1.3_01	Interpretation of DSC-curve of study 02L00109. BASF SE, -. non-GLP / Unpublished	Yes	BASF SE
Morita T, Takeda K, and Okumura K	1990	Annex II.1 - 8.5.2 / BPD ID A6.6.2_01	Evaluation of clastogenicity of formic acid, acetic acid and lactic acid on cultured mammalian cells. Mut Res 240, 195-202, -. - / Published	No	Public
██████████	1999	Annex II.1 - 8.4, _8.1, _8.2 / BPD ID A6.1.6_01	A sensory irritation study with Formi@ LHS in male mice.. ██████████, V98.1244. GLP / Unpublished	Yes	FATF
██████████	2007	Annex II.1 - 5.2.2, Annex II.1 - 5.2.2 / BPD ID A4.1_02	Method for the determination of formic acid in the air.. BASF AG, -. non-GLP / Unpublished	Yes	FATF
██████████	1985	Annex II.1 - 8.7.1 / BPD ID A6.1.1_01	Akute orale Toxizität von Ameisensäure 99 % für Ratten. Chem. ██████████, Report No. 0359. non-GLP / Unpublished	Yes	FATF
Murtaugh JJ, Bunch RL	1965	Annex II.1 - 10.1 / -	Acidic Components of Sewage Effluents and River Water. J Water Pollut Control Fed 37: 410-5, -. - / Published	No	Public
Naik RB, Stephens WP, Wilson DJ, Walker A, and Lee HA	1980	Annex II.1 - 8.12.2 / BPD ID A6.12.2_04	Ingestion of formic acid-containing agents - report of three fatal cases. Postgrad. Med. J. 56, 451-456, -. - / Published	No	Public
Neeb, P., Sauer, F., Horie, O. & Moortgat, G. R.	1997	CAR (ED) / -	Formation of hydroxymethyl hydroperoxide and formic acid in alkene ozonolysis in the presence of water vapor. Atmos. Environ. 31, 1417-1423, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	2007a	Annex II.1 - 3.1.4, 3.12, Annex III.1 - 3.1.3, 3.4.2.3, 4.16 / BPD ID A3_06	Expert Judgement: Formic acid 99-100 % - materials compatibility and odor. BASF AG, E-CZS/PC. non-GLP / Unpublished	Yes	FATF
██████████	2007b	Annex III.1 - 3.4.1.2 / BPD ID B3_02	Formic acid 85 % - Determination of storage stability. BASF AG , E-CZS/PC. non-GLP / Unpublished	Yes	BASF SE
██████████	2007c	Annex III.1 - 3.5.1, 3.5.2, 3.5.7 / BPD ID B3_03	Technical characteristics of the biocidal product Protectol FM 85 (formic acid 85 %). BASF AG , E-CZS/PC. non-GLP / Unpublished	Yes	BASF SE
NTP-CERHR expert panel	2004	Annex II.1 - 8.8, Annex II.1 - 8.13.2 / BPD ID A6.2_04	NTP-CERHR expert panel report on the reproductive and developmental toxicity of methanol. U.S. DHHS, NTP; Reprod. Toxicol. 18: 303-390, -. - / Published	No	Public
OECD	2007	Annex II.1 - 9.1 / BPD ID IIA4.2.1_01	SIDS Initial Assessment Report on the Ammonia Category. OECD, Paris, -. - / Published	Yes	FATF
Page LH, Ni JQ, Heber AJ, Mosier NS, Liu X, Joo HS, Ndegwa PM, Harrison JH	2014	Annex II.1 - 10.2 / 2014_Page LH et al_manure_anaerobic digestion	Characteristics of volatile fatty acids in stored dairy manure before and after anaerobic digestion. Biosystems Engineering 118: 16-28, -. - / Published	No	Public
██████████	1988a	Annex II.1 - 10.1.1.2.a, Annex II.1 - 10.1.1.2.b, Annex II.1 - 10.1.3.1.a, Annex II.1 - 10.1.3.2.a, Annex II.1 - 10.1.3.2.b, Annex II.1 - 10.1.5, Annex II.1 - 10.2.1, Annex II.1 - 10.2.8, Annex II.1 - 10.2.4, Annex II.1 - 10.2.6, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex II.1 - 9.6, Annex III.1- 10 / BPD ID A7.1.1.2.1_01	Report on the determination of the biological degradability of formic acid in the Modified OECD Screening Test. BASF AG, Lab. of Environm. Analytics & Ecology, 0048/88. non-GLP / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	1988b	Annex II.1 - 10.1.1.2.a, Annex II.1 - 10.1.1.2.b, Annex II.1 - 10.1.3.1.a, Annex II.1 - 10.1.3.2.a, Annex II.1 - 10.1.3.2.b, Annex II.1 - 10.1.5, Annex II.1 - 10.2.1, Annex II.1 - 10.2.8, Annex II.1 - 10.2.4, Annex II.1 - 10.2.6, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex II.1 - 9.6, Annex III.1- 10 / BPD ID A7.1.1.2.1_02	Report on the determination of the biological degradability of formic acid in the Modified OECD Screening Test. BASF AG, Lab. of Environm. Analytics & Ecology, 52/0048/88. non-GLP / Unpublished	Yes	FATF
██████████	1988c	Annex II.1 - 9.1.5, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3 / BPD ID A7.4.1.4_01, Annex III.1- 9.1 / BPD ID A7.4.1.4_01	Report on the Determination of the Respiration Activity of Activated Sludge by Formic Acid in the Short-Term Respiration Inhibition Test. BASF AG, Lab. of Environm. Analytics & Ecology, 01.0048/88. non-GLP / Unpublished	Yes	FATF
Rajan N, Rahim R, and Krishna Kumar S	1985	Annex II.1 - 8.12.2 / BPD ID A6.12.2_03	Formic acid poisoning with suicidal intent: a report of 53 cases. Postgrad. Med. J. 61, 35-36, -. - / Published	No	Public
██████████	1998	Annex II.1 - 8.9.1, _8.9.2, _8.9.3, ED-Assessment / BPD ID A6.4.1_01	Formi LHS: 13 week oral (dietary administration) toxicity study in the rat with a 4 week treatment-free period. ██████████, 1516/6-D6154. non-GLP / Unpublished	Yes	FATF
██████████	2007	Annex II.1 - 3.3 / BPD ID B3_01b	Physico-chemical properties of "Ameisensäure 85%". BASF AG, GKA Competence Center Analytics, 07L00172. GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
██████████	2007	Annex III.1 - 3.2 / BPD ID B3_01b	Physico-chemical properties of "Ameisensäure 85%". BASF AG, GKA Competence Center Analytics, 07L00172. GLP / Unpublished	Yes	BASF SE
██████████	2018	Annex III.1 - 6.7 / 1089285_13697_Version01.pdf	Protectol FM 85 - Quantitative surface test for the evaluation of bactericidal and fungicidal efficacy according to EN 13697 - Version01; report date: 11 May 2018. Labor LS SE & Co. KG, Bad Bocklet, Germany, L+S-No. 180411-0321-001. Guideline Study / Unpublished	Yes	BASF SE

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	2018	Annex III.1 - 6.7 / 1089285_1657_Version01.pdf	Protectol FM 85 - Quantitative suspension test for the evaluation of yeasticidal efficacy according to EN 1657 - Version01; report date: 11 May 2018. Labor LS SE & Co. KG, Bad Bocklet, Germany, L+S-No. 180411-0321-001. Guideline Study / Unpublished	Yes	BASF SE
██████████	2016	Annex III.1 - 6.7 / 0702966_14349_16438_engl_Version01.pdf	Protectol FM 85 - Quantitative surface test for the evaluation of bactericidal efficacy according to EN 14349 and of yeasticidal efficacy according to EN 16438 - Version01; report date: 08 Sep 2016. Labor L+S AG, Bad Bocklet, Germany, L+S-No. 160804-0106-001. Guideline Study / Unpublished	Yes	BASF SE
██████████	2016	Annex III.1 - 6.7 / BPR-6.7-05	Sample Protectol FM 85: Quantitative suspension test for the evaluation of the microbicidal efficacy according to EN 1276 and EN 1650; report date: 24 Mar 2016. Labor L+S AG, Bad Bocklet, Germany, L+S 0543119. non-GLP / Unpublished	Yes	BASF SE
██████████	2008a	Annex II.1 - 8.10.1, ED-Assessment / BPD ID A6.8.1_02	Natriumformiat (sodium formate) - Prenatal developmental toxicity study in Himalayan rabbits. Oral administration (Gavage).. ██████████ 40R0123/03089. GLP / Unpublished	Yes	FATF
██████████	2008b	Annex II.1 - 8.10.2, ED-Assessment / BPD ID A6.8.2_01	Natriumformiat (Sodium formate). Two-Generation Reproduction Toxicity Study in Wistar Rats. Administration via the Diet.. ██████████ 70R0123/03091. GLP / Unpublished	Yes	FATF
██████████	2005	Annex II.1 - 8.10.3, ED-Assessment / BPD ID A6.8.1_01	Sodium formate - Prenatal developmental toxicity study in Wistar rats . ██████████ 30R0123/03036. GLP / Unpublished	Yes	American Chemistry Council/US A
██████████	1988	Annex II.1 - 9.1.3.1 , Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.3_01	Algal growth inhibition test. BASF AG, Department of Ecology, 2/0290/88. non-GLP / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Sigurdsson J, Björnsson A, and Gudmundsson ST	1983	Annex II.1 - 8.12.2 / BPD ID A6.12.2_08	Formic acid burn - local and systemic effects. . Burns 9, 358-361, -. - / Published	No	Public
██████████	2017	Annex II.1 - 3.3 / BASF_BPR_Ann2_1	Acidity or Alkalinity of Protectol FM 99. Eurofins, Niefern-Öschelbronn, EAS Study Code S16-06390. GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
██████████	2017	Annex II.1 - 3.3 / BASF_BPR_Ann2_2	pH of Protectol FM 99 (aqueous dilution). Eurofins, Niefern-Öschelbronn, EAS Study Code S16-06389. GLP / Unpublished	Yes	BASF SE (LoA: Kemira / Taminco)
Spoelstra SF	1979	Annex II.1 - 10.2 / -	Volatile fatty acids in anaerobically stored piggery wastes. Neth. J. agric. Sci. 27, 60-66., -. - / Published	No	Public
Stavroukou, T., Muller, J. F., Peeters, J., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P., Hurtmans, D., De Maziere, M., Vigouroux, C., Deutscher, N., Griffith, D., Jones, N. & Paton-Walsh, C.	2012	CAR (ED) / -	Satellite evidence for a large source of formic acid from boreal and tropical forests. Nature Geoscience, 5 (1), 26-30, -. - / Published	No	Public
Takata Y, Tani M, Kato T, and Koike M	2011	Annex II.1 - 10.2 / -	Effects of land use and long-term organic matter application on low-molecular-weight organic acids in an Andisol. J. Soil Sci. Manage. 2(10), 292-298, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Tete E, Viaud V, and Walter C	2015	Annex II.1 - 10.2 / -	Organic carbon and nitrogen mineralization in a poorly-drained mineral soil under transient waterlogged conditions: an incubation experiment.. European Journal of Soil Science, 66, 427-437., -. - / Published	No	Public
Thompson M	1992	Annex II.1 - 8.9.1, _8.9.2, _8.9.3 / BPD ID A6.4.3_01	NTP Technical Report on Toxicity Studies of Formic Acid. administered by inhalation to F344/N rats and B6C3F1 mice. US Department of Health and Human Services . NTP US DHHS, Toxicity Report Series No: 19, NIH Publ. No: 92-3342, July 1992. - / Published	No	Public
██████████	1991	Annex II.1 - 9.1.5, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.4_02	Bacterial Growth Inhibition Test. BASF AG, Laboratory of Ecology, 9/0290/88. non-GLP / Unpublished	Yes	FATF
Van Hees PAW, Johansson E, and Jones DL	2008	Annex II.1 - 10.2 / -	Dynamics of simple carbon compounds in two forest soils as revealed by soil solution concentrations and biodegradation kinetics.. Plant Soil 310, 11-23., -. - / Published	No	Public
Verstraete AG, Vogelaers DP, van den Bogaerde JF, Colardyn FA, Ackerman CM and Buylaert WA	1989	Annex II.1 - 8.12.2 / BPD ID A6.12.2_02	Formic acid poisoning: Case report and in vitro study of the haemolytic activity. Am J Emerg Med 7, 286-290, -. - / Published	No	Public
von Muehlendahl KE, Oberdisse U and Krienke EG	1978	Annex II.1 - 8.12.2 / BPD ID A6.12.2_06	Local injuries by accidental ingestion of corrosive substances by children. Arch Toxicol 39, 299-314, -. - / Published	No	Public
██████████	2007	Annex II.1 - 9.1.4.1, Annex II.1 - 9.1.7, Annex II.1 - 9.1.7, Annex II.1 - 9.6, Annex III.1- 10.2 / BPD ID A7.4.2_01	Formic acid, BCFWIN v.2.17 calculations. ECT Oekotoxikologie GmbH, Flörsheim, Germany, -. non-GLP / Unpublished	Yes	FATF

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
██████████	2005	Annex II.1 - 9.1.1, Annex II.1 - 9.1.6, Annex II.1 - 9.1.6.1, Annex II.1 - 9.2.1; Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.1_02	Acute toxicity of ammonium formate to zebra fish (<i>Danio rerio</i>). ██████████ KEM-001/4-11. GLP / Unpublished	Yes	FATF
██████████	2005	Annex II.1 - 9.1.2, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID A7.4.1.2_02	Effect of ammonium formate on the immobilization of <i>Daphnia magna</i> . Fraunhofer-IME, KEM-001/4-20. GLP / Unpublished	Yes	FATF
██████████	2005	Annex II.1 - 9.1.3.1, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, Annex III.1- 9.1 / BPD ID FA. A7.4.1.3_02	Effect of ammonium formate on the growth of <i>Pseudokirchneriella subcapitata</i> . Fraunhofer-IME, KEM-001/4-30. GLP / Unpublished	Yes	FATF
Westphal F, Rochholz G, Ritz-Timme S, Bilzer N, Schütz HW, Kaatsch HJ	2001	Annex II.1 - 8.12.2 / BPD ID A6.12.2_01	Fatal intoxication with a decalcifying agent containing formic acid. <i>Int. J. Legal Med.</i> 114, 181-185, -. - / Published	No	Public
██████████	1999	Annex II.1 - 8.9.1, _8.9.2, _8.9.3, ED-Assessment / BPD ID A6.5_03	Formi LHS. Combined chronic toxicity and 104 week oral (dietary administration) oncogenicity study in the rat. Interim Draft study report ██████████ 1516/30-D6154. GLP / Unpublished	Yes	FATF
██████████	2002a	Annex II.1 - 8.9.1, _8.9.2, _8.9.3, Annex II.1 - 8.11.1, ED-Assessment / BPD ID A6.5_01	Formi LHS. Combined chronic toxicity and 104 week oral (dietary administration) oncogenicity study in the rat. ██████████, 1516/30-D6154. GLP / Unpublished	Yes	FATF
██████████	2002b	Annex II.1 - 8.11.2, ED-Assessment / BPD ID A6.7_02	Formi LHS. 80 week oral (dietary administration) oncogenicity study in the mouse. ██████████ 1516/33-D6154. GLP / Unpublished	Yes	FATF
Yang CC et al.	2008	Annex II.1 - 8.12.2 / FA_BPR_Ann_II_8_12_2_13.pdf	Formic acid: A rare but deadly source of carbon monoxide poisoning. <i>Clinical Toxicology</i> , 46:4, 287-289, -. - / Published	No	Public

Author(s)	Year	Section No / Reference No	Title. Source (where different from company) Company, Report No.	Data Protection Claimed (Yes/No)	Owner
Yelon JA, Simpson RL, and Gudjonsson O	1996	Annex II.1 - 8.12.2 / BPD ID A6.12.2_10	Formic acid inhalation injury: a case report.. J. Burn Care Rehab. 17, 241-242., -. - / Published	No	Public
Zeiger E, Anderson B, Haworth S, Lawlor T, and Mortelmans K	1992	Annex II.1 - 8.5.1 / BPD ID A6.6.1_01	Salmonella mutagenicity tests: V. Results from the testing of 311 chemicals. Environ. Molec. Mutagen. 19, Suppl 21, 2-141, -. - / Published	No	Public
██████████	1980	Annex II.1 - 8.7.2 / BPD ID A6.1.3_01	Bestimmung der akuten Inhalationstoxizität LC50 von Ameisensäure als Dampf bei 4-stündiger Exposition an Sprague-Dawley Ratten. ██████████ August 21, 1980, 16 pages Report No. 78/651. non-GLP / Unpublished	Yes	FATF
██████████	1980	Annex II.1 - 8.7.2 / BPD ID A6.1.3_01EN	Complete translation of BPD ID A6.1.3_01 into English (Date of translation: Aug 16, 2007). Acute inhalation toxicity LC50 of formic acid as vapor after 4-hour exposure in Sprague-Dawley rats. ██████████ Report No. 78/651; 16 pages, Non-GLP / Unpublished	Yes	FATF
Zepp RG, Hoigné J, Bader H	1987	Annex II.1 - 10.1.1.1.b / BPD ID A7.1.1.1.2_02	Nitrate-induced photooxidation of trace organic chemicals in water. Environ. Sci. Technol 21, 443-450, -. - / Published	No	Public
██████████	2007	Annex II.1 - 9.1.6, Annex II.1 - 9.1.6.1, Annex II.1 - 9.1.6.2.a, Annex II.1 - 9.2.1, Annex II.1 - 9.2.2, Annex II.1 - 9.2.3, ED-Assessment, Annex III.1- 9.1 / BPD ID A7.4.3.4_03	Final Report: Formic acid – Determination of the chronic effect on the reproduction of the water flea Daphnia magna STRAUS.. BASF AG, Experimental Toxicology and Ecology, 51E0274/073100. GLP / Unpublished	Yes	FATF

APPENDIX VI: CONFIDENTIAL INFORMATION

Please see separate document

APPENDIX VII: HUMAN HEALTH – READ-ACROSS JUSTIFICATION



RAAF final.docx