

Summary of product characteristics for a biocidal product

Product name: KILRAT PLUS rágcsálóirtó granulátum

Product type(s): PT14 - Rodenticides (Pest control)

PT14 - Rodenticides (Pest control)

Authorisation number: HU-2014-MA-14-00097-0000

R4BP 3 asset reference number: HU-0006720-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	1
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	2
4. Authorised use(s)	3
5. General directions for use	8
5.1. Instructions for use	8
5.2. Risk mitigation measures	10
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	10
5.4. Instructions for safe disposal of the product and its packaging	10
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	11
6. Other information	11

Administrative information

1.1. Trade names of the product

KILRAT PLUS rágcsálóirtó granulátum

1.2. Authorisation holder

Name and address of the authorisation holder	Name	Bábolna Bioenvironmental Centre Ltd.
	Address	Szállás u. 6. H-1107 Budapest Hungary
Authorisation number	HU-2014-MA-14-00097-0000	
R4BP 3 asset reference number	HU-0006720-0000	
Date of the authorisation	30/01/2014	
Expiry date of the authorisation	31/12/2024	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	Bábolna Környezetbiológiai Központ Kft.
Address of the manufacturer	Szállás u. 6. 1107 Budapest Hungary
Location of manufacturing sites	Köves János út 1-3. 2943 Bábolna Hungary

1.4. Manufacturer(s) of the active substance(s)

Active substance	11 - Brodifacoum
Name of the manufacturer	Dr. Tezza Srl
Address of the manufacturer	Via Tre Ponti 22 37050 S.Maria di Zevio Italy
Location of manufacturing sites	Via Tre Ponti 22 37050 S. Maria di Zevio Italy

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	Active Substance	56073-10-0	259-980-5	0,005

2.2. Type of formulation

GR - Granule

3. Hazard and precautionary statements

Hazard statements	<p>May damage the unborn child.</p> <p>May cause damage to organs ; through prolonged or repeated exposure (blood).</p>
Precautionary statements	<p>Keep out of reach of children.</p> <p>Do not handle until all safety precautions have been read and understood.</p> <p>Do not breathe dust.</p> <p>Avoid release to the environment.</p> <p>Wear protective gloves.</p> <p>IF exposed or concerned: Get medical advice.</p>

Dispose of contents to in accordance with local government.

4. Authorised use(s)

4.1 Use description

Use 1 - House mice and rats - trained professionals - indoor

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: -juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: Brown rat Development stage: -juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: Roof rat Development stage: -juveniles and adults
Field(s) of use	Indoor Indoor
Application method(s)	Method: Bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequencies	Application Rate: - Dilution (%): - Number and timing of application: <u>For mice:</u> 40 g / bait station. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters (for high levels of infestation) to 10 meters (for low levels of infestation). <u>For rat:</u> 100 g / bait station. If more than one bait station is needed, the minimum distance between bait stations

should be of 5 meters (for high levels of infestation) to 10 meters (for low levels of infestation).

Category(ies) of users

Trained professional

Pack sizes and packaging material

Labelled plastic (composite or HDPE) bottle sealed: 1.5 - 5 kg.
Labelled plastic (composite) sack, thermal welded: 1.5 - 25 kg.
Labelled plastic (composite) sack thermal welded - with inner plastic liner: 1.5 - 10 kg.
Labelled plastic composite (HDPE) bucket sealed: 1.5 - 25 kg.
Labelled plastic composite (HDPE) bucket sealed - with inner plastic liner: 1.5 - 10 kg.
Labelled carton sachet - with inner plastic liner: 1.5 - 10 kg.
Labelled cardboard - with inner plastic liner: 1.5 - 10 kg.

4.1.1 Use-specific instructions for use

A kezelési időszak után távolítsa el a megmaradó terméket.
Kövesse a vonatkozó helyi előírásokban/helyes kártevőirtási gyakorlatban szereplő további útmutatásokat.

4.1.2 Use-specific risk mitigation measures

Ahol lehetséges, a kezelés előtt tájékoztassa az esetlegesen közelben tartózkodókat (például a kezelt terület és környékének felhasználóit) a rágcsálóirtásról.
Vegye fontolóra a megelőző védelmi intézkedéseket (lyukak betömése, a rágcsálók táplálékforrásának eltávolítása, amennyire lehetséges) annak érdekében, hogy a rágcsálók a termékből többet fogyasszanak, és később kisebb valószínűséggel telepedjenek meg újból a területen.
A másodlagos mérgezés kockázatának csökkentése érdekében gyakran, a helyes kártevőirtási gyakorlat ajánlásaival összhangban távolítsa el a kezelés alatt elpusztult rágcsálókat.
Ne alkalmazza a terméket pulzáló / ritmusos kihelyezési mennyiséggel.

Ne alkalmazza a terméket tartósan (állandó kihelyezéssel).

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Ha az etetőállomásokat vízelvezető rendszerek közelében helyezi el, biztosítsa, hogy a csalétket ne érje víz.

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

4.2 Use description

Use 2 - House mice and rats - trained professionals – outdoor around buildings

Product type	PT14 - Rodenticides (Pest control)
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: House mouse Development stage: -juveniles and adults Scientific name: <i>Rattus norvegicus</i> Common name: Brown rat Development stage: -juveniles and adults Scientific name: <i>Rattus rattus</i> Common name: Roof rat Development stage: -juveniles and adults
Field(s) of use	Outdoor Outdoor around buildings

Application method(s)

Method: Bait application
Detailed description:
Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequencies

Application Rate: -
Dilution (%): -
Number and timing of application:

For mice:
40 g / bait station.
If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters (for high levels of infestation) to 10 meters (for low levels of infestation).

For rat:
100 g / bait station.
If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters (for high levels of infestation) to 10 meters (for low levels of infestation).

Category(ies) of users

Trained professional

Pack sizes and packaging material

Labelled plastic (composite or HDPE) bottle sealed: 1.5 - 5 kg.

Labelled plastic (composite) sack, thermal welded: 1.5 - 25 kg.
Labelled plastic (composite) sack thermal welded - with inner plastic liner: 1.5 - 10 kg.
Labelled plastic composite (HDPE) bucket sealed: 1.5 - 25 kg.
Labelled plastic composite (HDPE) bucket sealed - with inner plastic liner: 1.5 - 10 kg.
Labelled carton sachet - with inner plastic liner: 1.5 - 10 kg.

Labelled cardboard - with inner plastic liner: 1.5 - 10 kg.

4.2.1 Use-specific instructions for use

Protect bait from the weather. Place the bait stations in a place protected from rain and groundwater.
Replace any baits that have been damaged or soiled by the water in the feeder.
Remove the remaining product at the end of treatment period.

4.2.2 Use-specific risk mitigation measures

Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
Do not use the product in pulsed baiting treatments.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

-

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

-

5. General directions for use

5.1. Instructions for use

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.

Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).

Where possible, bait stations must be fixed to the ground or other structures.

Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).

When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.

Bait should be secured so that it cannot be dragged away from the bait station.

Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.

Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

Wear protective chemical resistant gloves during product handling phase

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.

If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.

If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.

Remove the remaining bait or the bait stations at the end of the treatment period.

Instructions for use that are "bait-specific":- Bait in sachets: Do not open the sachets containing the bait.

5.2. Risk mitigation measures

Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only"). Do not use in areas where resistance to the active substance can be suspected. Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment unless authorised for permanent baiting treatments. Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant. Do not wash the bait stations or utensils used in covered and protected bait points with water between applications. Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
Antidote: Vitamin K1 administered by medical/veterinary personnel only.
In case of:
Dermal exposure, wash skin with water and then with water and soap.
Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre
Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements.

Use of gloves is recommended.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
Store in places prevented from the access of children, birds, pets and farm animals.
Shelf life: 24 months

6. Other information

Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

This product contains a bittering agent and a dye.

Biocid number of expert report: 3625-1/2020/KBKHF

The label includes:

KILRAT PLUS rágcsálóirtó granulátum

Active substance: 0.005% brodifacoum

II. category pesticide

Authorisation holder: Bábolna Bio Ltd. (Budapest, Hungary)

Biocid license number: HU-2014-MA-14-00097-0000

as well as the following sections of this document:

2.2, 3, 4, (excluding packaging), and 5, 6 parts.

Plastic rod boxes shall bear the following text:

KILRAT PLUS rágcsálóirtó granulátum

Active substance: 0.005% brodifacoum

Antidote: K1 vitamíne

Authorisation holder: Bábolna Bio Ltd. (Budapest, Hungary)

Biocid license number: HU-2014-MA-14-00097-0000