

# Summary of product characteristics for a biocidal product

**Product name:** SALVESAFE E8

**Product type(s):** PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

PT03 - Veterinary hygiene (Disinfectants)

**Authorisation number:** EU-0020512-0000

**R4BP 3 asset reference number:** EU-0020512-0005

## Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	3
2. Product composition and formulation	3
2.1. Qualitative and quantitative information on the composition of the biocidal product	3
2.2. Type of formulation	3
3. Hazard and precautionary statements	3
4. Authorised use(s)	4
5. General directions for use	8
5.1. Instructions for use	8
5.2. Risk mitigation measures	8
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	9
5.4. Instructions for safe disposal of the product and its packaging	9
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	9
6. Other information	9

## Administrative information

### 1.1. Trade names of the product

SALVESAFE E8
--------------

### 1.2. Authorisation holder

**Name and address of the authorisation holder**

Name	SALVECO S.A.S.
Address	Avenue Pierre Mendès-France Z.A. Hellieule 4 F 88100 SAINT DIE DES VOSGES France

**Authorisation number**

EU-0020512-0000 1-2
---------------------

**R4BP 3 asset reference number**

EU-0020512-0005
-----------------

**Date of the authorisation**

28/08/2019
------------

**Expiry date of the authorisation**

28/08/2029
------------

### 1.3. Manufacturer(s) of the biocidal products

**Name of the manufacturer**

SALVECO S.A.S
---------------

**Address of the manufacturer**

Avenue Pierre Mendès-France, Z.I. Hellieule 4 F-88100 SAINT DIE DES VOSGES France
---

**Location of manufacturing sites**

Avenue Pierre Mendès-France, Z.I. Hellieule 4 F-88100 SAINT DIE DES VOSGES France
---

<b>Name of the manufacturer</b>	MULTIFILL BV BULKONT VANGST
<b>Address of the manufacturer</b>	CONSTRUCTIEWEG 25A 3641 SB MIJDRECHT Netherlands
<b>Location of manufacturing sites</b>	CONSTRUCTIEWEG 25A 3641 SB MIJDRECHT Netherlands

<b>Name of the manufacturer</b>	Diversey Netherlands Production BV
<b>Address of the manufacturer</b>	Rembrandtlaan 414 7545 ZW Enschede Netherlands
<b>Location of manufacturing sites</b>	Rembrandtlaan 414 7545 ZW Enschede Netherlands

<b>Name of the manufacturer</b>	Diversey UK Production Ltd
<b>Address of the manufacturer</b>	Cotes Park Industrial Estate, Somercotes DE55 4PA Alfreton United Kingdom
<b>Location of manufacturing sites</b>	Cotes Park Industrial Estate, Somercotes DE55 4PA Alfreton United Kingdom

<b>Name of the manufacturer</b>	Diversey España Production S.L.U
<b>Address of the manufacturer</b>	Avenida Conde Duque 5, 7 y 9, Poligono Industrial La Postura 28343 Valdemoro (Madrid) Spain
<b>Location of manufacturing sites</b>	Avenida Conde Duque 5, 7 y 9, Poligono Industrial La Postura 28343 Valdemoro (Madrid) Spain

<b>Name of the manufacturer</b>	Diversey Italy Production Srl
<b>Address of the manufacturer</b>	Strada Statale 235 26010 Bagnolo Cremasco (CR) Italy
<b>Location of manufacturing sites</b>	Strada Statale 235 26010 Bagnolo Cremasco (CR) Italy

<b>Name of the manufacturer</b>	Diversey Germany Production oHG
<b>Address of the manufacturer</b>	Morschheimer Strasse 12 67292 Kirchheimbolanden Germany
<b>Location of manufacturing sites</b>	Morschheimer Strasse 12 67292 Kirchheimbolanden Germany

#### 1.4. Manufacturer(s) of the active substance(s)

<b>Active substance</b>	1421 - Lactic acid
<b>Name of the manufacturer</b>	JUNGBUNGZLAUER S.A
<b>Address of the manufacturer</b>	Z. I Portuaire BP 32 67390 Marckolsheim France
<b>Location of manufacturing sites</b>	Z. I Portuaire BP 32 67390 Marckolsheim France

## 2. Product composition and formulation

### 2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Lactic acid		Active Substance		200-018-0	2,38

### 2.2. Type of formulation

AL - Any other liquid
-----------------------

## 3. Hazard and precautionary statements

**Hazard statements**

Safety data sheet available on request.

**Precautionary statements****4. Authorised use(s)****4.1 Use description****Use 1 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - spraying****Product type**

PT03 - Veterinary hygiene (Disinfectants)

**Where relevant, an exact description of the authorised use**

-

**Target organism(s) (including development stage)**

Scientific name: Bacteria  
Common name: Bacteria  
Development stage:

Scientific name: Yeasts  
Common name: Yeasts  
Development stage:

**Field(s) of use**

Indoor

Outdoor

**Application method(s)**

Method: Spraying  
Detailed description:  
Manual and automated spraying

**Application rate(s) and frequencies**

Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface.  
Dilution (%): Ready-to-use product  
Number and timing of application:  
Maximum 3 pre-milking and 3 post-milking applications per day

**Category(ies) of users**

Professional

**Pack sizes and packaging material**

Bottle, Plastic: HDPE 1L  
Jerrycan, Plastic: HDPE 5 - 60L  
Drum, Plastic: HDPE 220 L  
IBC, Plastic: HDPE 1000L

#### 4.1.1 Use-specific instructions for use

Before use, bring the product at least 20°C.  
Fill a manual or automate spraying device with ready to use product from the original packaging and close with spraying head.  
Pre-milking  
Apply the product by spray to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk.  
Post-milking  
The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

#### 4.1.2 Use-specific risk mitigation measures

Not applicable

#### 4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

#### 4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

#### 4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

#### 4.2 Use description

##### Use 2 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - foaming

##### Product type

PT03 - Veterinary hygiene (Disinfectants)

##### Where relevant, an exact description of the authorised use

-

##### Target organism(s) (including development stage)

Scientific name: Bacteria  
Common name: Bacteria  
Development stage:

Scientific name: Yeasts  
Common name: Yeasts  
Development stage:

<b>Field(s) of use</b>	Indoor Outdoor
<b>Application method(s)</b>	Method: Foam application Detailed description: Manual and automated foaming
<b>Application rate(s) and frequencies</b>	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking and 3 post-milking applications per day
<b>Category(ies) of users</b>	Professional
<b>Pack sizes and packaging material</b>	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

#### 4.2.1 Use-specific instructions for use

Before use, bring the product at least 20°C.  
Fill the foaming cup with ready to use product from the original packaging.  
Pre-milking  
Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk. Leave the teats skin dry.  
Post-milking  
The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

#### 4.2.2 Use-specific risk mitigation measures

Not applicable

#### 4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

#### 4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

#### 4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

### 4.3 Use description

#### Use 3 - Ready-to-use non-medical pre-milking and post-milking teat disinfection - dipping

<b>Product type</b>	PT03 - Veterinary hygiene (Disinfectants)
<b>Where relevant, an exact description of the authorised use</b>	-
<b>Target organism(s) (including development stage)</b>	Scientific name: Bacteria Common name: Bacteria Development stage:  Scientific name: Yeasts Common name: Yeasts Development stage:
<b>Field(s) of use</b>	Indoor  Outdoor
<b>Application method(s)</b>	Method: Dipping Detailed description: Manual and automated dipping
<b>Application rate(s) and frequencies</b>	Application Rate: Apply a maximum of 20 mL product per milking event to cover the treated surface. Dilution (%): Ready-to-use product Number and timing of application: Maximum 3 pre-milking and 3 post-milking applications per day
<b>Category(ies) of users</b>	Professional
<b>Pack sizes and packaging material</b>	Bottle, Plastic: HDPE 1L Jerrycan, Plastic: HDPE 5 - 60L Drum, Plastic: HDPE 220 L IBC, Plastic: HDPE 1000L

### 4.3.1 Use-specific instructions for use

Before use, bring the product at least 20°C.

Fill the manual or automated dipping cup with ready to use product from the original packaging.

#### *Pre-milking*

Apply the product to the full length of each teat. Leave to act at least 1 min. Wipe the product with a single-use paper towel/cloth to avoid any contamination of milk. Leave the teats skin dry.

#### *Post-milking*

The teat should be treated immediately after milking in such a way that the teats are fully covered with the product. Keep the animals standing or walking at least 5 minutes.

### 4.3.2 Use-specific risk mitigation measures

Not applicable

### 4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Section 5.3.

### 4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Section 5.4.

### 4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Section 5.5.

## 5. General directions for use

### 5.1. Instructions for use

Please refer to use-specific instructions for use

### 5.2. Risk mitigation measures

Not applicable.

### 5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

In case of eye contact: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
In case of skin contact: Wash with plenty of water.  
If swallowed: Rinse mouth. Do not induce vomiting. If symptoms occurs: Get medical advice/attention.  
  
If inhaled: Get medical attention if symptoms occur.

### 5.4. Instructions for safe disposal of the product and its packaging

Dispose of contents/container to in accordance with national regulation.

### 5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

Store in original container at room temperature. Protect from direct sun light. Protect from frost. Shelf-life: 2 years.

### 6. Other information

N/A