


Regulation (EU) No 528/2012 concerning the making available on the market and use of biocidal products

**PRODUCT ASSESSMENT REPORT OF A BIOCIDAL
PRODUCT FOR THE RENEWAL OF A NATIONAL
AUTHORISATION**



Product identifier in R4BP	ZAS CEBO FRESCO 005
Product type(s):	14 (Rodenticide)
Active ingredient(s):	Bromadiolone
Case No. in R4BP	BC-GA035313-70
Asset No. in R4BP	ES-0017722-0000
Evaluating Competent Authority	Spain
Internal registration/file no	ES/BB(NA)-2018-14-00452
Date	February 2018

Table of content

1	Conclusion	3
2	Summary of the product assessment	5
2.1	Administrative information	5
2.2	Composition and formulation.....	5
2.3	Classification and Labelling according to the Regulation (EC) No 1272/2008.....	7
2.4	Use(s) appropriate for further authorisation	8
2.5	General directions for use	16
3	Assessment of the product	20
3.1	Use(s) considered appropriate for authorisation after former assessment (uses currently under authorisation in Spain).....	20
3.2	Physical, chemical and technical properties.....	21
3.3	Physical hazards and respective characteristics.....	21
3.4	Methods for detection and identification.....	21
3.5	Efficacy against target organisms	21
3.6	Risk assessment for human health	22
3.7	Risk assessment for animal health.....	23
3.8	Risk assessment for the environment	24
3.9	Assessment of a combination of biocidal products	24
3.10	Comparative assessment	24
4	Confidential annex (Access level: “Restricted” to applicant and authority).....	25
4.1	Full composition of the product	25

1 Conclusion

The assessment presented in this report has shown that the ready-to-use product, ZAS CEBO FRESCO 005 with the active substance bromadiolone, at a level of 0.005% w/w, may be authorised for use as a rodenticide (product-type 14) since the conclusions of initial evaluation remain valid.

This product is an identical product to AGRORAT BD-5 PASTA PRO (asset number ES-0005692-0000). The applicant has submitted letters of access to AGRORAT BD-5 PASTA PRO and AGRORAT BD-5 PASTA in order to use the information/data.

However, the biocidal product ZAS CEBO FRESCO 005 contains 0.005 %w/w bromadiolone and the Commission Regulation (EU) 2016/1179 of 19 July 2016 amending, for the purposes of its adaptation to technical and scientific progress, Regulation (EC) No 1272/2008 of the European Parliament and of the Council on classification, labelling and packaging of substances and mixtures has been applied.

Due to national legislation in relation to categories of users which three categories of users are established (general public, professional and trained professional user) based on the qualification obtained, therefore the professional is extrapolated to the general public (under this national regulation the professional user is not bounded to use PPE when they apply the product). For that, the biocidal product rodenticides containing 0.005 %w/w bromadiolone only can be authorised by trained professional user because of the toxicological classification the use of PPE are mandatory. Given that, this legislation is national and in other Member States legislation could be different, each Competent Authority should consider that in order to grant the authorisation.

Therefore, ZAS CEBO FRESCO 005 is granted as a rodenticide product against house mice (*Mus musculus*) and brown rats (*Rattus norvegicus*). It is to be used indoors, outdoors around buildings, outdoor in open areas and waste dumps and sewers by trained professional. It is a ready to used paste bait to be used in tamper-resistant bait stations or anchored in sewers. The specific intended uses of the product are in section 2.4. of this assessment report..

The risk assessment for the environment has been performed for the intended uses in and around buildings, in open areas, around waste sites and in sewers since the concentration of the active substance is the same, the new evaluation shows that the conclusions for the first evaluation remain valid.

The overall conclusion is that the intended uses of ZAS CEBO FRESCO 005 do not pose an unacceptable risk to the sewage treatment plant, soil, air, surface water, sediment, and groundwater compartments.

However, an unacceptable risk is however identified for the primary and secondary poisoning of non-target vertebrates, and specific risk mitigation measures on the use of the product are required to reduce the risk for the environment. The measures include use of tamper resistant bait boxes, collection of unconsumed baits after termination of the control campaign and collection of dead rodents during and after the control campaign at frequent intervals, proper disposal of dead rodents and unused baits, restriction of the use in open areas to trained professional users only.

According to the renewal of anticoagulant active substance for trained professional users the product may be authorised for use in covered and protected bait points other than tamper resistant bait stations. The applicant has not submitted any additional information to include these application methods, so the ES CA does not authorise other use different to tamper resistant bait stations.

Please, note that this assessment report includes all uses requested by the applicant and assessed by ES CA, only as information for the concerned Member States.

Spanish CA only grants the use of ZAS CEBO FRESCO 005 according to the table 5 included in this assessment report due to our national risk mitigation measures.

2 Summary of the product assessment

2.1 Administrative information

2.1.1 Identifier in R4BP

ZAS CEBO FRESCO 005

2.1.2 Manufacturer(s) of the product

Name of manufacturer	IMPEX EUROPA, S.L
Address of manufacturer	Avda. Pontevedra, 39. 36600 Villagarcía de Arosa. Pontevedra. Spain
Location of manufacturing sites	Polígono industrial TRABANCA-BADIÑA, Parcelas 22-24 36600 Villagarcía de Arosa. Pontevedra. Spain

2.1.3 Manufacturer(s) of the active substance(s)

Active substance	Bromadiolone
Name of manufacturer	LABORATORIOS AGROCHEM S.L.
Address of manufacturer	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN
Location of manufacturing sites	C/ Tres Rieres, 10 08292 - Esparreguera (Barcelona) SPAIN

2.2 Composition and formulation

2.2.1 Qualitative and quantitative information on the composition

Table 1

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
-------------	------------	----------	------------	-----------	-------------

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Bromadiolone	3-[(1RS,3RS;1RS,3SR)-3-(4'-bromobiphenyl-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxycoumarin	Active substance	28772-56-7	249-205-9	0.005
-	-	Non-active substance	-	-	-

- The product contains a bittering agent and a dye.
 - Information on the full composition is provided in the confidential annex (see chapter 4).
- According to the information provided the product contains no nanomaterial as defined in Article 3 paragraph 1 (z) of Regulation No. 528/2012:

2.2.2 Information on the substance(s) of concern

No substance of concern was identified upon initial assessment (the application for authorisation was submitted and the assessment took place before the Biocidal Products Regulation 528/2012 entered into force).

2.2.3 Candidate(s) for substitution

No candidate for substitution was identified upon initial assessment (the application for authorisation was submitted and the assessment took place before the Biocidal Products Regulation 528/2012 entered into force).

Now that the Biocidal Products Regulation 528/2012 entered into force, the following substance(s) was/were identified as candidate(s) for substitution upon this renewal:

Bromadiolone does meet the exclusion criteria according to Article 5(1) BPR. Because the following exclusion criteria are met:

- toxic for reproduction category 1B
- persistent, bioaccumulative and toxic

And therefore, Bromadiolone does meet the conditions laid down in Article 10 BPR, and is consequently a candidate for substitution.

2.2.4 Type of formulation


Ready-to-use bait: paste.

2.3 Classification and Labelling according to the Regulation (EC) No 1272/2008

Table 2

Classification	
Hazard classes, Hazard categories	Hazard statements
Reproductive toxicity; Repr. 1B	H360D May damage the unborn child
Specific target organ toxicity — repeated exposure; STOT RE 1	H372 Causes damage to organs (blood) through prolonged or repeated exposure

Table 3

Labelling		
	Code	Pictogram / Wording
	GHS08	
Signal word	-	Danger
Hazard statements	H360D	May damage the unborn child
	H372	Causes damage to organs (blood) through prolonged or repeated exposure
Supplemental hazard information	-	-
Supplemental label elements	-	-
Precautionary statements	P201	Obtain special instructions before use.
	P202	Do not handle until all safety precautions have been read and understood.
	P264	Wash ... thoroughly after handling
	P270	Do not eat, drink or smoke when using this product.
	P280	Wear protective gloves/ protective clothing/eye protection/face protection
	P314	Get medical advice/attention if you feel unwell.
	P405	Store locked up.
	P501	Dispose of contents and/ or container as a hazardous waste to a registered establishment or undertaking, in accordance with current regulations.
Note	-	

2.4 Use(s) appropriate for further authorisation

In order to make proper use of the standard sentences for SPCs for rodenticides it is considered necessary to split the uses currently authorised in Spain further down:

Table 4

Use(s) considered appropriate for authorisation after former assessment (uses currently under authorisation in Spain)		Use(s) appropriate for further authorisation	
1	House mice and/or rats –general public – indoor and outdoor around buildings.	1	House mice and/or brown rats – trained professionals – indoor
2	House mice and/or rats – professional– indoor and outdoor around buildings.	2	House mice and/or brown rats – trained professionals – outdoor around buildings
		3	Brown rats – trained professionals – Outdoor open areas & waste dumps
3	House mice and/or rats –trained professional– indoor, outdoor around buildings and open areas.	4	Brown rats – trained professionals – Sewers

Uses authorized in Spain according national Risk Mitigation Measures

Table 5

Use(s) considered appropriate for authorisation after former assessment (uses currently under authorisation in Spain)	Use(s) appropriate for authorisation in Spain according national Risk Mitigation Measures.
House mice and/or rats –general public – indoor and outdoor around buildings.	House mice and/or brown rats – trained professionals – indoor
House mice and/or rats – professional– indoor and outdoor around buildings.	House mice and/or brown rats – trained professionals – outdoor around buildings
House mice and/or rats –trained professional– indoor, outdoor around buildings and open areas	Brown rats – trained professionals – Outdoor open areas & waste dumps

2.4.1 Use 1 - House mice and/or brown rats – trained professionals – indoor

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	Mus musculus (house mice) <i>Rattus norvegicus</i> (brown rat)
Field(s) of use	Indoor
Application method(s)	- Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequency	<p>Rats, bait boxes with 100-200 g per baiting point</p> <ul style="list-style-type: none"> · up to 200g of product using: <ul style="list-style-type: none"> - sachets of 20g - 1 tray of 160g or 2 trays of 80g. - caulking tube of 300g. · up to 100g of product using: <ul style="list-style-type: none"> - sachets of 10, 15 or 20g - 1 tray of 80g. - caulking tube of 300g. <p>Mice, bait boxes with 60-100 g per baiting point.</p>
Category(ies) of users	Trained professionals
Pack sizes and packaging material	<p>Minimum pack size of 3 kg.</p> <p>Number of packed bags per packaging: up to 25Kg</p> <p>Grams/kg of bait per packed bag: 10, 15 y 20g.</p> <p>Packaging material:</p> <ul style="list-style-type: none"> • Individual nonwoven tissue/cotton tea net sachets of 10, 15 and 20g. They are delivered inside kraft paper and plastic (polyethylene or polypropylene) bags, or carton box or plastic (HDPE) buckets of, 5, 10, 15, 20 or 25kg • Plastic (HDPE, PE or PP) tubes of 300g to apply via caulking gun, delivered inside carton boxes up to 10 units. • Pre-filled plastic (HDPE, PE or PP) trays of 80g delivered inside carton boxes up to 50 units. • Pre-filled plastic (HDPE, PE or PP) trays of 160g delivered inside carton boxes up to 25 units.

2.4.1.1 Use-specific instructions for use

- Remove the remaining product at the end of treatment period.
- Follow any additional instructions provided by the relevant code of best practice.

2.4.1.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use the product in pulsed baiting treatments.
- This product shall only be used indoors and in places that are not accessible to children or non-target animals.

2.4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to water drainage systems, ensure that bait contact with water is avoided.

2.4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See section 5.4.

2.4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See section 5.5.

2.4.2 Use 2 - Mice and/or brown rats – trained professionals – outdoor around buildings

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rat)
Field(s) of use	Outdoor around buildings.
Application method(s)	- Ready-to-use bait to be used in tamper-resistant bait stations

Application rate(s) and frequency	<p>Rats: bait boxes with 100-200 g per baiting point</p> <ul style="list-style-type: none"> · up to 200g of product using: <ul style="list-style-type: none"> - sachets of 20g - 1 tray of 160g or 2 trays of 80g. - caulking tube of 300g. · up to 100g of product using: <ul style="list-style-type: none"> - sachets of 10, 15 or 20g - 1 tray of 80g. - caulking tube of 300g. <p>Mice: bait boxes with 60-100 g per baiting point</p>
Category(ies) of users	Trained professionals
Pack sizes and packaging material ¹	<p>Minimum pack size of 3 kg.</p> <p>Number of packed bags per packaging: up to 25Kg</p> <p>Grams/kg of bait per packed bag: 10, 15 y 20g.</p> <p>Packaging material:</p> <ul style="list-style-type: none"> • Individual nonwoven tissue/cotton tea net sachets of 10, 15 and 20g. They are delivered inside kraft paper and plastic (polyethylene or polypropylene) bags, or carton box or plastic (HDPE) buckets of, 5, 10, 15, 20 or 25kg • Plastic (HDPE, PE or PP) tubes of 300g to apply via caulking gun, delivered inside carton boxes up to 10 units. • Pre-filled plastic (HDPE, PE or PP) trays of 80g delivered inside carton boxes up to 50 units. • Pre-filled plastic (HDPE, PE or PP) trays of 160g delivered inside carton boxes up to 25 units.

2.4.2.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the baiting points in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- Follow any additional instructions provided by the relevant code of best practice.

2.4.2.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign *[in accordance with the applicable code of good practice, if any]*.
- Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

2.4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

2.4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See section 5.4.

2.4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See section 5.5.

2.4.3 Use 3 – Brown rats – trained professionals – Outdoor open areas & waste dumps

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Rattus norvegicus</i> (brown rat)
Field(s) of use	Outdoor open areas Outdoor waste dumps
Application method(s)	- Ready-to-use bait to be used in tamper-resistant bait stations.
Application rate(s) and frequency	Rats: bait boxes with 100-200 g per baiting point · up to 200g of product using: <ul style="list-style-type: none"> - sachets of 20g - 1 tray of 160g or 2 trays of 80g. - caulking tube of 300g. · up to 100g of product using: <ul style="list-style-type: none"> - sachets of 10, 15 or 20g - 1 tray of 80g. - caulking tube of 300g.
Category(ies) of users	Trained professionals only
Pack sizes and packaging material	Minimum pack size of 3 kg. Number of packed bags per packaging: up to 25Kg Grams/kg of bait per packed bag: 10, 15 y 20g. Packaging material: <ul style="list-style-type: none"> • Individual nonwoven tissue/cotton tea net sachets of 10, 15 and 20g. They are delivered inside kraft paper and plastic (polyethylene or polypropylene) bags, or carton box or plastic (HDPE) buckets of, 5, 10, 15, 20 or 25kg • Plastic (HDPE, PE or PP) tubes of 300g to apply via caulking gun, delivered inside carton boxes up to 10 units. • Pre-filled plastic (HDPE, PE or PP) trays of 80g delivered inside carton boxes up to 50 units. • Pre-filled plastic (HDPE, PE or PP) trays of 160g delivered inside carton boxes up to 25 units.

2.4.3.1 Use-specific instructions for use

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- Follow any additional instructions provided by the relevant code of best practice.

2.4.3.2 Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign [in accordance with the applicable code of good practice, if any].
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use this product as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.
- Do not use this product in pulsed baiting treatments.
- Do not apply this product directly in the burrows.

2.4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

2.4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See section 5.4.

2.4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See section 5.5.

2.4.4 Use 4 – Brown rats – trained professionals – sewers

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Rattus norvegicus</i> (brown rat)
Field(s) of use	Sewers
Application method(s)	- Ready-to-use bait to be anchored or applied in bait stations preventing the bait from getting into contact with waste water.
Application rate(s) and frequency	Bait boxes with 100-200 g per manhole .
Category(ies) of users	Trained professionals only
Pack sizes and packaging material	<p>Minimum pack size of 3 kg. Number of packed bags per packaging: up to 25Kg Grams/kg of bait per packed bag: 10, 15 y 20g. Packaging material:</p> <ul style="list-style-type: none"> • Individual nonwoven tissue/cotton tea net sachets of 10, 15 and 20g. They are delivered inside kraft paper and plastic (polyethylene or polypropylene) bags, or carton box or plastic (HDPE) buckets of, 5, 10, 15, 20 or 25kg • Plastic (HDPE, PE or PP) tubes of 300g to apply via caulking gun, delivered inside carton boxes up to 10 units. • Pre-filled plastic (HDPE, PE or PP) trays of 80g delivered inside carton boxes up to 50 units. • Pre-filled plastic (HDPE, PE or PP) trays of 160g delivered inside carton boxes up to 25 units.

2.4.4.1 Use-specific instructions for use

- Baits must be applied in a way so that they do not come into contact with water and are not washed

away.

- Follow any additional instructions provided by the relevant code of best practice.

2.4.4.2 Use-specific risk mitigation measures

- [If national policy or legislation requires it] Place baits only in sewer systems which are connected to the sewage treatment plant.
- Do not use this product in pulsed baiting treatments.

2.4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See section 5.3.

2.4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

See section 5.4.

2.4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See section 5.5.

2.5 General directions for use

2.5.1 Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the

infestation.

- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (*see section 5.3 for the information to be shown on the label*).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves during product handling phase (glove material to be specified by the authorisation holder within the product information).
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of

traps as an alternative control measure.

- Place paste bait with a sufficiently elongated applicator (spatula) to reduce hand exposure - avoid reaching into the bucket.
- Specify how the equipment (e.g. spatula) shall be cleaned and how contact with residues of the bait can be avoided.

2.5.2 Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign [in accordance with the applicable code of good practice, if any]".
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment [unless authorised for permanent baiting treatments].
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

2.5.3 Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, always check for and remove contact lenses, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label [insert country specific information]. Contact a veterinary surgeon in case

of ingestion by a pet [insert country specific information]

- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre [insert national phone number]"
- Hazardous to wildlife.

2.5.4 Instructions for safe disposal of the product and its packaging

- At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements [The method of disposal shall be described specifically in the national SPC and be reflected on the product label].

2.5.5 Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 2 years.

2.5.6 Other information

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.

3 Assessment of the product

3.1 Use(s) considered appropriate for authorisation after former assessment (uses currently under authorisation in Spain)

3.1.1 Use 1 – House mice and/or brown rats – trained professional– in and around buildings and open areas and waste dumps.

Product Type(s)	14
Where relevant, an exact description of the use	Rodenticide
Target organism(s) (including development stage)	<i>Mus musculus</i> (house mice) <i>Rattus norvegicus</i> (brown rats)
Field(s) of use	In and around buildings, open areas, waste dumps.
Application method(s)	The biocidal product is ready to use paste bait in bait stations.
Application rate(s) and frequency	<p>For the control of rats, each bait point should be placed each 10 m² and usually contains:</p> <ul style="list-style-type: none"> · up to 200g of product using: - sachets of 20g - 1 tray of 160g or 2 trays of 80g. - caulking tube. · up to 100g of product using: - sachets of 10, 15 or 20g - 1 tray of 80g. - caulking tube. <p>For the control of mice, each bait point should be placed each 10 m² and usually contains:</p> <ul style="list-style-type: none"> · up to 80 g of bait using 1 tray of 80g. · up to 50 g of bait using sachets of 10, 15 y 20g or caulking tube.
Category(ies) of users	Trained Professional
Pack sizes and packaging material	<ul style="list-style-type: none"> -Individual nonwoven tissue/cotton tea net sachets of 10, 15 and 20g. They are delivered inside kraft paper and plastic (polyethylene or polypropylene) bags, or carton box or plastic (HDPE) buckets of 500g and 1, 2, 2.5, 5, 10, 15, 20 or 25kg for trained professional users. -Plastic (HDPE, PE or PP) tubes of 300g to apply via caulking gun, delivered inside carton boxes up to 10 units. -Pre-filled plastic (HDPE, PE or PP) trays of 80g delivered inside carton boxes up to 50 units. -Pre-filled plastic (HDPE, PE or PP) trays of 160g delivered inside carton boxes up to 25 units.

3.2 Physical, chemical and technical properties

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding physical, chemical and technical properties remains valid.

3.3 Physical hazards and respective characteristics

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding physical hazards and respective characteristics remains valid.

3.4 Methods for detection and identification

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding methods for detection and identification remains valid.

3.5 Efficacy against target organisms

The conclusion from the former assessment regarding efficacy against target organisms remains valid. This product is identical to AGRORAT BD-5 PASTA PRO. A letter of access has been submitted in order to use the information/data.

According to the test provided included in the letter of access, ES CA consider that the biocidal product is effective against rats (indoor, outdoors around buildings, outdoor open areas, waste dumps and sewers) and mice (indoor and outdoors around buildings).

3.6 Risk assessment for human health

3.6.1 Assessment of effects of the active substance on human health

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding effects of the active substance on human health remains valid.

3.6.2 Assessment of effects of the product on human health

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding effects of the product on human health remains valid.

3.6.3 Exposure assessment

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding the exposure remains valid.

3.6.4 Risk characterisation for human health

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding the risk characterisation for human health remains valid.

3.6.4.1 Risk for trained professional users

The conclusion from the former assessment regarding the risk characterisation for trained professional user remains valid.

3.6.4.2 Risk for professional users

Due to national legislation in relation to categories of users which three categories of users are established (general public, professional and trained professional user) based on the qualification obtained, therefore the professional is extrapolated to the general public (under this national regulation the professional user is not bounded to use PPE when they apply the product). For that, the biocidal product rodenticides containing 0.005 %w/w bromadiolone only can be authorised by trained

professional user because of the toxicological classification the use of PPE are mandatory. Given that, this legislation is national and in other Member States legislation could be different, each Competent Authority should consider that in order to grant the authorisation.

3.6.4.3 Risk for the general public

According to the Commission Regulation (EU) 2016/1179 of 19 July 2016 amending, for the purposes of its adaptation to technical and scientific progress, Regulation (EC) No 1272/2008 of the European Parliament and of the Council on classification, labelling and packaging of substances and mixtures, the biocidal product containing anticoagulant active substance cannot be authorised by general public if the concentration in the biocidal product is above the specific limit concentration ($\geq 0.003\%$).

3.6.4.4 Risk for consumers via residues in food

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding risks for consumers via residues in food remain valid.

3.6.4.5 Risk characterisation from combined exposure to several active substances or substances of concern within a biocidal product

The biocidal product does not contain other substances in quantities that would be of toxicological concern in the production formulation.

3.6.4.6 Summary of risk characterisation

The conclusion from the former assessment regarding risk characterisation remains valid, except to the authorisation for general public and professional user which have been removed to the authorisation in order to comply with the requirements laid down in Commission Regulation (EU) 2016/1179 of 19 July 2016 amending, for the purposes of its adaptation to technical and scientific progress, Regulation (EC) No 1272/2008 of the European Parliament and of the Council on classification, labelling and packaging of substances and mixtures.

3.7 Risk assessment for animal health

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding animal health remains valid.

3.8 Risk assessment for the environment

Neither new data was not provided nor had new guidance to be taken into account for re-assessment. Accordingly, the conclusion from the former assessment regarding the environment remains valid.

3.9 Assessment of a combination of biocidal products

A use with other biocidal products is not intended.

3.10 Comparative assessment

As bromadiolone is a Candidate for Substitution, a comparative assessment must be carried out as part of the evaluation process.

The Biocidal Products Committee of the European Chemicals Agency published its Opinion on Questions regarding the comparative assessment of anticoagulant rodenticides on 02 March 2017 (Document no. ECHA/BPC/145/2017).

The Decision states that:

- In the absence of anticoagulant rodenticides, the use of rodenticide biocidal products containing other active substances would lead to an inadequate chemical diversity to minimize the occurrence of resistance in the target harmful organisms. These products also show some significant practical or economical disadvantages for the relevant uses.
- There is insufficient scientific evidence to prove that non-chemical alternative methods of rodent control are sufficiently effective according to the criteria established in agreed Union guidance with a view to prohibit or restrict the authorised uses of anticoagulant rodenticides.

The Decision forms the basis of the COMMISSION IMPLEMENTING DECISION (EU) 2017/1532 of 7 September 2017 addressing questions regarding the comparative assessment of anticoagulant rodenticides in accordance with Article 23(5) of Regulation (EU) No 528/2012 of the European Parliament and of the Council.

On the basis of this comparative assessment, the authorisation of rodenticide products containing bromadiolone is justified.