

**Annex XV dossier**

**PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A  
CMR CAT 1 OR 1B, PBT, vPvB OR A SUBSTANCE OF AN  
EQUIVALENT LEVEL OF CONCERN**

**Substance Name(s): Tricosafluorododecanoic acid**

**EC Number(s): 206-203-2**

**CAS Number(s): 307-55-1**

**Submitted by: BAuA  
Federal Office for Chemicals  
Friedrich-Henkel-Weg 1 – 25  
44149 Dortmund  
Germany**



# CONTENTS

PREFACE.....	3
PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A CMR CAT 1A OR 1B, PBT, VPVB OR A SUBSTANCE OF AN EQUIVALENT LEVEL OF CONCERN .....	4
PART I JUSTIFICATION.....	6
1 IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES .....	6
1.1 Name and other identifiers of the substance .....	6
1.2 Composition of the substance .....	7
1.3 Physico-chemical properties.....	7
2 HARMONISED CLASSIFICATION AND LABELLING .....	10
3 ENVIRONMENTAL FATE PROPERTIES.....	11
3.1 Degradation .....	11
3.1.1 Abiotic degradation .....	11
3.1.2 Biodegradation .....	15
3.1.3 Summary and discussion on degradation .....	20
3.2 Environmental distribution .....	21
3.2.1 Adsorption/desorption .....	21
3.2.2 Volatilisation .....	21
3.2.3 Distribution modelling .....	22
3.3 Bioaccumulation.....	22
3.3.1 Aquatic bioaccumulation.....	22
3.3.2 Summary and discussion of bioaccumulation .....	31
4 HUMAN HEALTH HAZARD ASSESSMENT.....	32
5 ENVIRONMENTAL HAZARD ASSESSMENT .....	33
6 CONCLUSIONS ON THE SVHC PROPERTIES .....	34
6.1 PBT, vPvB assessment .....	34
6.1.1 Assessment of PBT/vPvB properties – comparison with the criteria of Annex XIII .....	34
6.1.2 Summary and overall conclusions on the PBT, vPvB properties .....	35
6.2 CMR assessment.....	36
6.3 Substances of equivalent level of concern assessment. ....	36
PART II .....	37
INFORMATION ON USE, EXPOSURE, ALTERNATIVES AND RISKS .....	37
ANNEX I - READ-ACROSS APPROACH.....	40
LITERATURE CITED .....	50

## TABLES

Table 1: Substance identity.....	6
Table 2: Overview of physicochemical properties .....	8
Table 3: Summary of photodegradation studies for C <sub>8</sub> -PFCA and its ammonium salt.....	13
Table 4: Screening tests for C <sub>12</sub> -PFCA.....	15
Table 5: Summary of simulations tests of C <sub>8</sub> -PFCA (PFOA) and its ammonium salt (APFO) and sodium salt ...	16
Table 6 Bioconcentration factors (BCF) of C <sub>12</sub> -PFCA.....	24
Table 7: Available BMF data of C <sub>12</sub> -PFCA; if not indicated otherwise BMFs refer to whole body .....	26
Table 8: Available Trophic Magnification Factors (TMF) for C <sub>12</sub> PFCA; if not indicated otherwise TMFs refer to whole body. ....	29
Table 9: BMF and TMF values for C <sub>12</sub> -PFCA for a terrestrial food chain (Müller et al., 2011). ....	31
Table 10: Alternative compounds, their product names, company and use for C <sub>8</sub> -PFCA and its salts. ....	38
Table 11: CAS-Numbers and similarity of chemical structures of long chain PFCAs. ....	40
Table 12: Basic substance information and physical chemical properties relevant to justify read across in the PBT assessment. ....	42
Table 13 : Information on BCF, BMF and TMF of C <sub>9-14</sub> PFCAs relevant to justify read across in the B assessment..	46

## FIGURES

Figure 1: BMFs for C <sub>11-13</sub> -PFCAs in a remote terrestrial food chain from two different locations (whole body, Müller et al., 2011). The study is reliable (reliability 2).....	48
Figure 2: TMFs for C <sub>11-13</sub> -PFCAs in a remote terrestrial food chain from two different locations (whole-body, Müller et al., 2011). The study is reliable (reliability 2).....	48

## PREFACE

In four provided dossiers, the intrinsic properties of four perfluorinated carboxylic acids (PFCAs) are assessed: C<sub>11-14</sub>-PFCAs. Many studies are only available for structurally similar shorter chain PFCAs such as C<sub>8</sub>-PFCA and C<sub>9</sub>-PFCA. In those cases where studies on the particular substance are missing, studies from either shorter or longer chain PFCAs are used in the provided dossiers by applying read-across. Read-across is based on the structural similarities and on the physicochemical properties, which follow a regular pattern. All PFCAs contain a carboxylic acids group and a perfluorinated carbon chain. The only difference is the number of CF<sub>2</sub>-groups in this chain. Details on the read-across approach, i.e. showing the trend of physicochemical properties and the structural similarities are given in Annex I.

## **PROPOSAL FOR IDENTIFICATION OF A SUBSTANCE AS A CMR CAT 1A OR 1B, PBT, VPvB OR A SUBSTANCE OF AN EQUIVALENT LEVEL OF CONCERN**

**Substance Name(s): Tricosafluorododecanoic acid**

**EC Number(s): 206-203-2**

**CAS number(s): 307-55-1**

- It is proposed to identify the substance(s) as vPvB according to Article 57 (e).

### **Summary of how the substance(s) meet(s) the CMR (Cat 1A or 1B), PBT or vPvB criteria, or is/are considered to be (a) substance(s) giving rise to an equivalent level of concern**

For C<sub>12</sub>-PFCA no abiotic degradation studies and only one standard screening study are available. Applying the read across approach, data from structurally similar compounds can be used to evaluate the degradation potential of the substance. C<sub>8-14</sub>-PFCAs contain a highly similar chemical structure, a perfluorinated carbon chain and a carboxylic acid group. The compounds differ only in the number of CF<sub>2</sub>-groups.

Comparing the physico-chemical properties of C<sub>8-14</sub>-PFCAs it becomes obvious that with increasing chain length water solubility decreases and the sorption potential increases. This trend is based on the increasing number of CF<sub>2</sub>-groups in the molecular structure. According to the read-across approach these chemicals follow a regular pattern as a result of structural similarity. Those substances may therefore be considered as a group or a category of substances and the read-across approach can be applied.

In general, the persistence of long chain PFCAs can be explained by the shielding effect of the fluorine atoms, blocking e.g. nucleophilic attacks to the carbon chain. High electronegativity, low polarizability and high bond energies make highly fluorinated alkanes to the most stable organic compounds. It is not expected that the carboxylic group in PFCAs alters this persistence of these chemicals. This fact is confirmed by a study which obtained a DT<sub>50</sub> of >92 years for C<sub>8</sub>-PFCA in water. Screening studies of C<sub>12</sub>-PFCA showed no biodegradation within 28 days. Non-standard tests with C<sub>8</sub>-PFCA could not detect any degradation products under environmentally relevant conditions. Moreover, a monitoring study showed that C<sub>8</sub>-PFCA remained in soil and groundwater, years after application of fire fighting foam which contained PFCAs. Furthermore, screening biodegradation studies on C<sub>8,9,12,14</sub>-PFCAs, one simulation study on aerobic aquatic biodegradation and monitoring studies from contaminated sites on C<sub>8</sub>-PFCA in soil and groundwater indicate that these substances may be persistent

Therefore, we conclude that C<sub>12</sub>-PFCA is not degraded in the environment and thus fulfils the P- and vP-criteria under REACH.

The available BCF-values of C<sub>12</sub>-PFCA are above 5000. Thus, the B as well as the vB-criteria according to Annex XIII of REACH are fulfilled.

In conclusion, C<sub>12</sub>-PFCA is a vPvB-substance according to Art. 57e) of REACH.

The substance has not yet been registered under REACH.

# PART I

## JUSTIFICATION

### 1 IDENTITY OF THE SUBSTANCE AND PHYSICAL AND CHEMICAL PROPERTIES


#### 1.1 Name and other identifiers of the substance

**Table 1: Substance identity**

<b>EC number:</b>	206-203-2
<b>EC name:</b>	Tricosafuorododecanoic acid
<b>CAS number (in the EC inventory):</b>	307-55-1
<b>CAS number:</b>	307-55-1
<b>CAS name:</b>	Dodecanoic acid, 2,2,3,3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11,12,12,12- tricosafuoro-
<b>IUPAC name:</b>	Tricosafuorododecanoic acid
<b>Index number in Annex VI of the CLP Regulation</b>	-
<b>Molecular formula:</b>	$C_{12}HF_{23}O_2$
<b>Molecular weight range:</b>	614.0984 g/mol
<b>Synonyms:</b>	Dodecanoic acid, tricosafuoro- $C_{12}$ -PFCA Perfluorododecanoic acid Perfluorolauric acid



**Structural formula:**



**1.2 Composition of the substance**

**Name:** Tricosfluorododecanoic acid

**Description:** Mono-constituent substance

**Degree of purity:** Registration dossiers or other information on concentration ranges and on any impurities are not available.

**1.3 Physico-chemical properties**

The following physico-chemical properties are mainly obtained by calculation. No further information regarding the given values or other properties is available.

**Table 2: Overview of physicochemical properties**

Property	Value	Remarks	Reference
Physical state at 20°C and 101.3 kPa	solid	According to melting point	
Melting/freezing point	112 – 114 °C	Experimental result	Huang B-N, Haas A., Lieb M., 1987
Boiling point	249 °C	Data from SRC PhysProp Database "PhysProp" data were obtained from Syracuse Research Corporation of Syracuse, New York (US)	Data from SRC PhysProp Database, 02/2012 "PhysProp" data were obtained from Syracuse Research Corporation of Syracuse, New York (US)
Vapour pressure	0.0043 mm Hg	Estimated value	
	9.40E-3 Torr at 25°C		Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (© 1994-2012 ACD/Labs)
Water solubility	4.81 E-006 mg/l	Estimated value	
	2.9E-5 g/L pH 1 at 25°C		Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (© 1994-2012 ACD/Labs)
	2.2E-4 g/L pH 2 at 25°C		
	2.0E-3 g/L pH 3 at 25°C		
	0.014 g/L pH 4 at 25°C		
	0.034 g/L pH 5 at 25°C		
	0.039 g/L pH 6 at 25°C		
0.040 g/L pH 7 at 25°C 0.041 g/L pH 8-10 at 25°C			
Partition coefficient n-octanol/water	10.16	Estimated value	
	logP 9.363±0.888 at 25°C	The value was most likely calculated for the non-ionised form.	Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (© 1994-2012 ACD/Labs)
Dissociation constant	0.52±0.10 pH:	-	Calculated using Advanced Chemistry Development (ACD/Labs) Software V11.02 (© 1994-2012 ACD/Labs)

The above presented experimentally determined or calculated values are in good agreement with the physical chemical information, which is available for the homologues of PFCAs. With increasing chain length the melting and boiling point increase, while no significant change can be found for the vapour pressure and dissociation constant for the C<sub>11-14</sub>-PFCAs based on the calculations given in Table 2. The water solubility decreases with increasing chain length. This is in agreement with the fact that the polarity of the substances decreases with an increasing chain length. It should be emphasised here that it is not possible to assess the calculated values of C<sub>11-14</sub>-PFCAs because there

are factors like special conformation of the molecules which have an influence on the real values, but which have not been taken into account for the calculation.

However, the calculated values have dimensions which would be theoretically expected for the C<sub>11-14</sub> PFCAs.

## **2 HARMONISED CLASSIFICATION AND LABELLING**

C<sub>12</sub>-PFCA is not classified according to Annex VI of Regulation (EC) No 1272/2008 (CLP Regulation).

Twenty-five notifications (3 aggregated notifications) have been submitted to the C&L Inventory. This information is publicly available via the ECHA website at: <http://echa.europa.eu/web/guest/information-on-chemicals/cl-inventory-database>.

### 3 ENVIRONMENTAL FATE PROPERTIES

#### 3.1 Degradation

##### 3.1.1 Abiotic degradation

###### 3.1.1.1 Hydrolysis

There are no studies on the hydrolysis of C<sub>12</sub>-PFCA available. Based on the data given in Annex 1, results of studies of structurally similar substances of the same chemical group could be used to evaluate the hydrolysis of C<sub>11</sub>-PFCA.

Two studies are available on shorter chain lengths PFCAs. Hydrolysis of perfluorinated octanoic acid (C<sub>8</sub>-PFCA; PFOA) and its ammonium salt (APFO) (CAS-No: 335-67-1, 3825-26-1) and perfluorinated nonanoic acid (C<sub>9</sub>-PFCA; PFNA) (CAS No: 375-95-1) were analyzed. The studies are summarized in the following:

C<sub>8</sub>-PFCA is hydrolytically stable under relevant environmental conditions. One study has been discussed in the OECD SIDS Initial Assessment Report for C<sub>8</sub>-PFCA (PFOA), which has been copied here in italic letters (OECD, 2006):

*The 3M Environmental Laboratory performed a study of the hydrolysis of APFO (3M Co., 2001a) (Reliability = 1). The study procedures were based on USEPA's OPPTS Guideline Document 835.2110; although the procedures do not fulfil all the requirements of the guideline, they were more than adequate for these studies. Results were based on the observed concentrations of APFO in buffered aqueous solutions as a function of time. The chosen analytical technique was high performance liquid chromatography with mass spectrometry detection (HPLC-MS).*

*During the study, samples were prepared and examined at six different pH levels from 1.5 to 11.0 over a period of 109 days. Experiments were performed at 50 °C and the results extrapolated to 25 °C. Data from two of the pH levels (3.0 and 11) failed to meet the data quality objective and were rejected. Also rejected were the data obtained for pH 1.5 because ion pairing led to artificially low concentrations for all the incubation periods. The results for the remaining pH levels (5.0, 7.0, and 9.0) indicated no clear dependence of the degradation rate of PFOA on pH. From the data pooled over the three pH levels, it was estimated that the hydrolytic half-life of PFOA at 25°C is greater than 92 years, with the most likely value of 235 years. From the mean value and precision of PFOA concentrations, it was estimated the hydrolytic half-life of PFOA to be greater than 97 years.*

A newer study showed no decomposition of C<sub>8,9</sub>-PFCAs in hot water in absence of S<sub>2</sub>O<sub>8</sub><sup>2-</sup>. After the addition of S<sub>2</sub>O<sub>8</sub><sup>2-</sup> to the reaction system efficient decomposition of PFCAs has been observed at 80 °C. After a reaction time of 6 hours, C<sub>8</sub>-PFCA and C<sub>9</sub>-PFCA were decomposed completely. The reaction products were mainly F<sup>-</sup> and CO<sub>2</sub> at a yield of 77.5 % ((moles of F<sup>-</sup> formed)/(moles of fluorine content in initial PFOA)) and 70.2 % ((moles of CO<sub>2</sub> formed)/(moles of carbon content in initial PFOA)), respectively for C<sub>8</sub>-PFCA. For C<sub>9</sub>-PFCA the reaction products were mainly F<sup>-</sup> and CO<sub>2</sub> at a yield of 88.9 % and 75.2 %, respectively. Short chain PFCAs were a minor reaction product. However, at higher temperatures (150°C) 12.3% of the initial C<sub>8</sub>-PFCA remained and the yields of F<sup>-</sup> and CO<sub>2</sub> were 24.6 and 37.0 %, respectively (Hori et al., 2008) (Reliability = 2).

The water solubility of C<sub>12</sub>-PFCA is lower than those of C<sub>8,9</sub>-PFCA, which can be explained by the expanded fluorinated carbon chain (Annex 1). However, the stability of the PFCAs is mainly based on the stability of the highly fluorinated carbon chain (Siegemund et al., 2000). Since C<sub>8</sub>-PFCA is hydrolytically stable, we estimate a comparable hydrolytically stability also for C<sub>12</sub>-PFCA.

Based on the read across rationale described in Annex 1, data on C<sub>8</sub>-PFCA is used as evidence for C<sub>12</sub>-PFCA to conclude that it is hydrolytically stable under environmental conditions.

### 3.1.1.2 Phototransformation/photolysis

Direct photolysis of a carbon fluorine chain is expected to be very slow, with stability expected to be sustained for more than 1000 years (Environment Canada, 2010).

#### 3.1.1.2.1 Phototransformation in air

There are no studies on phototransformation for C<sub>12</sub>-PFCA in air available. However, studies on C<sub>8</sub>-PFCA exist and are summarized below:

The following information was copied from the OECD SIDS Initial Assessment Report for C<sub>8</sub>-PFCA (PFOA) (OECD, 2006):

*Hurley et al. determined the rate constants of the reactions of OH radicals with a homologous series of perfluorinated acids (from trifluoroacetic acid to nonafluoropentanoic acid) in 700 Torr of air at 296 K (Hurley et al., 2004). For C<sub>3</sub> to C<sub>5</sub> chain length had no discernible impact on the reactivity of the molecule. The rate constant  $k(\text{OH} + \text{F}(\text{CF}_2)_n\text{COOH}) = (1.69 \pm 0.22) \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  for  $n = 2, 3, 4$ , respectively. Atmospheric lifetimes of  $\text{F}(\text{CF}_2)_n\text{COOH}$  with respect to reaction with OH radicals are estimated to be approximately 230 days for  $n = 1$  and 130 days for  $n > 1$ . (Calculation of lifetime by comparison with  $\text{CH}_3\text{CCl}_3$  (half-life 5.99 years,  $k = 1.0 \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ). The authors conclude, that the major atmospheric loss mechanism of perfluorinated carboxylic acids is dry and wet (particle mediated) deposition which occur on a time scale which is probably of the order of 10 days. Reaction with OH is a minor atmospheric loss mechanism for perfluorinated carboxylic acids*

#### 3.1.1.2.2 Phototransformation in water

There are no studies on phototransformation in water for C<sub>12</sub>-PFCA available. However, studies on C<sub>8-11</sub>-PFCA exist and are summarized below:

The photochemical decomposition of long-chain PFCAs in water by use of persulfate ion ( $\text{S}_2\text{O}_8^{2-}$ ) in water (C<sub>9</sub>-PFCA) and in an aqueous/liquid CO<sub>2</sub> biphasic system (C<sub>9-11</sub>-PFCAs) was examined by Hori et al. (Hori et al., 2005b) (Reliability = 2). In water and in the absence of  $\text{S}_2\text{O}_8^{2-}$  (direct photolysis) C<sub>9</sub>-PFCA decomposition of 64.5 % was determined. In the presence of  $\text{S}_2\text{O}_8^{2-}$  the decomposition increased to 100%. The decompositions after 12 hours in the biphasic system were 100% for C<sub>9</sub>-PFCA and C<sub>10</sub>-PFCA, and 77.1% for C<sub>11</sub>-PFCA. The reaction product was mainly F<sup>-</sup> (66.2 %, 73.4 % and 46.35 % of (moles of F<sup>-</sup> formed)/(moles of fluorine content in initial PFCA)) and the minor reaction products were short-chain PFCAs. Since the conditions in this study are not environmentally relevant, we did not describe this study in detail.

In addition to the study of Hori et al, further studies are available for C<sub>8</sub>-PFCA (PFOA) and its ammonium salt APFO (see table 3).

**Table 3: Summary of photodegradation studies for C<sub>8</sub>-PFCA and its ammonium salt**

Test Substance	Result	Remarks	Reliability	Reference
Ammonium salt of C <sub>8</sub> -PFCA	No photodegradation	Direct photolysis	2	(OECD, 2006);(3M Co., 1979)
Ammonium salt of C <sub>8</sub> -PFCA	No photodegradation	Direct and indirect (H <sub>2</sub> O <sub>2</sub> ; synthetic humic water, Fe <sub>2</sub> O <sub>3</sub> ) photolysis	1	(OECD, 2006);(3M Co., 2001b)
	Estimated half-life > 349 days	Indirect photolysis (Fe <sub>2</sub> O <sub>3</sub> )		
C <sub>8</sub> -PFCA		Short wave length (<300 nm) used for irradiation → limited relevance for an aqueous environment	2	(Hori et al., 2004)
	44.9% of the initial PFOA was decomposed after 24 hours	Direct photolysis; 0.48 MPa O <sub>2</sub>		
	35.5% of the initial PFOA was decomposed after 24 hours	Indirect photolysis (H <sub>2</sub> O <sub>2</sub> ); 0.48 MPa O <sub>2</sub>		
	100% of the initial PFOA was decomposed after 24 hours	Indirect photolysis (tungstic heteropolyacid photocatalyst); 0.48 MPa O <sub>2</sub>		
C <sub>8</sub> -PFCA		Short wave length (<300 nm) used for irradiation → limited relevance for an aqueous environment	2	(Hori et al., 2005a)
	16.8% of the initial PFOA was decomposed after 4 hours	Direct photolysis; 0.48 MPa O <sub>2</sub>		
	100% of the initial PFOA was decomposed after 4 hours	Indirect photolysis (S <sub>2</sub> O <sub>8</sub> <sup>2-</sup> ); 0.48 MPa O <sub>2</sub>		

The following information was copied from the OECD SIDS Initial Assessment Report for C<sub>8</sub>-PFCA (PFOA) (OECD, 2006):

*Direct photolysis of APFO was examined in two separate studies (3M Co., 1979; 3M Co., 2001b) and photodegradation was not observed in either study. In the 3M (1979) study, a solution of 50 mg/l APFO in 2.8 litres of distilled water was exposed to simulated sunlight at 22±2 °C. Spectral energy was characterized from 290-600 nm with a max output at ~360 nm. Direct photolysis of the test substance was not detected.*

*In the 3M (3M Co., 2001b) study, both direct and indirect photolysis were examined utilizing techniques based on USEPA and OECD guidance documents. To determine the potential for direct photolysis, APFO was dissolved in pH 7 buffered water and exposed to simulated sunlight. For indirect photolysis, APFO was dissolved in 3 separate matrices and exposed to simulated sunlight for periods of time from 69.5 to 164 hours. These exposures tested how each matrix would affect the photodegradation of APFO. One matrix was a pH 7 buffered aqueous solution containing H<sub>2</sub>O<sub>2</sub> as a well-characterized source of OH radicals. This tested the propensity of APFO to undergo indirect photolysis. The second matrix contained Fe<sub>2</sub>O<sub>3</sub> in water that has been shown to generate hydroxyl radicals via a Fenton-type reaction in the presence of natural and artificial sunlight. The third matrix contained a standard solution of humic material. Neither direct nor indirect photolysis of APFO was observed based on loss of starting material. Predicted degradation products were not detected above their limits of quantitation. There was no conclusive evidence of direct or indirect photolysis whose rates of degradation are highly dependent on the experimental conditions. Using the iron oxide (Fe<sub>2</sub>O<sub>3</sub>) photoinitiator matrix model, the APFO half-life was estimated to be greater than 349 days.*

*According to Hori et al., aqueous solutions of PFOA absorb light strongly from the deep UV-region to 220 nm (Hori et al., 2004). A weak, broad absorption band reaches from 220 to 270 nm (no absorption coefficient stated). The irradiation of a 1.35 mM PFOA solution (29.6 µmol) in water (under 0.48 MPa of oxygen) with light from a xenon-mercury lamp (no radiant flux stated) for 24 hours resulted in a ca. 44.9 % reduction (13.3 µmol) of PFOA concentration. Concentrations of CO<sub>2</sub> and fluoride increased simultaneously. Small amounts (0.1-5 µmol) of short chain perfluorinated hydrocarbon acids (C<sub>2</sub>-C<sub>7</sub>) were detected. The addition of the photocatalyst tungsten heteropolyacid ([PW<sub>12</sub>O<sub>40</sub>]<sup>-</sup>) or persulfate (S<sub>2</sub>O<sub>8</sub><sup>2-</sup>) (Hori et al., 2005a) accelerates the reaction rate. Due to the short wave length used for irradiation (< 300 nm) the photodegradation described may be of limited relevance for an aqueous environment but may be used as a technical process.*

### **3.1.1.2.3 Phototransformation in soil**

#### **3.1.1.3 Summary and discussion on abiotic degradation**

In general, the perfluorinated carboxylates are very stable. Since there are no degradation studies (under relevant environmental conditions) on the C<sub>12</sub>-PFCA available, data from similar substances need to be considered and discussed. Based on the data given in Annex 1, results of studies of structurally similar substances of the same chemical group could be used to evaluate the abiotic degradation of C<sub>12</sub>-PFCA.

The data on C<sub>8</sub>-PFCA indicate that abiotic degradation in the atmosphere is expected to be slow (atmospheric lifetime = 130 days; conclusion by analogy from short-chain perfluorinated acids). Under relevant environmental conditions C<sub>8</sub>-PFCA is hydrolytically stable (DT<sub>50</sub> > 92 years) and do



not undergo direct photodegradation in natural waters. The estimated DT<sub>50</sub> for indirect photolysis is 349 days.

Based on the read across rationale described in Annex 1, data on C<sub>8</sub>-PFCA are used as evidence for C<sub>12</sub>-PFCA to conclude that it is stable under environmental conditions and abiotic degradation is expected to be as low as for the chemically similar substance C<sub>8</sub>-PFCA.

### 3.1.2 Biodegradation

#### 3.1.2.1 Biodegradation in water

##### 3.1.2.1.1 Estimated data

##### 3.1.2.1.2 Screening tests

There is one study available for the C<sub>12</sub>-PFCA (Table 5). In a 28 day ready biodegradability test (OECD 301 C) using 100 mg/L C<sub>12</sub>-PFCA, and 30 mg/L activated sludge non-biodegradability was demonstrated. For C<sub>12</sub>-PFCA no biodegradation was observed (National Institute of Technology and Evaluation, 2007).

**Table 4: Screening tests for C<sub>12</sub>-PFCA.**

Test substance	Method	Result	Reliability	Reference
C <sub>12</sub> -PFCA	OECD 301 C	minus 16% in 28 days	2	(National Institute of Technology and Evaluation, 2002)

In summary, on the basis of the available screening test C<sub>12</sub> PFCA is not readily biodegradable.

##### 3.1.2.1.3 Simulation tests

For C<sub>12</sub>-PFCA no experimental degradation test is available. Based on the data given in Annex 1, results of studies of structurally similar substances of the same chemical group could be used to evaluate the hydrolysis of C<sub>12</sub>-PFCA.

Therefore, test results for C<sub>8</sub>-PFCA are discussed shortly in the following.

No environmental half-lives for C<sub>8</sub>-PFCA have been reported, even in the cases where corresponding tests have been performed (see table 5).

**Table 5: Summary of simulations tests of C<sub>8</sub>-PFCA (PFOA) and its ammonium salt (APFO) and sodium salt**

Test substance	Method	Result	Reliability	Reference
C <sub>8</sub> -PFCA	Closed-loop systems in laboratory scale;  Aerobic and anaerobic conditions	No elimination	3	(Meesters and Schroeder, 2004; Schröder, 2003)
Ammonium salt of C <sub>8</sub> -PFCA	Biodegradation in mixed bacterial culture and activated sludge  Aerobic conditions	< 0.6 % of <sup>14</sup> CO <sub>2</sub> was detected after 28 days	4	(Wang et al., 2005)
Sodium salt of C <sub>8</sub> -PFCA	Microcosm study  Aerobic conditions	No significant dissipation from water column after 35 days (initial concentration 0.3 mg/L; 1mg/L; 30 mg/L)  32% dissipation in 35 days (initial concentration 100 mg/L)	3	(Hanson et al., 2005)
C <sub>8</sub> -PFCA/ ammonium salt of C <sub>8</sub> -PFCA	1.Preliminary screening:  C <sub>8</sub> -PFCA serves as an electron acceptor under anaerobic conditions (in combination with different inocula)  2. Hypothesis refinement:  <sup>14</sup> C C <sub>8</sub> -PFCA serves as an electron acceptor under anaerobic conditions	No significant consumption of the initial C <sub>8</sub> -PFCA during 110 – 259 days  No loss of ammonium salt of C <sub>8</sub> -PFCA No production of <sup>14</sup> CO <sub>2</sub> No detection of radiolabel transformation products	2	(Liou et al., 2010)

In the OECD SIDS Initial Assessment Report it was concluded that C<sub>8</sub>-PFCA (PFOA) is not expected to undergo biodegradation (OECD, 2006). The following text in italic letters was copied from there:

*Schroeder (2003), and Meesters and Schroeder (2004) investigated the biochemical degradation of PFOA in sewage sludge in laboratory scale reactors. After 25 days under aerobic conditions PFOA (initial concentration 5 mg/l) was not eliminated by metabolic processes, mineralization processes or by adsorption (Meesters and Schroeder, 2004; Schröder, 2003).*

Wang et al. studied the biodegradation of fluorotelomer alcohols. However, <sup>14</sup>C-labelled C<sub>8</sub>-PFCA ammonium salt was used as starting material in this study, too. The authors analyzed the headspace of sealed vessels containing mixed bacterial cultures and vessels containing activated sludge from a domestic sewage treatment plant under continuous air flow. The mixed bacterial culture from industrial wastewater treatment sludge was enriched using 8:2 telomere alcohol and <sup>14</sup>C-labelled C<sub>8</sub>-PFCA ammonium salt, respectively. However, for using C<sub>8</sub>-PFCA ammonium salt as a starting material no detailed information are available from the report. The authors describe that potential biodegradation products were separated and quantified by LC/ARC (on-line liquid chromatography/accurate radioisotope counting). Transformation products were identified by quadrupole time of flight mass spectrometry. Only <0.6 % of <sup>14</sup>CO<sub>2</sub> was detected after 28 days. The report contains no graphs or further data to re-evaluate this statement. Although the study seems to be very well documented for <sup>14</sup>C labelled 8:2 FTOH, we can only flag the study with a reliability of 3-4, since details on C<sub>8</sub>-PFCA ammonium salt are not available. The documentation for the results obtained with C<sub>8</sub>-PFCA is missing in the report. However the result indicates that C<sub>8</sub>-PFCA ammonium salt is not biodegradable within 28 days (Wang et al., 2005).

Hanson et al. performed a microcosm study. Microcosms were approximately 1.2 m deep with a water depth of 1 m, a diameter of 3.9 m, and a surface area of 11.95 m<sup>2</sup>. Each microcosm had a capacity of approximately 12m<sup>3</sup> of water. Sediment consisted of a 1:1:1 mixture of sand, loam and organic matter (mainly composted manure). The total carbon content of the sediment was 16.3%. Microcosms were circulated for 2 weeks from a well-fed irrigation pond prior to the experiments. Nominal concentrations of 0.3, 1, 30, and 100 mg/L C<sub>8</sub>-PFCA, as the sodium salt, plus controls were added to the microcosms. Each exposure was randomly assigned to three separate microcosms from a total of 15 microcosms. Immediately prior to treatment, water flow into each microcosm from the main irrigation pond ceased, creating a closed system relative to the other microcosms and the irrigation pond.

Water chemistry and PFOA analysis were taken at the same time on a regularly basis. Temperature and dissolved oxygen content were measured daily. Water samples were collected with a metal integrated water column sampler. Integrated subsamples from at least 4 randomly selected locations in each microcosm were collected to a total volume of 4 L. Samples were stored at 4 °C until analysis. Water samples were analyzed by ion chromatography. The mobile phase was 0.5 mM NaOH, 5 % methanol, and 5% acetonitrile with a flow rate of 0.4 mL/min. Injection volumes varied from 5, 10, 75, and 200 µl for the 100, 30, 1 and 0.3 mg/L microcosms, respectively. For each set of samples analyzed five standards and one quality control sample were included at the beginning of each run and again at the end. Radioactive labelling was not performed. Over a 35-day field study C<sub>8</sub>-PFCA showed no significant dissipation from the water column. However, at the highest concentration (100 mg/L) a partitioning from the water column into other compartments is suspected (32% dissipation in 35 days) (Hanson et al., 2005). Since the documentation of the procedure was insufficient in our opinion the study is not reliable (reliability 3).

Liou et al. investigated the anaerobic biodegradability of C<sub>8</sub>-PFCA respectively its ammonium salt. In a two-phase experiment (preliminary screening, hypothesis refinement) the use of C<sub>8</sub>-PFCA as a

physiological electron acceptor (electron donor: acetate, lactate, ethanol or hydrogen gas) was studied. Additionally, the possibility of co-metabolism of C<sub>8</sub>-PFCA during reductive dechlorination of trichloroethene and during various physiological conditions (aerobic, nitrate-reducing, iron-reducing, sulfate reducing, and methanogenic) was analyzed. Five different inoculums were used (from a municipal waste-water treatment plant, industrial site sediment, an agricultural soil, and soils from two fire training areas). Environmental samples used as inoculum sources in the biodegradation experiments were aseptically gathered (sterile spatula) placed in 0.5 L sterilized canning jars (filled to the brim), stored on ice in the field, and maintained at 4 °C before being transferred to an anaerobic hood where samples were degassed and dispensed as slurries in biodegradation assays. Soils and sludges were gathered from: the Ithaca sewage treatment plant; a water-saturated drainage ditch adjacent to the DuPont Chambers Works waste treatment facility in Salem County, New Jersey, previously shown to carry out reductive dechlorination (Fung et al., 2009); the Cornell agricultural field station (Collamer silt loam, Ithaca, NY), the Ithaca fire training facility, and the Rochester, NY fire training facility (the latter two sites were chosen due to potential contamination with fluorinated fire retardant chemicals) (Liou et al., 2010).

For the serum bottle –based biodegradation assays treatments occurred in triplicates (160 ml serum bottles with 100 mL of media; live ± C<sub>8</sub>-PFCA and abiotic controls, autoclaved for 1 h). For the <sup>14</sup>C-PFOA experiments, 15-mL serum bottles were utilized (50% O<sub>2</sub>-free N<sub>2</sub> headspace, 50% inoculated anaerobic test medium) with non-radioactive C<sub>8</sub>-PFCA and <sup>14</sup>C- C<sub>8</sub>-PFCA (4.5 ICi/mL test medium) to give a final concentration of 100 mg/L C<sub>8</sub>-PFCA medium. For establishing the various terminal electron-accepting processes, a standard anaerobic procedure was used. The anaerobic mineral salts buffer (plus vitamins and trace minerals) was used as diluents for the various inoculums sources (5% wt/volume) with addition of electron donors (10 mM sodium acetate ± 40 mM sodium lactate or 0.6 mM ethanol or 2 atm H<sub>2</sub>) or electron acceptors [O<sub>2</sub> as air headspace or O<sub>2</sub>- free N<sub>2</sub> headspace in each serum bottle with additions of 30 mM nitrate or 4 mg/mL FeOOH or 10 mM sulfate or 0.4 mM trichloroethene (TCE) or no addition (for the methanogenic treatment)]. Samples (1.0 mL) were periodically removed from each serum bottle, placed in 4-mL glass vials sealed with Al-backed caps, immediately mixed with an equal volume of methanol and stored at -20 °C until analysis. Accumulated batches of samples from serum vials were analyzed for concentrations of PFOA, <sup>14</sup>C- C<sub>8</sub>-PFCA, fluoride, nitrate, sulfate, and potential C<sub>8</sub>-PFCA transformation products. Headspace gases were sampled with a gas-tight syringe (250 mL) and analyzed for TCE, vinyl chloride and methane. In the radiotracer study, dissolved <sup>14</sup>C activity in the anaerobic medium and in the 0.4 N KOH solution retrieved from the internal reservoir to trap <sup>14</sup>CO<sub>2</sub> were determined by scintillation counting. To assay potential microbial inhibition by C<sub>8</sub>-PFCA, triplicate serum- bottle assays inoculated with 5% Ithaca sewage were prepared, as above. Anaerobic preparations (±100 ppm C<sub>8</sub>-PFCA) were assayed for methanogenesis. Aerobic preparations containing 15 ppm naphthalene were sampled as above and analyzed by high-performance liquid chromatography (HPLC). After filtration through nylon acrodisc filters, naphthalene was separated at room temperature. Methanol–water (1:1) was the mobile phase at a flow rate of 1.5 mL/ min. The eluent was monitored by UV VIS at 340 nm. Quantification was done by comparison to authentic standards (Liou et al., 2010). C<sub>8</sub>-PFCA quantification was performed by LC/MS/MS following a standard procedure. Potential C<sub>8</sub>-PFCA metabolites were screened by applying LC/MS.

In no combination of the inoculum source, electron donor or physiological conditions a significant percentage of the initial C<sub>8</sub>-PFCA (100 ppm and 100 ppb) was consumed (110 - 259 days). In a test with <sup>14</sup>C labelled C<sub>8</sub>-PFCA ammonium salt (inoculum = sewage), no loss of C<sub>8</sub>-PFCA ammonium salt was detected, no <sup>14</sup>CO<sub>2</sub> was produced and no radiolabelled C<sub>8</sub>-PFCA ammonium salt transformation product was indicated. Co-metabolism of C<sub>8</sub>-PFCA during reductive dechlorination of trichloroethene was suggested by a drop in C<sub>8</sub>-PFCA concentration in the 100 ppb treatment after

a 65-d incubation. However, extensive analysis failed to determine corroborating transformation products. In summary, under conditions which were examined in this study, C<sub>8</sub>-PFCA is environmentally persistent (Liou et al., 2010).

In conclusion, one non-standard aerobic degradation simulation study and one non-standard anaerobic degradation simulation study on C<sub>8</sub>-PFCA demonstrate the high persistence of the compound. Based on the read across rationale described in Annex 1, data on C<sub>8</sub>-PFCAs can be used as evidence of persistence for C<sub>12</sub>-PFCA.

### 3.1.2.2 Biodegradation in sediments

### 3.1.2.3 Biodegradation in soil

There are no degradation studies on C<sub>12</sub>-PFCA available. Based on the data given in Annex 1, results of studies of structurally similar substances of the same chemical group could be used to evaluate the hydrolysis of C<sub>12</sub>-PFCA. A number of studies are available for C<sub>8</sub>-PFCA (PFOA) which were already discussed in the OECD SIDS Initial Assessment Report. The following text was copied from there (italic letters) (OECD, 2006):

*Moody and Field (1999) conducted sampling and analysis of samples taken from groundwater 1 to 3 meters below the soil surface in close proximity to two fire-training areas with a history of aqueous film forming foam use. Perfluorooctanoate was detected at maximum concentrations ranging from 116 to 6750 µg/l at the two sites many years after its use at those sites had been discontinued. These results suggest that PFOA can leach to groundwater (Moody and Field, 1999).*

*Extensive site specific monitoring of soil and ground water concentrations of PFOA and related substances was conducted by 3M, DuPont Daikin and others. PFOA in soil has been shown to persist for decades and to be a long term source of groundwater and surface water contamination (see for example (DuPont Co., 2003; 3M Co., 2005)).*

*At the DuPont Washington Works site soil contaminated by perfluorochemical waste has been shown to contain ppm levels of PFOA 3 decades after application ceased. The underlying groundwater also contains ppm levels of PFOA (DuPont Co., 1999).*

*Extensive field monitoring data generated by 3M at the Decatur, AL site have also shown that PFOA is persistent in soils. Soil samples were collected from a former sludge application area of the 3M Decatur, AL facility also show soil contamination and underlying groundwater contamination up to ppm levels decades after application ceased.*

Moody et al. investigated groundwater at a former fire-training area at Wurtsmith Air Force Base which was used between 1950s and 1993. Before sampling, the soil and groundwater in the area has been studied in detail. Groundwater samples were collected from two types of monitoring wells. All samples were collected in high density polypropylene bottles. Samples were shipped on ice without preservation and stored at 4 °C prior to analysis. Perfluorocarboxylate concentrations were measured as described in the following: Strong anion exchange disks were used to extract perfluorocarboxylates (6 to 8 carbons) from groundwater. The perfluorocarboxylates were simultaneously eluted from the disks and derivatized to their methyl esters by treatment with iodomethane for direct analysis by electron impact gas chromatography-mass spectrometry (GC-MS). A single analysis was conducted for each groundwater sample. The detection limit (defined as a signal-to-noise ratio greater than 3) and quantification limit (defined as a signal-to-noise ratio

greater than 10) for perfluorocarboxylates were 3 and 13 mg/ L, respectively, using 2-chlorolepidine as the internal standard. Additionally, electron capture negative chemical ionization GC-MS was employed to confirm the identity of PFOA, in groundwater samples (Moody et al., 2003). Depending on the location of sampling, the concentrations of C<sub>8</sub>-PFCA were between 8 and 105 µg/L in groundwater. The authors estimated that perfluorinated surfactants had been in the groundwater for at least five years and possibly for as long as 15 years. This showed that C<sub>8</sub>-PFCA did not degrade under the environmental conditions at this site (for both soil and groundwater) (Reliability = 2) (Moody et al., 2003).

The anaerobic biodegradability of C<sub>8</sub>-PFCA and its ammonium salt, respectively, in soil from two fire training areas was investigated by Liou et al. (see above 3.1.2.1.3 Simulation tests). No significant amount of the initial PFOA was dissipated after 259 days.

In conclusion, the available data on C<sub>8</sub>-PFCA demonstrate the high persistence of the compound. Based on the read across rationale described in Annex 1, data on C<sub>8</sub>-PFCAs can be used as evidence of persistence for C<sub>12</sub>-PFCA.

#### **3.1.2.4 Summary and discussion on biodegradation**

A screening study proves that C<sub>12</sub>-PFCA is not readily biodegradable.

Results from non standard degradation studies of C<sub>8</sub>-PFCA, used for read-across approach as described in Annex1, indicate that the structurally similar compound is not biodegradable. The results of one non-standard aerobic biodegradation simulation test, one non-standard anaerobic biodegradation simulation test and field monitoring data on C<sub>8</sub>-PFCA from contaminated sites provide evidence that biodegradation in water, soil and sediment occurs. Since the stability of PFCAs is in general mainly based on the stability of the fluorinated carbon chain it can be concluded that also for C<sub>12</sub>-PFCA no biodegradation in water, soil and sediment can be expected. Thus, it can be assumed that C<sub>12</sub>-PFCA is not biodegradable.

#### **3.1.3 Summary and discussion on degradation**

In conclusion, for C<sub>12</sub>-PFCA there is only one study available showing that the substance is not readily biodegradable. Therefore, data from chemically similar compounds should be considered in a read-across approach (please see Annex 1 for further details). The degradation potential of substances differing only in the number of carbons in the fluorinated carbon chain has been analyzed in some studies. Generally, it is known that the bond between carbon and fluorine is one of the most stable ones in organic chemistry.

A number of studies for the shorter chain C<sub>8</sub>-PFCA show that this substance is very persistent and does not undergo abiotic or biotic degradation at all under environmental conditions. Moreover, also for the C<sub>9</sub>-PFCA, it was shown that no readily biodegradation occurs. Phototransformation was analysed for C<sub>8-11</sub>-PFCA (please see the Data Matrix in Annex 1). However, the test was not obtained under relevant environmental conditions. The persistence is mainly based on the common structural feature: the fluorinated carbon chain.

##### Abiotic degradation

The data on C<sub>8</sub>-PFCA indicate that abiotic degradation in the atmosphere is expected to be slow (atmospheric lifetime = 130 days). The hydrolytic half-life of C<sub>8</sub>-PFCA at 25°C is greater than 92 years, with the most likely value of 235 years under relevant environmental conditions (*3M Co.*,

2001a). No photodegradation of C<sub>8</sub>-PFCA has been observed in studies conducted under relevant environmental conditions. The estimated DT<sub>50</sub> for indirect photolysis is 349 days.

### Biotic degradation

A standard screening tests is available for C<sub>12</sub>-PFCA. No biodegradation at all has been observed within 28 days. For C<sub>8</sub>-PFCA test results differ from “no biodegratation” to 13% biodegradation of the ammonium salt. Thus, it can be concluded that C<sub>8,12</sub>-PFCAs are not readily biodegradable.

For C<sub>8</sub>-PFCA a non-standard aerobic biodegradation simulation test, one non-standard anaerobic biodegradation simulation test and field monitoring data from contaminated sites provide evidence that no biodegradation in water, soil and sediment occurs.

### Conclusion

PFCAs are synthetic compounds which contain a structural feature: a perfluorinated carbon chain combined with a carboxylic group. The chemical structure of these compounds differs only in the number of perfluorinated carbons in the carbon chain.

The stability of organic fluorine compounds has been described in detail by Siegemund et al., 2000: When all valences of a carbon chain are satisfied by fluorine, the zig-zag-shaped carbon skeleton is twisted out of its plane in the form of a helix. This situation allows the electronegative fluorine substituents to envelope the carbon skeleton completely and shield it from chemical attack. Several other properties of the carbon-fluorine bond contribute to the fact that highly fluorinated alkanes are the most stable organic compounds. These include polarizability and high bond energies, which increase with increasing substitution by fluorine. The influence of fluorine is greatest in highly fluorinated and perfluorinated compounds. Properties that are exploited commercially include high thermal and chemical stability (Siegemund et al., 2000).

Based on their molecular properties it is, thus, not surprising, that researchers could not measure degradation of the intensively studied C<sub>8</sub>-PFCA or its salts. Considering the organic chemistry of this substance group it seems to be very likely that a carbon chain being some CF<sub>2</sub>-groups longer is as persistent as a shorter chain C<sub>8</sub>-PFCAs. We therefore conclude that C<sub>9-14</sub>-PFCAs are as resistant to degradation as it has been shown for C<sub>8</sub>-PFCA.

In summary, using the described read across approach, we conclude that C<sub>12</sub>-PFCA is a very persistent synthetic compound which is resistant to abiotic and biotic degradation.

## **3.2 Environmental distribution**

### **3.2.1 Adsorption/desorption**

*Not relevant for this dossier.*

### **3.2.2 Volatilisation**

*Not relevant for this dossier.*

### 3.2.3 Distribution modelling

*Not relevant for this dossier.*

## 3.3 Bioaccumulation

### 3.3.1 Aquatic bioaccumulation

#### 3.3.1.1 Bioconcentration factor BCF

Bioconcentration is the process by which a chemical is accumulated by an organism as a result of exposure to the chemical in water – it often refers to a condition usually achieved under laboratory and steady state conditions. The BCF is typically calculated as the ratio of the measured chemical concentrations in the organism and the water once a steady state has been achieved:

$$BCF = \frac{C_{Biota}}{C_{Water}}$$

The BCF can alternatively be determined kinetically by using the uptake rate  $k_1$  and the depuration rate  $k_2$ :

$$BCF = \frac{k_1}{k_2}$$

There are two studies available which determined the BCFs of C<sub>12</sub>-PFCA.

In the first study carp were exposed to C<sub>12</sub>-PFCA (National Institute of Technology and Evaluation, 2007). The test was conducted in accordance with the OECD 305 guideline this means the test was conducted in a flow through test system, the concentration of the test substance was analytically checked and the pH was within the range 6.0 to 8.5. The uptake period was 60 days. The depuration period was 45 days. Steady state BCFs (whole body) were in the range from 10000 - 16000 for C<sub>12</sub>-PFCA (Table 6). This laboratory study is reliable (reliability 2).

In the second study rainbow trout were exposed in a flow-through system for 12 days followed by a depuration time of 33 days in fresh water to determine tissue distribution and bioconcentration (Martin et al., 2003a). For determination of bioconcentration, juvenile fish (5-10g) were exposed simultaneously to PFCAs of varying chain length. No adverse effects were observable based on fish mortality, growth and liver somatic index. The exposure concentration of each PFCA was analytically checked. PFCA concentrations were stable throughout the uptake phase. There was an initial decrease between 0.25 h and 24 h which is considered to be caused by the rapid uptake of the PFCAs. The mean waterborne concentrations were 0.20 µg/L for C<sub>12</sub>-PFCA. The concentration was relatively stable. A direct analysis was possible as the concentration was above the limit of detection. At 7 occasions during uptake period and 9 occasions during depuration phase, three fish from the exposure tank and one fish from the control were removed to determine the kinetics of uptake and depuration. The BCFs were determined on the basis of the uptake and depuration kinetics and results are given in Table 6. All tissue concentrations were corrected for growth dilution. Additionally, for the tissue distribution study, four immature trout (30 – 48 g) were exposed in separate tanks but under the same uptake conditions (Martin et al., 2003a).



This tissue distribution study showed that unlike lipophilic organic compounds PFCA did not preferentially accumulate in adipose tissue. Hence a lipid-normalisation of the BCFs would not be reasonable. C<sub>11</sub>-C<sub>14</sub>-PFCA concentrations were highest in blood, kidney, liver and gall bladder and low in the gonads, adipose and muscle tissue. Within the blood, the plasma contained between 94 – 99 % of PFCA, with only a minor fraction detectable in the cellular fraction. Recovery from hearts and spleen was low (<10%). Based on high blood, liver and gall bladder concentrations and slow depuration the authors assume that PFCA enter the enterohepatic recirculation in fish. That means the compounds are continuously transferred between the different organs (Martin et al., 2003a).

BCFs were calculated for different body compartments. Though, bioaccumulation should preferably be based on whole body. According to the authors carcass BCFs closely approximate the whole-body BCF. However, compartment-specific BCFs can be more relevant if there is a potential for direct organ-specific toxicity. PFCA cause hepatomegaly in rodents (Kudo et al., 2000) which is an indicator for hepatotoxicity. Thus, from a toxicological perspective, BCFs based on concentrations in individual organs, such as the liver, may be more relevant when the potential for direct organ-specific toxicity (i.e., liver toxicity) is being predicted. No statistically significant difference was found between the liver somatic index of exposed and control fish. However, bioaccumulation tests are not designed for showing toxic effects. The calculated kinetic BCFs are summarized in Table 6.

In a recent study, BCFs have been calculated on the basis of BMF data (Inoue et al., 2012). A recently published comparison of BCFs and biomagnification factors (BMFs) investigated 9 substances in a laboratory fish feeding study with carp (Inoue et al. 2012). Five substances showed BCFs larger than 5000 but only two of these substances were likely to biomagnify. Based on linear regression conducted with their data the authors suggest that a BMF of 0.31 indicates a high bioaccumulation potential. Therefore, we used this approach in a similar way to calculate further BCFs based on BMFs of a study by Martin et al (Martin et al., 2003b) (please find details of the study in section 3.3.1.2).

The calculated BCFs from BMFs of the fish feeding study (Martin et al., 2003b) are smaller than the measured BCF (Martin et al., 2003a) but in a similar order of magnitude. Deviations from the measured BCFs may be due to an erroneous regression. Though, according to the authors the results of the study are highly suggestive. More data would be necessary to support their findings. On the other hand the elimination rates were different in the two studies of Martin et al. (Martin et al., 2003a; Martin et al., 2003b), which may be due to the different fish size and may explain the differences.

There are also other mechanistic models available to roughly estimate a kinetic bioconcentration factor ( $BCF_{Kin}$ ) from data generated in the dietary study. These models depend on physical-chemical input parameters such as the log  $K_{OW}$ . As this parameter cannot be sufficiently estimated for PFCA these approaches should not be used (Weisbrod et al., 2009).

**Table 6 Bioconcentration factors (BCF) of C<sub>12</sub>-PFCA.**

Species/foodweb	BCF	Reliability	Reference
Rainbow trout (carcass)	18000 ± 2700	2	(Martin et al., 2003a)
Rainbow trout (blood)	40000 ± 4500		
Rainbow trout (liver)	18000 ± 2900		
Carp (whole)	10000-16000	2	(National Institute of Technology and Evaluation, 2007)
Juvenile rainbow trout (Carcass)	5761-7327*	3	(Inoue et al., 2012; Martin et al., 2003b)

\* Calculated BCFs based on BMF values for C<sub>12</sub>-PFCA (0.43±0.062). The BMFs were measured by Martin et al. 2003b (please find details of the study in section 3.3.1.2) and the calculation was done as suggested by (Inoue et al., 2012).

#### Conclusion:

All reported BCFs are above 5000 indicating a high bioaccumulation potential of the C<sub>12</sub>-PFCA.

#### **3.3.1.2 Biomagnification factors (BMFs)**

Besides bioconcentration also biomagnification describes the potential of a chemical to bioaccumulate. Biomagnification factors (BMFs) can be measured in the laboratory in a fashion similar to that used in the OECD and US-EPA bioconcentration test protocols. Organisms are exposed to a chemical preliminary via diet. The BMF test typically includes an uptake phase, where levels of chemicals are followed over time, ideally until the chemical concentration in the organism no longer changes with time (i.e., reaching the steady-state). If a steady-state cannot be reached in the experiment, the uptake phase is followed by a depuration phase where organisms are exposed to uncontaminated food. The rate of decline in chemical concentration over time measured in the depuration phase can then be used to derive the chemical uptake rate from which a hypothetical steady-state concentration can be estimated (Conder et al., 2011).

The laboratory-derived dietary BMF is calculated using the ratio of the chemical concentrations in the test animals at steady-state and their diet:

$$BMF = \frac{C_{biota}}{C_{diet}}$$

where chemical concentration in the organism (C<sub>biota</sub>) and its diet (C<sub>diet</sub>) are appropriately normalized, if needed, (e.g., lipid- or protein-normalized) (Conder et al., 2011).

BMF values based on field studies are based on the ratio of the concentration in the predator and the prey:

$$BMF_{(field)} = \frac{C_{predator}}{C_{prey}}$$

Problems arise with increasing body size of predators because analysis is based on tissue or serum samples. Whole-body analysis is not feasible for ethical reasons, i.e. a whole whale would be needed, and due to the challenging logistics with respect to sampling and laboratory constraints. Therefore, some of the derived BMF-values are restricted to certain tissue samples rather than whole body samples. BMF values based on liver samples may be over estimative. From a toxicological perspective concentrations in individual organs, such as the liver, may be more relevant when the potential for direct organ-specific toxicity (i.e., liver toxicity) is being predicted. Whole body values may be estimated if the tissue mass fraction is known for the organism regarded. There may however be some uncertainties due to inter individual and geographical differences (Houde et al., 2006).

Martin et al. (2003b) exposed juvenile rainbow trout (*Oncorhynchus mykiss*) for 34 days to PFCAs in the diet, followed by a 41 day depuration period. Though, the authors describe their results as BAF the results of this study should rather be assigned as BMFs according to the above mentioned definition as uptake only derived from the diet. During the uptake period, animals were daily fed with spiked food at a rate of 1.5 % food per body weight. Spiked food concentrations were 1.1 µg/g for C<sub>12</sub>-PFCA. Water samples collected before and after feeding revealed no traces of PFCAs in water. At 6 occasions during uptake period and during depuration period, fish were removed to determine the kinetics of uptake and depuration. The authors estimated the steady state to less than 34 days. Carcass and liver concentrations were determined by using liquid chromatography-tandem mass spectrometry, and kinetic rates were calculated to determine bioaccumulation parameters. Bioaccumulation (carcass) increased with increasing chain length but was not larger than one: 0.43 ± 0.062 for C<sub>12</sub>-PFCA (see also Table 68). This indicates that a dietary exposure will not result in biomagnification in juvenile trout. The authors assume that the lack of observed biomagnification was likely due to the small size of fish used in the study, resulting in more rapid chemical elimination to water, relative to body size and and that their natural feeding rate is too low. This more rapid chemical elimination would reduce the BMF stronger than what would be observed for larger species or size classes (Martin et al., 2003b).

Furthermore BMFs were estimated from field studies. Studies are described below and results are shown in Table 7.

Transfer of PFCAs was elucidated in Lake Ontario including one 4-membered pelagic food chain (Martin et al., 2004). Whole body samples were collected. Two macroinvertebrates (*Diporeia* and *Mysis*) were considered as primary prey whereas rainbow trout inhabited the top predator's position. Due to the inherent uncertainties correlated with constitution of diet 4 individual combinations of rainbow trout and its prey were regarded. As this study was conducted with fish uptake of PFCAs may not have occurred exclusively over diet but also over the gills. Thus, the factors may be more accurately addressed as BAF. A striking finding of this study was the unexpectedly high content of PFCAs in the benthic invertebrate *Diporeia* occupying the lowest trophic level. The mechanism leading to this exceptional accumulation still needs to be unravelled. The author's hypothesis is that sediments are a major source for PFCAs. Results are given in Table 7.

Tomy et al. also investigated beluga whale, ringed seal, fish pelagic amphipod and arctic copepod of the Western Canadian Arctic. As the authors state themselves differences in sampling years may influence the interpretation of the food web transfer. On the other hand the Arctic as a remote area may be less prone to temporal changes and the existence of point sources there is unlikely. The

derived BMF-values (see Table 11) are restricted to the liver and the resulting BMF may be over estimative. From a toxicological perspective concentrations in individual organs, such as the liver, may be more relevant when the potential for direct organ-specific toxicity (i.e., liver toxicity) is being predicted (Tomy et al., 2009).

Houde et al. examined PFCA serum concentrations in bottlenose dolphins at two different habitats. The authors claim that utilization of serum or liver concentrations of dolphins will overestimate the BMF by a factor of 10 - 30. Whole body concentrations were estimated on the basis of tissue distribution. Samples were collected between 2002 and 2004. Unfortunately, concentrations in other representative fish species originated from different years, thus, entailing additional uncertainty when assessing BMF through the food chain. On the other hand it may be assumed that media and biota were continuously exposed to PFCA in this area throughout the years. The results are summarized in Table 8 (Houde et al., 2006).

Butt et al. conducted a study in the Canadian Arctic. Ringed seal liver samples were provided by local hunters from 11 different locations. The age of the animals was determined via tooth aging and for a few samples the age was estimated using length-age correlations. Stable isotope analysis was done with  $^{15}\text{N}$  to  $^{14}\text{N}$  and  $^{13}\text{C}$  to  $^{12}\text{C}$ . Based on liver samples from polar bears obtained from another study and ringed seal data measured in this study BMFs were calculated (see Table 8). The polar bear sample sites were associated with ringed seal populations. However, the sample collection year for ringed seal populations varied from 2002 to 2005, and it is possible that interpretation of spatial trends may be confounded by temporal variations of PFCA concentration within seal populations (Butt et al., 2008).

Various predator prey relationships in the Westerschelde (Netherlands) were investigated by van Heuvel-Greve and co-workers. Samples of harbour seal plasma and whole body samples of herring, seabass and flounder as well as zooplankton were collected in 2007 and 2008. The trophic level was estimated based on stable isotope ( $^{15}\text{N}$ ) analysis. BMFs were considerable for harbour seal as well as for the sediment dwelling flounder (Environment Canada Health Canada, 2010; van den Heuvel-Greve et al., 2009).

**Table 7: Available BMF data of C<sub>12</sub>-PFCA; if not indicated otherwise BMFs refer to whole body**

Location	Species/foodweb	BMF	Reference	Reliability
Canadian Arctic	Polar bear (liver)/ ringed seal (liver)	2.3-3.6	Butt et al. 2008	2
US, South Carolina, Charleston	Seatrou /pinfish	0.1	Houde et al. 2000	2
US, South Carolina	Dolphin/striped mullet	0.2		
US, South Carolina	Dolphin/pinfish	0.1		
US, South Carolina	Dolphin/red drum	0.4		
US, South Carolina	Dolphin / atlantic croaker	1.8		
US, South Carolina	Dolphin / spotfish	0.6		
US, South Carolina	Dolphin / seatrout	0.6		
US Sarasota Bay, Florida	striped mullet / zooplankton	89		

ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC

US Sarasota Bay, Florida	Pigfish/ zooplankton	2.5		
US Sarasota Bay, Florida	Sheephead/ zooplankton	156		
US Sarasota Bay, Florida	Pinfish/ zooplankton	2.5		
US Sarasota Bay, Florida	Seatrout/ zooplankton	35		
US Sarasota Bay, Florida	Seatrout/striped mullet	0.4		
US Sarasota Bay, Florida	Seatrout/pigfish	14		
US Sarasota Bay, Florida	Seatrout / sheephead	0.2		
US Sarasota Bay, Florida	Seatrout/pinfish	14		
US Sarasota Bay, Florida	Dolphin/striped mullet	0.1		
US Sarasota Bay, Florida	dolphin /pigfish	2.0		
US Sarasota Bay, Florida	Dolphin/ sheephead	0.0		
US Sarasota Bay, Florida	dolphin/pinfish	2.0		
US Sarasota Bay, Florida	Dolphin/seatrout	0.1		
Western Canadian Arctic	Ringed seal (liver)/ arctic cod (liver)	0.1	Tomy et al. 2009	2
Western Canadian Arctic	Beluga whale (liver)/ arctic cod (liver)	3.2		
Western Canadian Arctic	Beluga whale (liver)/ Pacific herring (liver)	7.9		
Western Canadian Arctic	Beluga whale (liver)/ arctic cisco (liver)	4.0		
Western Canadian Arctic	Arctic cod (liver)/ marine arctic copepod (whole body)	1.2		
Western Canadian Arctic	Arctic cod (liver)/ marine pelagic amphipod (whole body)	1.3		
Lake Ontario	Lake trout /alewife	1.9	Martin et al. 2004	2
Lake Ontario	Lake trout/ smelt	1.0		
Lake Ontario	Lake trout / sculpin	0.28		
Lake Ontario	Lake trout / prey (weighted)	1.6		
Laboratory	juvenile rainbow trout(Carcass)	0.43±0.062	Martin et al. 2003b	2

Conclusion:

The biomagnification potential of C<sub>12</sub>-PFCAs was investigated in several field studies and one laboratory study. Field studies investigating the biomagnifications potential between different

predator/prey-relationships showed BMFs well above one indicating biomagnification. Biomagnification was greater in homeotherms than in poikilotherms. Especially for dolphin, beluga whale, polar bear, arctic cod and seatrout, sheephead, and striped mullet BMFs greater than one have been reported.

### 3.3.1.3 Trophic magnification factors (TMFs)

The trophic magnification factor (TMF) is a measure to evaluate biomagnification occurring in food webs. In the Guidance Document on Information Requirements, Chapter R.7.10.1.1, TMF is defined as the concentration increase in organisms with an increase of one trophic level. Again, a TMF greater than one indicates accumulation within the food chain. As already discussed in the BMF chapter sample collection is often restricted to tissue or serum samples with increasing body size of predators due to ethical reasons and due to the challenging logistics with respect to sampling and laboratory constraints.

Martin et al. examined PFCA contents in the food web from Lake Ontario in Canada (Martin et al., 2004). Adult lake trout (top predator) were collected at various years and locations in Lake Ontario. Samples of prey fish (sculpin, smelts and alewife) and macroinvertebrates (*Mysis sp.*, *Diporeia sp.*) were collected at one location in October 2002. Lake trout samples analyzed represented individual whole fish homogenates. The other species were processed as composites of whole individuals. TMFs are shown in Table 8.

Houde et al. investigated the food web of bottlenose dolphins. The authors sampled different biota, i.e. croaker, pinfish, spotfish, spotted seatrout, striped mullet and samples from bottlenose dolphins, as well as water and surface sediment. Sample collection was conducted between 2002 and 2004. Based on stable isotope ( $^{15}\text{N}$ ) analysis the trophic level of each biota sample was determined. PFCAs were analysed in plasma and liver of dolphins and afterwards a whole body burden was calculated. For prey whole body homogenates were analysed for PFCA (Houde et al., 2006). For results see Table 8.

Kelly et al. measured PFCAs in the Canadian Arctic marine food web. The authors used concentrations in sediment and in different organisms (lichens, macroalgae, bivalves, fish, seaducks, and marine mammals) to calculate TMFs. Sample collection was conducted between 1999 and 2003. PFCAs were measured in different tissues/fluids of the beluga whale including blood, muscle liver, milk and also in fetuses. The authors could show that PFCAs especially accumulate in protein rich compartments such as blood and liver and that the TMF of PFCAs correlates with the partitioning behaviour between protein and water and protein and air. Comparisons of different food webs show that the TMF is below one in the case of piscivorous food webs if air breathing organisms are excluded but becomes larger than one if air breathing organisms are taken into account (see Table 8) (Kelly et al., 2009).

Loi et al. investigated a subtropical food web in a nature reserve including phytoplankton, zooplankton, gastropod, worm and shrimp, and liver samples of fish and water bird. Samples were collected between 2008 and 2010. Surface water and sediment samples were collected concurrently with the biota samples. BSAF were calculated based on the assumption that sediment was the major exposure pathway for worms. The study investigated PFCAs with different chain length.  $\text{C}_{12}$ -PFCA was detected in sediment only (Loi et al., 2011).

**Table 8: Available Trophic Magnification Factors (TMF) for C<sub>12</sub> PFCA; if not indicated otherwise TMFs refer to whole body.**

Location	Species/foodweb	TMF	Reference	Reliability
US, South Carolina	Dolphin plasma croaker, pinfish, spotfish, spotted seatrout	0.7 ± 0.8	Houde et al. 2006	2
US, South Carolina,	Whole dolphin burden	0.6 ± 0.8		
Hudson Bay (north-eastern Canada	Sediment/ macroalgae/ bivalves/ fish/ seaduck/ beluga whale	3.89 - 5.87 2.34 - 3.76 (protein corrected)	Kelly et al. 2009	2
Hudson Bay (north-eastern Canada	Sediment/ macroalgae/ bivalves/ fish	0.71 - 1.44 (protein corrected)		
Lake Ontario	Diporeia/slimy sculpin	No significant association with trophic level	Martin et al. 2004	2
Lake Ontario	Mysis/alewife/rainbow smelt/lake trout	No significant association with trophic level		
Mai Po Marshes Nature Reserve in Hong Kong	Tidal shrimp pond brackish food web	1.38	Loi et al., 2011	2

**Conclusion:**

A number of field studies are available which analyzed the trophic magnification potential of C<sub>12</sub>-PFCA. For food chains of beluga whale and in a subtropical pelagic food web TMFs greater than one have been reported, indicating trophic biomagnification. Trophical magnification was greater if the food chain contained homeotherms. TMFs were smaller in the case of piscivorous food webs and if air breathing organisms are excluded but became larger if air breathing organisms were taken into account (Kelly et al., 2009). Thus, C<sub>12</sub>-PFCA biomagnifies in some food webs analyzed within the food chain.

#### **3.3.1.4 Terrestrial bioaccumulation**

Müller et al. conducted a terrestrial food web study consisting of lichen and plants as well as tissue samples (liver muscle and kidney) from caribou, and wolves from two remote northern areas in Canada. Some samples are not from the same season. This food web is considered as relatively well documented example (Kelly and Gobas, 2003). The study illustrates a considerable carry over between plants and caribou. Caribou are a major human food source in numerous arctic communities. This food-chain may also be considered comparable to the pasture-cow food-chain in temperate regions. The results of the study, BMFs as well as TMFs are shown in Table 9.

Tissue concentrations and whole body concentrations were used for calculations. Tissue based BMFs differ considerably. Therefore it is concluded that BMFs based on whole body concentrations are more appropriate (Müller et al., 2011).



**Table 9: BMF and TMF values for C<sub>12</sub>-PFCA for a terrestrial food chain (Müller et al., 2011).**

	Food chain	Location	
		Porcupine	Bathurst
BMF	Caribou (whole)/ lichen	2.9	11
BMF	Caribou(whole)/ lichen	4.5	8
BMF	Wolf (whole)/ caribou (whole)	1.2	1.4
TMF	Wolf (whole)/ caribou (whole)/ lichen	1.4	2.2
TMF	Wolf (whole)/ caribou (whole) / vegetation	1.3	2.0

Conclusion:

The terrestrial BMF and TMF of C<sub>12</sub>-PFCAs are greater than one for the remote Arctic food chain lichen – caribou – wolf, indicating trophic biomagnification.

**3.3.2 Summary and discussion of bioaccumulation**

For C<sub>12</sub>-PFCA reported BCFs are well above the trigger value of 5000 according to REACH Annex XIII. Thus, it can be concluded that C<sub>12</sub>-PFCA is a very bioaccumulative substances. Moreover, BMF and TMF- values greater than one have been reported for a number of food chains, indicating a potential for biomagnification.

#### **4 HUMAN HEALTH HAZARD ASSESSMENT**

*Not relevant for this dossier.*

## **5 ENVIRONMENTAL HAZARD ASSESSMENT**

*Not relevant for this dossier.*

## 6 CONCLUSIONS ON THE SVHC PROPERTIES

### 6.1 PBT, vPvB assessment

#### 6.1.1 Assessment of PBT/vPvB properties – comparison with the criteria of Annex XIII

##### 6.1.1.1 Persistence

For C<sub>12</sub>-PFCA there is only one study available showing that the substance is not readily biodegradable. Further data on the degradability of this substance are not available. Therefore, data from chemically similar compounds are considered in a read-across approach (please see Annex 1 for further details). The degradation potential of substances differing only in the number of carbons in the perfluorinated carbon chain has been analyzed in some studies which indicate that long chain PFCAs are resistant to degradation in the environment.

##### Abiotic degradation

The data on C<sub>8</sub>-PFCA indicate that abiotic degradation in the atmosphere is expected to be slow (atmospheric lifetime = 130 days). The hydrolytic half-life of C<sub>8</sub>-PFCA at 25°C is greater than 92 years, with the most likely value of 235 years under relevant environmental conditions (*3M Co., 2001a*). No photodegradation of C<sub>8</sub>-PFCA has been observed in studies conducted under relevant environmental conditions. The estimated DT<sub>50</sub> for indirect photolysis is 349 days.

##### Biotic degradation

A standard screening test is available for C<sub>12</sub>-PFCA. No biodegradation at all has been detected for within 28 days. For C<sub>8</sub>-PFCA test results differ from “no biodegradation” to 13% biodegradation of the ammonium salt. Thus, it can be concluded that C<sub>8,12</sub>-PFCAs are not readily biodegradable.

For C<sub>8</sub>-PFCA a non-standard aerobic biodegradation simulation test, one non-standard anaerobic biodegradation simulation test and field monitoring data from contaminated sites provide evidence that no biodegradation in water, soil and sediment occurs.

##### Conclusion

PFCAs are synthetic compounds which contain a structural feature: a perfluorinated carbon chain combined with a carboxylic group. The perfluorinated carbon chain is a synthetic feature, there are no natural sources known. The chemical structure of these compounds differs only in the number of perfluorinated carbons in the carbon chain.

The stability of organic fluorine compounds has been described in detail by Siegemund et al., 2000: When all valences of a carbon chain are satisfied by fluorine, the zig-zag-shaped carbon skeleton is twisted out of its plane in the form of a helix. This situation allows the electronegative fluorine substituents to envelope the carbon skeleton completely and shield it from chemical attack. Several other properties of the carbon-fluorine bond contribute to the fact that highly fluorinated alkanes are the most stable organic compounds. These include polarizability and high bond energies, which increase with increasing substitution by fluorine. The influence of fluorine is greatest in highly fluorinated and perfluorinated compounds. Properties that are exploited commercially include high thermal and chemical stability (Siegemund et al., 2000).

Comparing the physico-chemical properties of C<sub>8-14</sub>-PFCAs it becomes obvious that with increasing chain length water solubility decreases and the sorption potential increases. This trend is based on the increasing number of CF<sub>2</sub>-groups in the molecular structure. The molecular reason for the persistence of highly fluorinated chemicals is the shielding effect of the substituted fluorine atoms described by Siegemund et al., 2000. Thus, using the described read across approach, we conclude that C<sub>12</sub>-PFCA is a very persistent synthetic compound which is resistant to abiotic and biotic degradation and fulfils both, the P and the vP criteria of Annex XIII.

#### **6.1.1.2 Bioaccumulation**

Bioaccumulative substances are defined in Annex XIII with a BCF >2000. If substances have a BCF >5000 they fulfil the criteria of being very bioaccumulative.

For C<sub>12</sub>-PFCA reported BCFs are well above the trigger value of 5000 according to REACH Annex XIII. Thus, it can be concluded on the basis of the studies that C<sub>12</sub>-PFCA is a very bioaccumulative substances. Moreover, BMF and TMF values greater than one have been reported for a number of food chains, indicating a potential for biomagnification.

#### **6.1.1.3 Toxicity**

*Not relevant for this dossier.*

### **6.1.2 Summary and overall conclusions on the PBT, vPvB properties**

C<sub>12</sub>-PFCA is not readily biodegradable. Further degradation studies are not available. Applying the read across approach, data from structurally similar compounds can be used to evaluate the degradation potential of the substance. C<sub>8-14</sub>-PFCAs contain a highly similar chemical structure, a perfluorinated carbon chain and a carboxylic acid group. The compounds differ only in the number of CF<sub>2</sub>-groups.

Comparing the physico-chemical properties of C<sub>8-14</sub>-PFCAs it becomes obvious that with increasing chain length water solubility decreases and the sorption potential increases. This trend is based on the increasing number of CF<sub>2</sub>-groups in the molecular structure. According to the read-across approach these chemicals follow a regular pattern as a result of structural similarity. Those substances may therefore be considered as a group or a category of substances and the read-across approach can be applied.

In general, the persistence of long chain PFCAs can be explained by the shielding effect of the fluorine atoms, blocking e.g. nucleophilic attacks to the carbon chain. High electronegativity, low polarizability and high bond energies make highly fluorinated alkanes to the most stable organic compounds. It is not expected that the substitution of a functional group – the carboxylic group in PFCAs – alters this persistence of these chemicals. This fact is confirmed by a study which obtained a DT<sub>50</sub> of >92 years for C<sub>8</sub>-PFCA in water. Screening studies of C<sub>8,9,12,14</sub>-PFCA showed no biodegradation within 28 days. Non-standard tests with C<sub>8</sub>-PFCA could not detect any degradation products under environmentally relevant conditions. Moreover, a monitoring study showed that C<sub>8</sub>-PFCA remained in soil and groundwater, years after application of fire fighting foam which contained PFCAs.

Therefore, we conclude that C<sub>12</sub>-PFCA is not degraded in the environment and thus fulfils the P- and vP-criteria under REACH.

The available BCF-values of C<sub>12</sub>-PFCA are above 5000. Thus, the B as well as the vB-criteria according to Annex XIII of REACH are fulfilled.

In conclusion, C<sub>12</sub>-PFCA is a vPvB-substance according to Art. 57e) of REACH.

## **6.2 CMR assessment**

*Not relevant for this dossier.*

## **6.3 Substances of equivalent level of concern assessment.**

*Not relevant for this dossier.*

## PART II

### INFORMATION ON USE, EXPOSURE, ALTERNATIVES AND RISKS

#### INFORMATION ON MANUFACTURE, IMPORT/EXPORT AND USES –CONCLUSIONS ON EXPOSURE

Information on environmental exposure of the environment with C<sub>11-14</sub>-PFCAs can be seen from time trends of concentrations in monitoring studies. It has to be kept in mind that concentrations of PFCAs in different environmental media can be caused by degradation of precursors as well. Nevertheless, concentration trends give a hint on the relevance of PFCAs.

A study determined PFCAs in livers of melon-headed whales (*Peponocephala electra*) collected along the coast of Japan, from three mass strandings that occurred during the past 25 years (Hart et al., 2008). Whereas in 1982 C<sub>11,12</sub>-PFCAs were below the limit of detection, they were detected in 2006.

Concentrations of PFCAs were determined in liver of harbour seals (n = 68) collected from the northwest Atlantic between 2000 and 2007 showed an increasing trend (Shaw et al., 2009).

The temporal trends of PFCAs were determined in lake trout collected between 1979 and 2004 from Lake Ontario. From 1998 on concentrations show an increasing trend for C<sub>11-14</sub>-PFCAs (Furdui et al., 2008).

Significant annual increases for C<sub>11,12,13</sub>-PFCAs (5.9%, 8.5% and 5.2%, respectively) were found in polar bears (*Ursus maritimus*) within the period 1984 – 2006 (Dietz et al., 2008).

These increasing trends outside of Europe are supported by two studies performed with samples from the German Specimen Bank. An increasing trend of concentrations of C<sub>11-14</sub>-PFCAs was found in fish samples within the years 1995 – 2010. Fish samples originated from different location and the increasing trend was found for all locations (Theobald et al., 2011). This increasing trend is supported by data from a biomonitoring study with human sera samples from two German cities. These data are not yet published and are therefore confidential. Serum samples cover the period from 1982 – 2009 and an increasing trend of concentrations was found for C<sub>11,12</sub>-PFCAs. This trend is also supported by concentrations in serum samples from a Norwegian study (Haug et al., 2009).

There is evidence that C<sub>11-14</sub>-PFCAs are in use and that exposure of the environment is taking place. Furthermore PFOA and long chain PFCA are used in the production of fluoropolymers and fluorotelomers and as additives and components in consumer and industrial products (Environment Canada Health Canada. 2010b).

#### CURRENT KNOWLEDGE ON ALTERNATIVES

Only little information is available on substitutes. It can be assumed that some substitutes for C<sub>8</sub>-PFCA might also be usable for C<sub>11-14</sub>-PFCAs.

In general, PFCAs with eight carbon atoms can be replaced with shorter chain fluorinated chemicals containing six or less carbon atoms. Non-fluorinated alternatives are available as well, i.e.

propylated aromatics (naphthalene or biphenyls) and aliphatic alcohols (sulphosuccinate and fatty alcohol ethoxylates) (Danish Ministry of the Environment, 2005; van der Putte et al., 2010; Walters and Santillo, 2006). In the table 11 below known C<sub>8</sub>-PFCA alternatives are summarized.

**Table 10: Alternative compounds, their product names, company and use for C<sub>8</sub>-PFCA and its salts.**

Alternative compound	Product name	Company	Used for /Used in	Ref.
PFBS or based on different C <sub>4</sub> -perfluoro-compounds	Novec®	3M	Paint and coatings industry. Electronic coating, industrial and commercial cleaning, cleaner for solder flux residue, degreasing applications	(Poulsen et al., 2005; van der Putte et al., 2010; Walters and Santillo, 2006)
Dodecafluoro-2-methylpentan-3-one(CF <sub>3</sub> -CF <sub>2</sub> -C(O)-CF(CF <sub>3</sub> ) <sub>2</sub> )	Novec®	3M	Fire-fighting fluid	(Poulsen et al., 2005; Walters and Santillo, 2006)
C6-fluorocompounds	Forafac®	DuPont	Fire-fighting foam	(Poulsen et al., 2005)
CF <sub>3</sub> or C <sub>2</sub> F <sub>5</sub> pendant fluoroalkyl polyethers	PolyFox®	OMNOVA Solutions Inc.	Surfactant and flow, level and wetting additive for coating formulations. Also used in floor polish	(Poulsen et al., 2005)
Propylated aromatics (naphthalenes or biphenyls)	Ruetasolv®	Rütgers Kurehe Solvents GmbH	Water repelling agents for rust protection systems, marine paints, coatings, etc.	(Poulsen et al., 2005; Walters and Santillo, 2006)
Aliphatic alcohols (sulphosuccinate and fatty alcohol ethoxylates)	Emulphor®, Lutensit®	BASF	Levelling and wetting agents	(Poulsen et al., 2005)
Sulfosuccinates	EDAPLAN® LA451	Münzing Chemie	Paint and coatings industry: Wetting agents for water based applications,	(Poulsen et al., 2005)



ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC

			e.g. wood primers	
Sulfosuccinate	Hydropal at®875	Cognis	Paint and coating industry: Wetting and dispersing agents	(Poulsen et al., 2005)
Silicone Polymers	WorléeAdd®	Welrée-Chemie	Wetting agents in paint and ink industry	(Poulsen et al., 2005)
Branched fluoro ethers			Can be applied for all products	(van der Putte et al., 2010)
Short-chain fluorinated technologies (six or less carbons)	Capstone	DuPont	Commercially available in home furnishings, fire fighting foam, fluorosurfactants, paper packaging, textiles, stone and tile, and leather end uses	<sup>1</sup>
Ammonium 4,8-dioxa-3H-perfluorononanoate	ADONA	3M	Emulsifier used in the aqueous emulsion polymerization of fluoropolymers made from tetrafluoroethylene (TFE)	(Gordon, 2011)

<sup>1</sup> [http://www.oecd.org/document/34/0,3746,en\\_21571361\\_44787844\\_44799586\\_1\\_1\\_1\\_1,00.html](http://www.oecd.org/document/34/0,3746,en_21571361_44787844_44799586_1_1_1_1,00.html)

## ANNEX I - READ-ACROSS APPROACH

In general, the read-across approach can be applied if substances whose physicochemical and/or toxicological and/or ecotoxicological properties are likely to be similar or follow a regular pattern as a result of structural similarity. Those substances may be considered as a group or a category of substances. According to ECHA's practical guide 6 "How to report read-across and categories" similarities may be due to a common functional group, common precursor or breakdown products, constant pattern in changing potency or common constituents or chemical class.

### Structural similarities of C8-14 PFCAs and some salts

In Table 11 the chemical structures of the C8-14-PFCAs are displayed. All PFCAs contain a carboxylic acid group and a perfluorinated carbon chain. The compounds differ only in the number of carbon atoms within the fluorinated carbon chain. Thus, we conclude, that all the C8-14-PFCAs belong to the same chemical class and contain not only a common functional group but are highly similar according to their chemical structure.

**Table 11: CAS-Numbers and similarity of chemical structures of long chain PFCAs.**

Name	Abbreviation	CAS-No	IUPAC Name	Chemical structure
PFOA	C <sub>8</sub> -PFCA	335-67-1	Octanoic acid, pentadecafluoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>6</sub> -COOH
PFNA	C <sub>9</sub> -PFCA	375-95-1	Nonanoic acid, heptadecafluoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>7</sub> -COOH
PFDA	C <sub>10</sub> -PFCA	335-76-2	Decanoic acid, nonadecafluoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>8</sub> -COOH
PFUnDA	C <sub>11</sub> -PFCA	2058-94-8	Undecanoic acid, heneicosafuoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>9</sub> -COOH
PFDoDA	C <sub>12</sub> -PFCA	307-55-1	Dodecanoic acid, tricosafuoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>10</sub> -COOH
PFTrDA	C <sub>13</sub> -PFCA	72629-94-8	Tridecanoic acid, pentacosafuoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>11</sub> -COOH
PFTeDA	C <sub>14</sub> -PFCA	376-06-7	Tetradecanoic acid, heptacosafuoro-	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>12</sub> -COOH

### Dissociation of C<sub>8-14</sub>-PFCAs and its salts in aqueous media

Under environmental conditions in aqueous media the free perfluorinated carboxylic acids (PFCAs) stay in equilibrium with their conjugate bases, the perfluorinated carboxylates. The fraction of each species depends on the acid dissociation constant ( $pK_a$ ) and the pH of the environmental compartment. Salts of PFCAs, which are sometimes used in laboratory experiments, will be in equilibrium with the corresponding acid in aqueous phases as well. Currently used techniques for analysis and quantification of PFCAs in i.e. environmental samples are not able to distinguish

between both of the species. Therefore, reported concentrations always include the acids as well as the bases. If reported concentrations are used for the determination of bioaccumulation factors or for experiments determining the persistency, aqueous phase concentrations include both species. Experimental determination of  $pK_a$  is difficult for PFCAs, i.e. because of the surface active properties. Calculated values should be taken with care, because for most of the models it is unclear whether PFCAs are within their applicability domain. For assessing the intrinsic properties of the PFCA within this dossier the exact knowledge of the fraction of each species is not required, because both of the species will be available independently from the starting conditions.

Furthermore, due to the uncertainties of  $pK_a$  values it is not wise to calculate partition coefficients under environmental pH conditions. We would like to mention that there is an ongoing scientific discussion showing that the partitioning of PFCAs in the environment can be described by the properties of the neutral PFCAs only (Webster and Ellis 2011).

#### Physicochemical properties and partition coefficients of C<sub>8-14</sub>-PFCAs and some salts

The experimental determination of i.e. partition coefficients is difficult for example because of the surface active properties of the ionic PFCAs. The presence of ionic PFCAs depends on the dissociation of PFCAs in aqueous media. Nevertheless, there are models available, i.e. COSMOtherm calculating partitioning coefficients of neutral PFCAs. Again, whether neutral PFCAs are present in aqueous media depends in the dissociation of the acids. Air-water as well as octanol-water partition coefficients are of course different for PFCAs with 8 to 14 carbon atoms but they show a clear increasing trend with chain length (see **Table 12** below, Wang et al., 2011). This can be explained by the increasing molecular volume with each additional CF<sub>2</sub>-unit. The trend of the fate of PFCAs with chain length is supported by information on sorption of PFCAs on sediment. Sorption increases with increasing chain length (Higgins and Luthy, 2006) also under environmental conditions (Ahrens et al., 2010) (see **Table 12**).

**Table 12: Basic substance information and physical chemical properties relevant to justify read across in the PBT assessment.**

Abbreviation	C <sub>8</sub> -PFCA			C <sub>9</sub> -PFCA	C <sub>10</sub> -PFCA	C <sub>11</sub> -PFCA	C <sub>12</sub> -PFCA	C <sub>13</sub> -PFCA	C <sub>14</sub> -PFCA
Acronym	PFOA	APFO	NaPFO	PFNA	PFDA	PFUnDA	PFDoDA	PFTTrDA	PFTeDA
IUPAC Name	Octanoic acid, pentadecafluoro-	ammonium pentadecafluoro-octanoate	pentadeca octanoic acid sodium salt	Nonanoic acid, heptadecafluoro-	Decanoic acid, nonadecafluoro-	Undecanoic acid, heneicosafuoro-	Dodecanoic acid, tricosafuoro-	Tridecanoic acid, pentacosafuoro-	Tetradecanoic acid, heptacosafuoro-
Chemical Structure	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>6</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>6</sub> -COO-NH <sub>4</sub> <sup>+</sup>	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>6</sub> -COO-Na <sup>+</sup>	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>7</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>8</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>9</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>10</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>11</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>12</sub> -COOH
CAS No	335-67-1	3825-26-1	335-95-5	375-95-1	335-76-2	2058-94-8	307-55-1	72629-94-8	376-06-7
<b>Physico-chemical data</b>									
Molecular Weight g/mol	414.09	431.1		464.08	514.08	564.0909	614.0984	664.1059	714.11
Partitioning Coefficient logKow				2.3 – 2.48 (exp)	2.65 – 2.87 (exp)	3.19 – 3.41	logP 9.363±0.888 at 25°C (calc)	logP 10.093±0.901 at 25 °C (calc)	logP 10.823±0.914 at 25 °C (calc)
	5.30 (calc., COSMOtherm, Wang et al., 2011)			5.9 (calc., COSMOtherm, Wang et al., 2011)	6.5 (calc., COSMOtherm, Wang et al., 2011)	7.2 (calc., COSMOtherm, Wang et al., 2011)	7.8 (calc., COSMOtherm, Wang et al., 2011)	8.25 (calc., COSMOtherm, Wang et al., 2011)	8.90 (calc., COSMOtherm, Wang et al., 2011)
log K <sub>OA</sub>	7.23 (calc., COSMOtherm, Wang et al., 2011)			7.50 (calc., COSMOtherm, Wang et al., 2011)	7.77 (calc., COSMOtherm, Wang et al., 2011)	8.08 (calc., COSMOtherm, Wang et al., 2011)	8.36 (calc., COSMOtherm, Wang et al., 2011)	8.63 (calc., COSMOtherm, Wang et al., 2011)	8.87 (calc., COSMOtherm, Wang et al., 2011)
log K <sub>AW</sub>	-1.93 (calc., COSMOtherm, Wang et al., 2011)			-1.58 (calc., COSMOtherm, Wang et al., 2011)	-1.27 (calc., COSMOtherm, Wang et al., 2011)	-0.92 (calc., COSMOtherm, Wang et al., 2011)	-0.58 (calc., COSMOtherm, Wang et al., 2011)	-0.38 (calc., COSMOtherm, Wang et al., 2011)	0.03 (calc., COSMOtherm, Wang et al., 2011)
Dissociation constant	2.5 2.8 in 50% aqueous ethanol 1.5 - 2.8					0.52±0.10; (calculated)	0.52±0.10 (calculated)	0.52±0.10; (calculated)	0.52±0.10; (calculated)
Partition coefficients log Kd (sediment and overlapping)	0.04 (Ahrens et al., 2010)*			0.6 (Ahrens et al., 2010) *	1.8 (Ahrens et al., 2010) *	3.0 (Ahrens et al., 2010) *			

ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC

Abbreviation	C <sub>8</sub> -PFCA			C <sub>9</sub> -PFCA	C <sub>10</sub> -PFCA	C <sub>11</sub> -PFCA	C <sub>12</sub> -PFCA	C <sub>13</sub> -PFCA	C <sub>14</sub> -PFCA
Acronym	PFOA	APFO	NaPFO	PFNA	PFDA	PFUnDA	PFDoDA	PFTrDA	PFTeDA
dissolved phase)									
Log K <sub>oc</sub> (sediment organic carbon-normalized distribution coefficient)	2.06 (Higgins and Luthy, 2006)# 1.09 (Ahrens et al., 2010) *			2.39 (Higgins and Luthy, 2006) # 2.4 (Ahrens et al., 2010) *	2.76 (Higgins and Luthy, 2006) # 3.6 (Ahrens et al., 2010) *	3.3 (Higgins and Luthy, 2006) # 4.8 (Ahrens et al., 2010) *			
Water solubility	9.5 g/L (25 °C) 4.14 g/L (22 °C)	0.033 mol/L, 14.2 g/L at 2.5 °C (Nielsen 2012)	0.036 mol/L at 8.0 °C at critical micelle concentration (Nielsen 2012)			1.2E-4 g/L; pH 1 at 25 °C 9.0E-4 g/L; pH 2 at 25 °C 8.5E-3 g/L; pH 3 at 25 °C 0.056 g/L; pH 4 at 25 °C 0.14 g/L; pH 5 at 25 °C 0.16 g/L; pH 6-10 at 25 °C (calculated)	2.9E-5 g/L pH 1 at 25 °C 2.2E-4 g/L pH 2 at 25 °C 2.0E-3 g/L pH 3 at 25 °C 0.014 g/L pH 4 at 25 °C 0.034 g/L pH 5 at 25 °C 0.039 g/L pH 6 at 25 °C 0.040 g/L pH 7 at 25 °C 0.041 g/L pH 8-10 at 25 °C (calculated)	7.3E-6 g/L; pH 1 at 25 °C 5.5E-5 g/L; pH 2 at 25 °C 5.1E-4 g/L; pH 3 at 25 °C 3.5E-3 g/L; pH 4 at 25 °C 8.6E-3 g/L; pH 5 at 25 °C 0.0100 g/L; pH 6-10 at 25 °C (calculated)	1.9E-6 g/L; pH 1 at 25 °C 1.4E-5 g/L; pH 2 at 25 °C 1.3E-4 g/L; pH 3 at 25 °C 9.3E-4 g/L; pH 4 at 25 °C 2.2E-3 g/L; pH 5 at 25 °C 2.6E-3 g/L; pH 6-10 at 25 °C (calculated)
Vapour pressure	4.2 Pa (25 °C) for PFOA extrapolated from measured data 2.3Pa (20 °C) for PFOA extrapolated from measured data 128 Pa (59.3 °C) for PFOA measured	0.0081 Pa at 20 °C, calculated from measured data <0.1 hPa at 20 °C 0.012 Pa at 25 °C 0.0028 Pa at 25 °C (Nielsen 2012)				0.6 to 99.97 kPa (112 to 237.7 °C) (calculated)	9.40E-3 Torr at 25 °C (calculated)	3.59E-3 Torr at 25 °C (calculated)	1.37E-3 Torr at 25 °C (calculated)

ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC

Abbreviation	C <sub>8</sub> -PFCA			C <sub>9</sub> -PFCA	C <sub>10</sub> -PFCA	C <sub>11</sub> -PFCA	C <sub>12</sub> -PFCA	C <sub>13</sub> -PFCA	C <sub>14</sub> -PFCA
Acronym	PFOA	APFO	NaPFO	PFNA	PFDA	PFUnDA	PFDoDA	PFTrDA	PFTeDA
<b>Stability</b>									
Phototransformation in water DT50	No photodegradation detected under relevant env. conditions	No photodegradation detected under relevant env. conditions		No photodegradation tested under relevant env. conditions 100 % after 12 h by use of persulfate ion (S2O8 <sup>2-</sup> ) in water	No photodegradation tested under relevant env. Conditions 100 % after 12 h by use of persulfate ion (S2O8 <sup>2-</sup> ) in water	No photodegradation tested under relevant env. Conditions 77% after 12 h by use of persulfate ion (S2O8 <sup>2-</sup> ) in water			
Hydrolysis DT50	>97 yr								
Direct photolysis		No photodegradation							
indirect photolysis		No photodegradation (H <sub>2</sub> O <sub>2</sub> ; synthetic humic water, Fe <sub>2</sub> O <sub>3</sub> )  estimated half-life > 349 days (Fe <sub>2</sub> O <sub>3</sub> )							
ready biodegradability screening test	not readily biodegradable (OECD 301 C,F)	not readily biodegradable (OECD 301 B)		not readily biodegradable (OECD 301 F)			not readily biodegradable (OECD 301 C)		not readily biodegradable (OECD 301 C)
Simulation tests	No elimination by metabolic processes, mineralization or adsorption								
Biodegradation in soil, sediment	No degradation detected								

\* pH of the water samples analyzed 7.1-8.3 Temp.: 15.3 – 17.7 °C

ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC

**Table 13 : Information on BCF, BMF and TMF of C<sub>9-14</sub> PFCAs relevant to justify read across in the B assessment.**

<b>Abbreviation</b>	<b>C<sub>9</sub>-PFCA</b>	<b>C<sub>10</sub>-PFCA</b>	<b>C<sub>11</sub>-PFCA</b>	<b>C<sub>12</sub>-PFCA</b>	<b>C<sub>13</sub>-PFCA</b>	<b>C<sub>14</sub>-PFCA</b>
Acronym	PFNA	PFDA	PFUnDA	PFDoDA	PFTrDA	PFTeDA
IUPAC Name	Nonanoic acid, heptafluoro-	Decanoic acid, nonafluoro-	Undecanoic acid, heneicosfluoro-	Dodecanoic acid, tricosfluoro-	Tridecanoic acid, pentacosfluoro-	Tetradecanoic acid, heptacosfluoro-
Chemical Structure	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>7</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>8</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>9</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>10</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>11</sub> -COOH	CF <sub>3</sub> (CF <sub>2</sub> ) <sub>12</sub> -COOH
CAS No	375-95-1	335-76-2	2058-94-8	307-55-1	72629-94-8	376-06-7
<b>Physico-chemical data</b>						
Molecular Weight g/mol	464.08	514.08	564.0909	614.0984	664.1059	714.11
Partitioning Coefficient log K <sub>OW</sub>	2.3 – 2.48 (exp)	2.65 – 2.87 (exp)		logP 9.363±0.888 at 25°C (calc)	logP 10.093±0.901 at 25 °C (calc)	logP 10.823±0.914 at 25 °C (calc)
	5.9 (calc., COSMOtherm, Wang et al., 2011)	6.5 (calc., COSMOtherm, Wang et al., 2011)	7.2 (calc., COSMOtherm, Wang et al., 2011)	7.8 (calc., COSMOtherm, Wang et al., 2011)	8.25 (calc., COSMOtherm, Wang et al., 2011)	8.90 (calc., COSMOtherm, Wang et al., 2011)
log K <sub>OA</sub>	7.50 (calc., COSMOtherm, Wang et al., 2011)	7.77 (calc., COSMOtherm, Wang et al., 2011)	8.08 (calc., COSMOtherm, Wang et al., 2011)	8.36 (calc., COSMOtherm, Wang et al., 2011)	8.63 (calc., COSMOtherm, Wang et al., 2011)	8.87 (calc., COSMOtherm, Wang et al., 2011)
log K <sub>AW</sub>	-1.58 (calc., COSMOtherm, Wang et al., 2011)	-1.27 (calc., COSMOtherm, Wang et al., 2011)	-0.92 (calc., COSMOtherm, Wang et al., 2011)	-0.58 (calc., COSMOtherm, Wang et al., 2011)	-0.38 (calc., COSMOtherm, Wang et al., 2011)	0.03 (calc., COSMOtherm, Wang et al., 2011)
<b>BCF</b>						
MITI (fish)			2300 - 3700	10000 - 16000		16000 - 17000
Lumbriculus variegatus / sediment	0.64 ± 0.05	0.06 ± 0.04				
Rainbow trout (carcass)			2700 ± 400	18000 ± 2700		23000 ± 5300
Rainbow trout (blood)		2700±350	11000 ± 1400	40000 ± 4500		30000 ± 4200
Rainbow trout (liver)		1100 ± 180	4900 ± 770	18000 ± 2900		30000 ± 6000
Carp (whole)			2300 - 3700	10000-16000		16000 - 17000
Juvenile rainbow trout (carcass)		450 ± 6 62	4044-5132*	5761-7327*		10388-15857*
<b>BMF</b>	0.13-111	0.21-87	0.21 - 353	0.1 – 156	0.35 – 9	0.33 – 8.5
<b>TMF</b>	1.9-7.0	1.5-20	0.75 – 31.2	0.6 – 3.76	1.4 – 2.45	0.23 – 2.6




## ANNEX XV – IDENTIFICATION OF TRICOSAFLUORODODECANOIC ACID AS SVHC


---

\* Calculated BCFs based on BMF values for C<sub>11</sub>-PFCA ( $0.28 \pm 0.04$ ), C<sub>12</sub>-PFCA ( $0.43 \pm 0.062$ ) and C<sub>14</sub>-PFCA ( $1.0 \pm 0.25$ ).

# pH of sediment analysed: 5.7 to 7.6



**Figure 1: BMFs for C<sub>11-13</sub>-PFCAs in a remote terrestrial food chain from two different locations (whole body, Müller et al., 2011). The study is reliable (reliability 2).**



**Figure 2: TMFs for C<sub>11-13</sub>-PFCAs in a remote terrestrial food chain from two different locations (whole-body, Müller et al., 2011). The study is reliable (reliability 2).**



## LITERATURE CITED

3M Co. 1978. *Biodegradation (ABS/LAS Shake Culture Test)*. St. Paul, MN. Project number 9970612613. U.S. Environmental Protection Agency Administrative Record 226-0489.

3M Co. 1979. *Photolysis study using simulated sunlight. FC-143 Photolysis study using simulated sunlight* U.S. Environmental Protection Agency Administrative Record 226-0490.

3M Co. 1980. *Environmental Laboratory Acute Toxicity Testing: FC-12*. U.S. Environmental Protection Agency Administrative Record 226-0504.

3M Co. 1985. *Ready Biodegradation of FX-1001 (BOD/COD)*. Lab Request No. C1006. Environmental Laboratory. St. Paul, MN. U.S. Environmental Protection Agency Administrative Record 226-0494.

3M Co. 2001a. *Hydrolysis Reactions of Perfluorooctanoic Acid (PFOA)*. 3M Lab Request Number E00-1851. U.S. Environmental Protection Agency Administrative Record 226-1030a090.

3M Co. 2001b. *Screening study on the aqueous photolytic degradation of perfluorooctanoic acid [PFOA]*. U.S. Environmental Protection Agency Administrative Record 226-1030a091.

3M Co. 2005. *Analysis of PFBS, PFHS, PFOS and PFOA in Water Samples collected at 3M Guin*. Amended Analytical Report E05-0662. U.S. Environmental Protection Agency Administrative Record 226-3571.

Ahrens L, Taniyasu S, Yeung LWY, Yamashita N, Lam PKS, Ebinghaus R. 2010. *Distribution of polyfluoroalkyl compounds in water, suspended particulate matter and sediment from Tokyo Bay, Japan*. *Chemosphere* 79: 266-272.

Butt CM, Mabury SA, Kwan M, Wang X, Muir DC. 2008 Mar. *Spatial trends of perfluoroalkyl compounds in ringed seals (Phoca hispida) from the Canadian Arctic*. *Environ Toxicol Chem* 27(3):542-553.

Conder JM, Gobas FAPC, Borga K, Muir DCG, Powell DE. 2011. *Use of trophic magnification factors and related measures to characterize bioaccumulation potential of chemicals*. *Integr Environ Assess Manag*:n/a.

Danish Ministry of the Environment. 2005. *More environmentally friendly alternatives to PFOS-compound and PFOA*.

Dietz R, Bossi R, Rigét FF, Sonne C, Born EW. 2008. *Increasing Perfluoroalkyl Contaminants in East Greenland Polar Bears (Ursus maritimus): A New Toxic Threat to the Arctic Bears*. *Environ Sci Technol* 42(7):2701-2707.

DuPont Co. 1997. *Evaluation of the Biodegradability of C-8 Using the Modified Sturm Test (OECD 301 B)*.

DuPont Co. 1999. *RCRA Facility Investigation Report. DuPont Washington Works, West Virginia USEPA Permit Number WVD04-587-5291. Docket # OPPT-2003-0012-0184*.

DuPont Co. 2003. *Surface Water Monitoring Report for Washington Works Facility and Local, Letart and Dry Run Landfills*. Washington, WV. U.S. Environmental Protection Agency Administrative Record 226-1508.

Environment Canada. 2010. *Draft Ecological Screening Assessment Report. Long-Chain (C9-C20) PErfluorocarboxylic Acids, their Salts and their Precursors-*.

Environment Canada Health Canada. 2010. *Draft screening assessment perfluorooctanoic acid , its salts, and its precursors*.

Environment Canada Health Canada. 2010b. *Risk management scope for Perfluorooctanoic Acid (PFOA) management scope for Perfluorooctanoic Acid (PFOA),its Salts, and its Precursors, and long-Chain (C9-C20) Perfluorocarboxylic Acids (PFCAs), their Salts, and their Precursors*.

Furdui VI, Helm PA, Crozier PW, Lucaciu C, Reiner EI, Marvin CH, Whittle DM, Mabury SA, Tomy GT. 2008. *Temporal trends of perfluoroalkyl compounds with isomer analysis in lake trout from Lake Ontario (1979-2004)*. *Environ Sci Technol* 42(13):4739-4744.

Gordon SC. 2011. *Toxicological evaluation of ammonium 4,8-dioxa-3H-perfluorononanoate, a new emulsifier to replace ammonium perfluorooctanoate in fluoropolymer manufacturing*. *Regulatory Toxicology and Pharmacology* 59(1):64-80.

Hanson M, Small J, Sibley P, Boudreau T, Brain R, Mabury S, Solomon K. 2005. *Microcosm Evaluation of the Fate, Toxicity, and Risk to Aquatic Macrophytes from Perfluorooctanoic Acid (PFOA)*. *aect* 49(3):307-316.

Hart K, Kannan K, Isobe T, Takahashi S, Yamada TK, Miyazaki N, Tanabe S. 2008 *Time trends and transplacental transfer of perfluorinated compounds in melon-headed whales stranded along the Japanese coast in 1982, 2001/2002, and 2006*. *Environ Sci Technol* 42(19):7132-7137.

Haug LS, Thomsen C, Becher G. 2009. *Time Trends and the Influence of Age and Gender on Serum Concentrations of Perfluorinated Compounds in Archived Human Samples*. *Environ. Sci. Technol* 43 (6): 2131–2136.

Haukas M, Berger U, Hop H, Gulliksen B, Gabrielsen GW. 2007. *Bioaccumulation of per- and polyfluorinated alkyl substances (PFAS) in selected species from the Barents Sea food web*. *Environ Pollut* 148(1):360-371.

Higgins CP and Luthy RG. 2006, *Sorption of Perfluorinated Surfactants on Sedimentss*. *Environ. Sci. Technol.* 40: 7251-7256.

Hori H, Hayakawa E, Einaga H, Kutsuna S, Koike K, Ibusuki T, Kiatagawa H, Arakawa R. 2004. *Decomposition of environmentally persistent perfluorooctanoic acid in water by photochemical approaches*. *Environ Sci Technol* 38(22):6118-6124.

Hori H, Nagaoka Y, Murayama M, Kutsuna S. 2008. *Efficient decomposition of perfluorocarboxylic acids and alternative fluorochemical surfactants in hot water*. *Environ Sci Technol* 42(19):7438-7443.

Hori H, Yamamoto A, Hayakawa E, Taniyasu S, Yamashita N, Kutsuna S, Kiatagawa H, Arakawa R. 2005a. *Efficient decomposition of environmentally persistent perfluorocarboxylic acids by use of persulfate as a photochemical oxidant*. *Environ Sci Technol* 39(7):2383-2388.

- Hori H, Yamamoto A, Kutsuna S. 2005b. Efficient photochemical decomposition of long-chain perfluorocarboxylic acids by means of an aqueous/liquid CO<sub>2</sub> biphasic system. *Environmental Science and Technology* 39(19):7692-7697.
- Houde M, Bujas TA, Small J, Wells RS, Fair PA, Bossart GD, Solomon KR, Muir DC. 2006. Biomagnification of perfluoroalkyl compounds in the bottlenose dolphin (*Tursiops truncatus*) food web. *Environ Sci Technol* 40(13):4138-4144.
- Hurley MD, Andersen MPS, Wallington TJ, Ellis DA, Martin JW, Mabury SA. 2004. Atmospheric chemistry of perfluorinated carboxylic acids: Reaction with OH radicals and atmospheric lifetimes. *Journal of Physical Chemistry A* 108(4):615-620.
- Inoue Y, Hashizume N, Yoshida T, Murakami H, Suzuki Y, Koga Y, Takeshige R, Kikushima E, Yakata N, Otsuka M. 2012. Comparison of Bioconcentration and Biomagnification Factors for Poorly Water-Soluble Chemicals Using Common Carp (*Cyprinus carpio* L.). *Arch Environ Contam Toxicol*.
- Kelly BC, Gobas FAPC. 2003. An arctic terrestrial food-chain bioaccumulation model for persistent organic pollutants  
1. *Environ Sci Technol* 37(13):2966-2974.
- Kelly BC, Ikonomou MG, Blair JD, Surridge B, Hoover D, Grace R, Gobas FAPC. 2009. Perfluoroalkyl contaminants in an Arctic marine food web: Trophic magnification and wildlife exposure. *Environ Sci Technol* 43(11):4037-4043.
- Kudo N, Bandai N, Suzuki E, Katakura M, Kawashima Y. 2000. Induction by perfluorinated fatty acids with different carbon chain length of peroxisomal beta-oxidation in the liver of rats. *Chem Biol Interact* 124(2):119-132.
- Liou JS, Szostek B, Derito CM, Madsen EL. 2010. Investigating the biodegradability of perfluorooctanoic acid. *Chemosphere* 80(2):176-183.
- Loi EI, Yeung LW, Taniyasu S, Lam PK, Kannan K, Yamashita N. 2011. Trophic magnification of poly- and perfluorinated compounds in a subtropical food web. *Environ Sci Technol* 45(13):5506-5513.
- Martin JW, Mabury SA, Solomon KR, Muir DC. 2003a. Bioconcentration and tissue distribution of perfluorinated acids in rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol Chem* 22(1):196-204.
- Martin JW, Mabury SA, Solomon KR, Muir DC. 2003b. Dietary accumulation of perfluorinated acids in juvenile rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol Chem* 22(1):189-195.
- Martin JW, Whittle DM, Muir DC, Mabury SA. 2004. Perfluoroalkyl contaminants in a food web from Lake Ontario. *Environ Sci Technol* 38(20):5379-5385.
- Meesters RJ, Schroeder HF. 2004. Perfluorooctane sulfonate - a quite mobile anionic anthropogenic surfactant, ubiquitously found in the environment. *Water Sci Technol* 50(5):235-242.
- Moody CA, Field JA. 1999. Determination of perfluorocarboxylates in groundwater impacted by fire-fighting activity. *Environmental Science and Technology* 33(16):2800-2806.

Moody CA, Hebert GN, Strauss SH, Field JA. 2003. Occurrence and persistence of perfluorooctanesulfonate and other perfluorinated surfactants in groundwater at a fire-training area at Wurtsmith Air Force Base, Michigan, USA. *J Environ Monit* 5(2):341-345.

Müller CE, De Silva AO, Small J, Williamson M, Wang X, Morris A, Katz S, Gamberg M, Muir DC. 2011. Biomagnification of Perfluorinated Compounds in a Remote Terrestrial Food Chain: Lichen-Caribou-Wolf. *Environ Sci Technol* 45(20):8665-8673.

National Institute of Technology and Evaluation. 2002. Biodegradation and Bioconcentration of Existing Chemical Substances under the Chemical Substances Control Law. Information on the chemical published in the Official Bulletin of Economy, Trade and Industry.

National Institute of Technology and Evaluation. 2007. Biodegradation and Bioconcentration of Existing Chemical Substances under the Chemical Substances Control Law. Information on the chemical published in the Official Bulletin of Economy, Trade and Industry.

Nielsen CJ. 2012. PFOA Isomers, Salts and Precursors. Literature study and evaluation of physico-chemical properties. Klif project no. 3012013. [www.klif.no/publikasjoner/2944/ta2944.pdf](http://www.klif.no/publikasjoner/2944/ta2944.pdf).

Oakes KD, Sibley PK, Solomon KR, Mabury SA, Van der Kraak GJ. 2004. Impact of perfluorooctanoic acid on fathead minnow (*Pimephales promelas*) fatty acyl-CoA oxidase activity, circulating steroids, and reproduction in outdoor microcosms. *Environ Toxicol Chem* 23(8):1912-1919.

OECD. 2006. SIDS Initial Assessment Report after SIAM 22 - Ammonium Perfluorooctanoate & Perfluorooctanoic Acid. 1-210.

OECD. 2007. SIDS Dossier - Ammonium Perfluorooctanoate & Perfluorooctanoic acid.

Pace Analytical. 1987. Ready Biodegradation of FC-126 (BOD/COD). 3M Company Lab Request No. E1282. Minneapolis, MN. U.S. Environmental Protection Agency Administrative Record 226-0495.

Poulsen PB, Jensen AA, Wallström E. 2005. More environmentally friendly alternatives to PFOS-compounds and PFOA. Danish Ministry of the Environment.

Powley CR, George SW, Russell MH, Hoke RA, Buck RC. 2008. Polyfluorinated chemicals in a spatially and temporally integrated food web in the Western Arctic. *Chemosphere* 70(4):664-672.

Schröder HF. 2003. Determination of fluorinated surfactants and their metabolites in sewage sludge samples by liquid chromatography with mass spectrometry and tandem mass spectrometry after pressurised liquid extraction and separation on fluorine-modified reversed-phase sorbents. *Journal of Chromatography A* 1020(1):131-151.

Shaw S, Berger ML, Brenner D, Tao L, Wu Q, Kannan K. 2009. Specific accumulation of perfluorochemicals in harbor seals (*Phoca vitulina concolor*) from the northwest Atlantic. *Chemosphere* 74(8):1037-1043.

Siegemund G, Schwertfeger W, Feiring A, Smart B, Behr F, Vogel H, McKusick B. 2000. Fluorine compounds, organic. *Ullmann's Encyclopedia of Industrial Chemistry*.

Stasinakis AS, Petalas AV, Mamais D, Thomaidis NS. 2008. Application of the OECD 301F respirometric test for the biodegradability assessment of various potential endocrine disrupting chemicals. *Bioresour Technol* 99(9):3458-3467.

Theobald N, Schäfer S, Baaß A-C. 2011. Perfluorierte Verbindungen in archivierten Fischproben der Umweltprobenbank des Bundes. FKZ 301 02 038, <http://www.umweltprobenbank.de/de/documents/publications/16904>.

Tomy GT, Pleskach K, Ferguson SH, Hare J, Stern G, Macinnis G, Marvin CH, Loseto L. 2009. Trophodynamics of some PFCs and BFRs in a western Canadian Arctic marine food web. *Environmental Science and Technology* 43(11):4076-4081.

van den Heuvel-Greve M, Leonards P, Brasseur S, Kotterman M, Zabel A, Vethaak D. 2009. Bioaccumulation of perfluorinated compounds in a harbour seal food web in the Westerschelde, the Netherlands: a field study. In: Poster presentation at SETAC North America, New Orleans.

van der Putte I, Murin M, van Velthoven M, Affourtit F. 2010. Analysis of the risks arising from the industrial use of Perfluorooctanoic Acid (PFOA) and Ammonium Perfluorooctanoate (APFO) and from their use in consumer articles. Evaluation of the risk reduction measures for potential restrictions on the manufacture, placing on the market and use of PFOA and APFO. European Commission, DG Enterprise and Industry.

Walters A, Santillo D. 2006. Uses of perfluorinated substances. Greenpeace.

Wang N, Szostek B, Buck RC, Folsom PW, Sulecki LM, Capka V, Berti WR, Gannon JT. 2005. Fluorotelomer alcohol biodegradation-direct evidence that perfluorinated carbon chains breakdown. *Environ Sci Technol* 39(19):7516-7528.

Wang Z, MacLeod M, Cousins IT, Scheringer M, Hungerbühler K. 2011. Using COSMOtherm to predict physicochemical properties of poly- and perfluorinated alkyl substances (PFASs). *Environ. Chem.* 8: 389-398.

Webster EM, Ellis DA. 2011. Equilibrium Modeling: A pathway to understanding observed perfluorocarboxylic and perfluorosulfonic acid behavior. *Environmental Toxicology and Chemistry* 30: 2229-2236.

Weisbrod AV, Woodburn KB, Koelmans AA, Parkerton TF, McElroy AE. 2009. Evaluation of bioaccumulation using in vivo laboratory and field studies. *Integr Environ Assess Manag* 5:598-623.