

EN

ANNEX

SUMMARY OF PRODUCT CHARACTERISTICS FOR A BIOCIDAL PRODUCT

FACORAT PASTA

Product type(s)

PT14: Rodenticides

Authorisation number: RO/2019/0128/MRA/IT/2014/00199/AUT

R4BP asset number: RO-0009813-0000

1. ADMINISTRATIVE INFORMATION

1.1. Trade name(s) of the product

Trade name(s)	FACORAT PASTA
---------------	---------------

1.2. Authorisation holder

Name and address of the authorisation holder	Name	ST ARKO IMPORT EXPORT SRL
	Address	Str. Pinteza Viteazu nr. 9 400404 Cluj-Napoca Romania
Authorisation number	RO/2019/0128/MRA/IT/2014/00199/AUT	
<i>R4BP asset number</i>	RO-0009813-0000	
Date of the authorisation	05/02/2014	
Expiry date of the authorisation	30/12/2026	

1.3. Manufacturer(s) of the product

Name of manufacturer	INDUSTRIALCHIMICA SRL
Address of manufacturer	Via Sorgaglia, 40 35020 Arre (PD) Italy
Location of manufacturing sites	Via Sorgaglia, 40 35020 Arre (PD) Italy

1.4. Manufacturer(s) of the active substance(s)

Active substance	Brodifacoum
Name of manufacturer	ACTIVA SRL
Address of manufacturer	Via Feltre 32 20132 Milan Italy
Location of manufacturing sites	Tezza S.r.l-via Tre Ponti 22 37050 S. Maria di Zevio Italy

2. PRODUCT COMPOSITION AND FORMULATION

2.1. Qualitative and quantitative information on the composition of the product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
Brodifacoum	3-[3-(4'-bromobiphenyl-4-yl)-1,2,3,4-tetrahydro-1-naphthyl]-4-hydroxycoumarin	active substance	56073-10-0	259-980-5	0,005 % (w/w)

2.2. Type(s) of formulation

RB Bait (ready for use)

3. HAZARD AND PRECAUTIONARY STATEMENTS

Hazard statements	H360D: May damage the unborn child. H373: May cause damage to organs to BLOOD through prolonged or repeated exposure {1:state route of exposure if it is conclusively proven that no other routes of exposure cause the hazard:}.
Precautionary statements	P102: Keep out of reach of children. P202: Do not handle until all safety precautions have been read and understood. P280: Wear protective gloves. P308+P313: IF exposed or concerned: Get medical advice. P501: Dispose of contents to in accordance with national and international regulations.

4. AUTHORISED USE(S)

4.1. Use description

Table 1. House mice - professionals - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: juveniles to adult
Field(s) of use	indoor use Indoor
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: 40g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): 0 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	industrial ; professional
Pack sizes and packaging material	<p>Bait product of 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.</p> <p>Packaging material: plastic composite-triple layer (polyester/PET met/polyethylene)</p> <p>Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional) Compatibility of the product with the proposed packaging materials (Yes/No) Labelled or Printed Bottle from 1.5 Kg- to 5 kg</p>

	Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled Plastic Bucket from 1.5 Kg -to 25 kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes Labelled Plastic Bucket- with inner plastic liner from 1.5 - to 10 kg - max size inner liner 10 Kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes Labelled or Printed Carton sachet with plastic inner liner from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled or Printed Cardboard with inner plastic liner 1,5-10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated cardboard Professional, trained professional Yes
--	--

4.1.1. Use-specific instructions

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.1.2. Use-specific risk mitigation measures

4.1.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.1.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

4.2. Use description

Table 2. Rats - professionals - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juvenile to adult Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: other: juvenile to adult
Field(s) of use	indoor use Indoor
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: 60-100g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Dilution (%): 0 Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Category(ies) of users	industrial ; professional
Pack sizes and packaging material	Bait product of 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Packaging material: plastic composite-triple layer (polyester/PET met/polyethylene)

Type of packaging
Size/volume of the packaging
Material of the packaging
Type and material of closure(s)
Intended user (e.g. professional, non-professional)
Compatibility of the product with the proposed packaging materials (Yes/No)

Labelled or Printed Bottle
 from 1.5 Kg- to 5 kg
 Plastic composite (HDPE)
 Bottle sealed
 Professional, trained professional
 Yes

Labelled or Printed Bag/Sack
 from 1.5 Kg - to 25 kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes

Labelled or Printed Bag/Sack - with Inner plastic liner
 from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes

Labelled Plastic Bucket
 from 1.5 Kg -to 25 kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes

Labelled Plastic Bucket- with inner plastic liner
 from 1.5 - to 10 kg - max size inner liner 10 Kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes

Labelled or Printed Carton sachet with plastic inner liner
 from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each
 Carton + Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes

Labelled or Printed Cardboard with inner plastic liner
 1,5-10 kg single inner plastic bags of up to 10 Kg each
 Carton + Plastic composite
 Prefabricated cardboard
 Professional, trained professional
 Yes

--	--

4.2.1. Use-specific instructions

The bait stations should be visited at least every 5 to 7 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.2.2. Use-specific risk mitigation measures

4.2.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.2.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

4.3. Use description

Table 3. House mice and rats - professionals -outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juvenile to adult Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: other: juvenile to adult Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: other: juvenile to adult
Field(s) of use	outdoor use outdoor around buildings
Application method(s)	Method: bait application Detailed description: Ready-to-use bait to be used in tamper-resistant bait stations
Application rate(s) and frequency	Application rate: House mouse: 40g of bait per bait point. <i>Rattus</i> : 60-100g of bait per bait point. If more than one bait station is

	<p>needed, the minimum distance between bait stations should be of 5 meters.</p> <p>Dilution (%): 0</p> <p>Number and timing of application: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.</p>
Category(ies) of users	industrial ; professional
Pack sizes and packaging material	<p>Bait product of 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.</p> <p>Packaging material: plastic composite-triple layer (polyester/PET met/polyethylene)</p> <p>Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional) Compatibility of the product with the proposed packaging materials (Yes/No)</p> <p>Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled Plastic Bucket from 1.5 Kg -to 25 kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes</p> <p>Labelled Plastic Bucket- with inner plastic liner</p>

	<p>from 1.5 - to 10 kg - max size inner liner 10 Kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes Labelled or Printed Carton sachet with plastic inner liner from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled or Printed Cardboard with inner plastic liner 1,5-10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated cardboard Professional, trained professional Yes</p>
--	--

4.3.1. Use-specific instructions

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2. Use-specific risk mitigation measures

Do not apply this product directly in the burrows.

4.3.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.3.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months

4.4. Use description

Table 4. House mice and rats - trained professionals - indoor

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	<p>Scientific name: Rattus norvegicus Common name: brown rat Development stage: juvenile to adult</p> <p>Scientific name: Rattus rattus Common name: roof rat Development stage: other: juvenile to adult</p> <p>Scientific name: Mus musculus Common name: house mouse Development stage: other: juvenile to adult</p>
Field(s) of use	<p>indoor use</p> <p>indoor</p>
Application method(s)	<p>Method: bait application</p> <p>Detailed description: Bait formulations:- Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations.</p>
Application rate(s) and frequency	<p>Application rate: Rate:Rats :- High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point.. Mice 40 gr of bait per bait point</p> <p>Dilution (%): 0</p> <p>Number and timing of application: Rate:Rats :- High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point.. Mice 40 gr of bait per bait point</p>
Category(ies) of users	trained professional
Pack sizes and packaging material	<p>Bait product of 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Packaging material: plastic composite-triple layer (polyester/PET met/polyethylene)</p> <p>Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional)</p>

	<p>Compatibility of the product with the proposed packaging materials (Yes/No)</p> <p>Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled Plastic Bucket from 1.5 Kg -to 25 kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes</p> <p>Labelled Plastic Bucket- with inner plastic liner from 1.5 - to 10 kg - max size inner liner 10 Kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes</p> <p>Labelled or Printed Carton sachet with plastic inner liner from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Cardboard with inner plastic liner 1,5-10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated cardboard Professional, trained professional Yes</p>
--	--

4.4.1. Use-specific instructions

Remove the remaining product at the end of treatment period.

4.4.2. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product in pulsed baiting treatments.
- do not use the product as permanent baits for the prevention of rodent infestation or monitoring activities

4.4.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to water drainage system, ensure that bait contact with water is avoided.

4.4.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.4.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section “General directions for use (trained professionals)”, are implemented.

4.5. Use description

Table 5. Mice and rats - trained professionals - outdoor around buildings

Product type	PT14: Rodenticides
Where relevant, an exact description of the authorised use	Not relevant for rodenticides
Target organism(s) (including development stage)	Scientific name: <i>Rattus norvegicus</i> Common name: brown rat Development stage: juvenile to adult Scientific name: <i>Rattus rattus</i> Common name: roof rat Development stage: other: juvenile to adult Scientific name: <i>Mus musculus</i> Common name: house mouse Development stage: other: juvenile to adult
Field(s) of use	outdoor use outdoor around buildings
Application method(s)	Method: bait application

	Detailed description: Bait formulations:- Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations.
Application rate(s) and frequency	<p>Application rate: Rate:Rats :- High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point.. Mice 40 gr of bait per bait point</p> <p>Dilution (%): 0</p> <p>Number and timing of application: Rate:Rats :- High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point.. Mice 40 gr of bait per bait point</p>
Category(ies) of users	trained professional
Pack sizes and packaging material	<p>Bait product of 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Packaging material: plastic composite-triple layer (polyester/PET met/polyethylene)</p> <p>Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional) Compatibility of the product with the proposed packaging materials (Yes/No)</p> <p>Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack from 1.5 Kg - to 25 kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled or Printed Bag/Sack - with Inner plastic liner from 1.5 Kg - to 10 kg - max size inner liner 10 Kg Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes</p> <p>Labelled Plastic Bucket from 1.5 Kg -to 25 kg Labelled plastic (HDPE) bucket</p>

	<p>Rectangular or truncated cone bucket sealed Professional, trained professional Yes Labelled Plastic Bucket- with inner plastic liner from 1.5 - to 10 kg - max size inner liner 10 Kg Labelled plastic (HDPE) bucket Rectangular or truncated cone bucket sealed Professional, trained professional Yes Labelled or Printed Carton sachet with plastic inner liner from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated bags or serial production bags , both thermal welded Professional, trained professional Yes Labelled or Printed Cardboard with inner plastic liner 1,5-10 kg single inner plastic bags of up to 10 Kg each Carton + Plastic composite Prefabricated cardboard Professional, trained professional Yes</p>
--	---

4.5.1. Use-specific instructions

Remove the remaining product at the end of treatment period.

4.5.2. Use-specific risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Do not use the product in pulsed baiting treatments.
- do not use this product directly in the burrows

- do not use the product as permanent baits for the prevention of rodent infestation or monitoring activities

4.5.3. Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.4. Where specific to the use, the instructions for safe disposal of the product and its packaging

4.5.5. Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section “General directions for use (trained professionals)”, are implemented.

5. GENERAL DIRECTIONS FOR USE¹

5.1. Instructions for use

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves (rubber) during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.- Instructions for use that are "bait-specific":- Bait in sachets: Do not open the sachets containing the bait.

¹Instructions for use, risk mitigation measures and other directions for use under this section are valid for any authorised uses.

5.2. Risk mitigation measures

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment unless authorised for permanent baiting treatments.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre
- Hazardous to wildlife.

5.4. Instructions for safe disposal of the product and its packaging

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

6. OTHER INFORMATION

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.