

Rezumatul caracteristicilor produsului pentru un produs biocid

Denumirea produsului: CROPTER(AGRORAT DIFE- 5 PASTA)

Tipul (tipurile) de produs (produse): TP 14 - Rodenticide

TP 14 - Rodenticide

TP 14 - Rodenticide

TP 14 - Rodenticide

TP 14 - Rodenticide

TP 14 - Rodenticide

Numărul curent: RO/2019/0186/MRA/IT 2012/2-00005/AUT

Numărul curent R4BP 3: RO-0016298-0000

Cuprins

Informații administrative	1
1.1. Denumirea comercială a produsului	1
1.2. Detinatorul autorizației	1
1.3. Producătorul (Producătorii) produselor biocide	1
1.4. Producatorul (producatorii) substanței (substanțelor) active	1
2. Compoziția și formularea produsului	2
2.1. Informații calitative și cantitative despre compoziția produsului biocid	2
2.2. Tipul preparatului	2
3. Fraze de pericol și de precauție	2
4. Utilizare (utilizări) autorizată (autorizate)	3
5. Direcții generale de utilizare	18
5.1. Instrucțiuni de utilizare	18
5.2. Măsuri de reducere a riscurilor	19
5.3. Particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului	20
5.4. Instrucțiuni pentru eliminarea în siguranță a produsului și a ambalajului acestuia	20
5.5. Condiții de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare	20
6. Alte informații	20

Informații administrative

1.1. Denumirea comercială a produsului

CROPTER, AGRORAT DIFE- 5 PASTA

1.2. Detinatorul autorizației

Numele și adresa detinatorului autorizației	Numele	Laboratorios Agrochem S.L.
	Adresa	C/Tres Rieres, 10 - Pol. Ind. Sud 08292 Esparreguera Spania
Numărul curent	RO/2019/0186/MRA/IT 2012/2-00005/AUT	
Numărul curent R4BP 3	RO-0016298-0000	
Data autorizației	05/05/2010	
Data de expirare a autorizației	01/07/2024	

1.3. Producătorul (Producătorii) produselor biocide

Numele producatorului	Activa S.r.l.
Adresa producătorului	Via Feltre, 32 20132 Milano Italia
Adresa locurilor de producție	Vebi Istituto Biochimico srl - Via Desman, 43 35010 Borgoricco (PD) Italia
	I.N.D.I.A. Industrie Chimiche srl - Via Sorgaglia, 25 35020 Arre (PD) Italia

1.4. Producatorul (producatorii) substanței (substanțelor) active

Substanța activă	26 - Difenacum
Numele producătorului	ACTIVA SRL
Adresa producătorului	Via Feltre 32 20132 Milan Italia
Adresa locurilor de producție	Tezza S.r.l.-via Tre Ponti 22 37050 S. Maria di Zevio Italia

2. Compoziția și formularea produsului

2.1. Informații calitative și cantitative despre compoziția produsului biocid

Denumirea comună	Denumirea IUPAC	Funcția	Numărul CAS	Numărul CE	Continut (%)
Difenacum	3-(3-biphenyl-4-yl-1,2,3,4-tetrahydro-1-naphthyl)-4-hydroxycoumarin	Substanța activă	56073-07-5	259-978-4	0,005

2.2. Tipul preparatului

RB - Bait (gata de utilizare)

3. Fraze de pericol și de precauție

Fraze de pericol	<p>Poate dăuna fătului.</p> <p>Poate provoca leziuni ale organelor to BLOOD în caz de expunere prelungită sau repetată.</p>
Fraze de precauție	<p>A nu se lăsa la îndemâna copiilor.</p> <p>A nu se manipula decât după ce au fost citite și înțelese toate măsurile de securitate.</p> <p>Purtați mănuși de protecție.</p> <p>ÎN CAZ DE expunere sau de posibilă expunere:Consultați medicul.</p> <p>Aruncați conținutul la in accordance with national and international regulations.</p>

4. Utilizare (utilizări) autorizată (autorizate)

4.1 Modul de utilizare

Utilizarea 1 - House mice - professionals - indoor

Tipul produsului	TP 14 - Rodenticide
Descrierea exactă a utilizării autorizate, după caz	Not relevant for rodenticides
Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)	denumirea științifică: Mus musculus denumirea comună: House mouse etapă de dezvoltare: juveniles to adult
Domeniul de utilizare	de interior Indoor
Metoda (metodele) de aplicare	Metoda: cerere momeală Descriere detaliată: Ready-to-use bait to be used in tamper-resistant bait stations
Doza (dozele) și frecvența de aplicare	Frecvența de aplicare: 40g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Diluare (%): 0 Numărul și calendarul de aplicare: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Categoria (categoriile) de utilizatori	industrial profesional
Dimensiunile ambalajului și materialul de ambalare	Bait product of 10 or 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg. Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional) Compatibility of the product with the proposed packaging materials (Yes/No) Labelled or Printed Bottle from 1.5 Kg- to 5 kg Plastic composite (HDPE) Bottle sealed Professional, trained professional

Yes
Labelled or Printed Bag/Sack
from 1.5 Kg - to 25 kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled or Printed Bag/Sack - with Inner plastic liner
from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled Plastic Bucket
from 1.5 Kg -to 25 kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes
Labelled Plastic Bucket- with inner plastic liner
from 1.5 - to 10 kg - max size inner liner 10 Kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes
Labelled or Printed Carton sachet with plastic inner liner
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled or Printed Cardboard with inner plastic liner
1,5-10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated cardboard
Professional, trained professional
Yes

4.1.1 Instrucțiuni specifice de utilizare

The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.1.2 Măsuri specifice de reducere a riscului de utilizare

4.1.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.1.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

--

4.1.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months

4.2 Modul de utilizare

Utilizarea 2 - Rats - professionals - indoor

Tipul produsului	TP 14 - Rodenticide
Descrierea exactă a utilizării autorizate, după caz	Not relevant for rodenticides
Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)	denumirea științifică: Rattus norvegicus denumirea comună: Brown rat etapă de dezvoltare: juvenile to adult denumirea științifică: Rattus rattus denumirea comună: Roof rat etapă de dezvoltare: other: juvenile to adult
Domeniul de utilizare	de interior Indoor
Metoda (metodele) de aplicare	Metoda: cerere momeală Descriere detaliată: Ready-to-use bait to be used in tamper-resistant bait stations
Doza (dozele) și frecvența de aplicare	Frecvența de aplicare: 60-100g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters. Diluare (%): 0 Numărul și calendarul de aplicare: If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
Categoria (categoriile) de utilizatori	industrial profesional
Dimensiunile ambalajului și materialul de ambalare	

Bait product of 10 or 20 gr units
Minimum pack size of 1.5 kg up to 25 kg.
Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.

Type of packaging

Size/volume of the packaging

Material of the packaging

Type and material of closure(s)

Intended user (e.g. professional, non-professional)

Compatibility of the product with the proposed packaging materials (Yes/No)

Labelled or Printed Bottle

from 1.5 Kg- to 5 kg

Plastic composite (HDPE)

Bottle sealed

Professional, trained professional

Yes

Labelled or Printed Bag/Sack

from 1.5 Kg - to 25 kg

Plastic composite

Prefabricated bags or serial production bags , both thermal welded

Professional, trained professional

Yes

Labelled or Printed Bag/Sack - with Inner plastic liner

from 1.5 Kg - to 10 kg - max size inner liner 10 Kg

Plastic composite

Prefabricated bags or serial production bags , both thermal welded

Professional, trained professional

Yes

Labelled Plastic Bucket

from 1.5 Kg -to 25 kg

Labelled plastic (HDPE) bucket

Rectangular or truncated cone bucket sealed

Professional, trained professional

Yes

Labelled Plastic Bucket- with inner plastic liner

from 1.5 - to 10 kg - max size inner liner 10 Kg

Labelled plastic (HDPE) bucket

Rectangular or truncated cone bucket sealed

Professional, trained professional

Yes

Labelled or Printed Carton sachet with plastic inner liner

from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each

Carton + Plastic composite

Prefabricated bags or serial production bags , both thermal welded

Professional, trained professional

Yes

Labelled or Printed Cardboard with inner plastic liner

1,5-10 kg single inner plastic bags of up to 10 Kg each

Carton + Plastic composite

Prefabricated cardboard

Professional, trained professional

Yes

4.2.1 Instrucțiuni specifice de utilizare

The bait stations should be visited at least every 5 to 7 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

4.2.2 Măsuri specifice de reducere a riscului de utilizare

4.2.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait stations close to water drainage systems, ensure that bait contact with water is avoided.

4.2.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.2.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months

4.3 Modul de utilizare

Utilizarea 3 - House mice and rats - professionals -outdoor around buildings

Tipul produsului

TP 14 - Rodenticide

Descrierea exactă a utilizării autorizate, după caz

Not relevant for rodenticides

Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)

denumirea științifică: Rattus norvegicus
denumirea comună: Brown rat
etapă de dezvoltare: juvenile to adult

denumirea științifică: Rattus rattus
denumirea comună: Roof rat
etapă de dezvoltare: other: juvenile to adult

denumirea științifică: Mus musculus
denumirea comună: House mouse
etapă de dezvoltare: other: juvenile to adult

Domeniul de utilizare

în aer liber

outdoor around buildings

Metoda (metodele) de aplicare

Metoda: cerere momeală
 Descriere detaliata:
 Ready-to-use bait to be used in tamper-resistant bait stations

Doza (dozele) și frecvența de aplicare

Frecvența de aplicare: House mouse: 40g of bait per bait point. Rattus: 60-100g of bait per bait point. If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.
 Diluare (%): 0
 Numărul și calendarul de aplicare:
 If more than one bait station is needed, the minimum distance between bait stations should be of 5 meters.

Categoria (categoriile) de utilizatori

industrial
 profesional

Dimensiunile ambalajului și materialul de ambalare

Bait product of 10 or 20 gr units
 Minimum pack size of 1.5 kg up to 25 kg.
 Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.

Type of packaging
Size/volume of the packaging
Material of the packaging
Type and material of closure(s)
Intended user (e.g. professional, non-professional)
Compatibility of the product with the proposed packaging materials (Yes/No)
 Labelled or Printed Bottle
 from 1.5 Kg- to 5 kg
 Plastic composite (HDPE)
 Bottle sealed
 Professional, trained professional
 Yes
 Labelled or Printed Bag/Sack
 from 1.5 Kg - to 25 kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes
 Labelled or Printed Bag/Sack - with Inner plastic liner
 from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes
 Labelled Plastic Bucket
 from 1.5 Kg -to 25 kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes
 Labelled Plastic Bucket- with inner plastic liner
 from 1.5 - to 10 kg - max size inner liner 10 Kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes
 Labelled or Printed Carton sachet with plastic inner liner
 from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each
 Carton + Plastic composite

Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled or Printed Cardboard with inner plastic liner
1,5-10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated cardboard
Professional, trained professional
Yes

4.3.1 Instrucțiuni specifice de utilizare

- Protect bait from the atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.
- The bait stations should be visited [for mice - at least every 2 to 3 days at] [for rats - only 5 to 7 days after] the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
- Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

4.3.2 Măsurile specifice de reducere a riscului de utilizare

Do not apply this product directly in the burrows.

4.3.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait stations close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.3.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.3.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months

4.4 Modul de utilizare

Utilizarea 4 - House mice and rats - trained professionals - indoor

Tipul produsului	TP 14 - Rodenticide
Descrierea exactă a utilizării autorizate, după caz	Not relevant for rodenticides
Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)	<p>denumirea științifică: Rattus norvegicus denumirea comună: Brown rat etapă de dezvoltare: juvenile to adult</p> <p>denumirea științifică: Rattus rattus denumirea comună: Roof rat etapă de dezvoltare: other: juvenile to adult</p> <p>denumirea științifică: Mus musculus denumirea comună: House mouse etapă de dezvoltare: other: juvenile to adult</p>
Domeniul de utilizare	<p>de interior</p> <p>indoor</p>
Metoda (metodele) de aplicare	<p>Metoda: cerere momeală Descriere detaliată: Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations.</p>
Doza (dozele) și frecvența de aplicare	<p>Frecvența de aplicare: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point.. Permanent baiting 60-100 gr of bait per bait point. Diluar (%) : 0 Numărul și calendarul de aplicare: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.</p>
Categoria (categoriile) de utilizatori	instruiți profesional
Dimensiunile ambalajului și materialul de ambalare	<p>Bait product of 10 or 20 gr units Minimum pack size of 1.5 kg up to 25 kg. Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.</p> <p>Type of packaging Size/volume of the packaging Material of the packaging Type and material of closure(s) Intended user (e.g. professional, non-professional)</p>

Compatibility of the product with the proposed packaging materials (Yes/No)

Labelled or Printed Bottle
from 1.5 Kg- to 5 kg
Plastic composite (HDPE)
Bottle sealed
Professional, trained professional
Yes

Labelled or Printed Bag/Sack
from 1.5 Kg - to 25 kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes

Labelled or Printed Bag/Sack - with Inner plastic liner
from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes

Labelled Plastic Bucket
from 1.5 Kg -to 25 kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes

Labelled Plastic Bucket- with inner plastic liner
from 1.5 - to 10 kg - max size inner liner 10 Kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes

Labelled or Printed Carton sachet with plastic inner liner
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes

Labelled or Printed Cardboard with inner plastic liner
1,5-10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated cardboard
Professional, trained professional
Yes

4.4.1 Instrucțiuni specifice de utilizare

Remove the remaining product at the end of treatment period.

For permanent baiting :

- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.

- Follow any additional instructions provided by the relevant code of best practice.

4.4.2 Măsuri specifice de reducere a riscului de utilizare

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product in pulsed baiting treatments.

In case of permanent baiting:

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.4.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait points close to water drainage system, ensure that bait contact with water is avoided.

4.4.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.4.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.5 Modul de utilizare

Utilizarea 5 - Mice and rats - trained professionals - outdoor around buildings

Tipul produsului	TP 14 - Rodenticide
Descrierea exactă a utilizării autorizate, după caz	Not relevant for rodenticides
Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)	<p>denumirea științifică: <i>Rattus norvegicus</i> denumirea comună: Brown rat etapă de dezvoltare: juvenile to adult</p> <p>denumirea științifică: <i>Rattus rattus</i> denumirea comună: Roof rat etapă de dezvoltare: other: juvenile to adult</p> <p>denumirea științifică: <i>Mus musculus</i> denumirea comună: House mouse etapă de dezvoltare: other: juvenile to adult</p>

Domeniul de utilizare

în aer liber
 outdoor around buildings

Metoda (metodele) de aplicare

Metoda: cerere momeală
 Descriere detaliata:
 Bait formulations:
 - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations

Doza (dozele) și frecvența de aplicare

Frecvența de aplicare: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Mice - 40 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.
 Diluare (%): 0
 Numărul și calendarul de aplicare:
 Bait products
 Rats
 - High infestation: 100 g of bait per baiting point.
 - Low infestation: 60 g of bait per baiting point.
 Mice
 - 40 g of bait per baiting point.
 Permanent baiting 60-100 gr of bait per bait point.

Categoria (categoriile) de utilizatori

instruiți profesional

Dimensiunile ambalajului și materialul de ambalare

Bait product of 10 or 20 gr units
 Minimum pack size of 1.5 kg up to 25 kg.
 Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.

Type of packaging
Size/volume of the packaging
Material of the packaging
Type and material of closure(s)
Intended user (e.g. professional, non-professional)
Compatibility of the product with the proposed packaging materials (Yes/No)

Labelled or Printed Bottle
 from 1.5 Kg- to 5 kg
 Plastic composite (HDPE)
 Bottle sealed
 Professional, trained professional
 Yes

Labelled or Printed Bag/Sack
 from 1.5 Kg - to 25 kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes

Labelled or Printed Bag/Sack - with Inner plastic liner
 from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
 Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes

Labelled Plastic Bucket

from 1.5 Kg -to 25 kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes
 Labelled Plastic Bucket- with inner plastic liner
 from 1.5 - to 10 kg - max size inner liner 10 Kg
 Labelled plastic (HDPE) bucket
 Rectangular or truncated cone bucket sealed
 Professional, trained professional
 Yes
 Labelled or Printed Carton sachet with plastic inner liner
 from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each
 Carton + Plastic composite
 Prefabricated bags or serial production bags , both thermal welded
 Professional, trained professional
 Yes
 Labelled or Printed Cardboard with inner plastic liner
 1,5-10 kg single inner plastic bags of up to 10 Kg each
 Carton + Plastic composite
 Prefabricated cardboard
 Professional, trained professional
 Yes

4.5.1 Instrucțiuni specifice de utilizare

Remove the remaining product at the end of treatment period.

For permanent baiting :

- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.
- Follow any additional instructions provided by the relevant code of best practice.

For application in covered and protected bait points:

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Follow any additional instructions provided by the relevant code of best practice.

4.5.2 Măsuri specifice de reducere a riscului de utilizare

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign

- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.

- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.

- Do not use the product in pulsed baiting treatments.

-do not apply this product directly in the burrows

In case of permanent baiting:

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.

- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the

assessment of the risk for re-infestation.

4.5.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.5.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.5.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

4.6 Modul de utilizare

Utilizarea 6 - Rats - trained professionals - Outdoor open areas

Tipul produsului	TP 14 - Rodenticide
Descrierea exactă a utilizării autorizate, după caz	Not relevant for rodenticides
Organism(organisme)-țintă (inclusiv stadiul de dezvoltare)	denumirea științifică: Rattus norvegicus denumirea comună: Brown rat etapă de dezvoltare: juvenile to adult denumirea științifică: Rattus rattus denumirea comună: Roof rat etapă de dezvoltare: other: juvenile to adult
Domeniul de utilizare	în aer liber outdoor open areas
Metoda (metodele) de aplicare	Metoda: cerere momeală Descriere detaliată: Bait formulations: - Ready-to-use bait to be used in tamper-resistant bait stations or covered and protected baiting points as long as they provide the same level of protection for non-target species and humans as tamper-resistant bait stations

Doza (dozele) și frecvența de aplicare

Frecvența de aplicare: Bait products Rats - High infestation: 100 g of bait per baiting point. - Low infestation: 60 g of bait per baiting point. Permanent baiting 60-100 gr of bait per bait point.
Diluare (%): 0
Numărul și calendarul de aplicare:
Bait products
Rats
- High infestation: 100 g of bait per baiting point.
- Low infestation: 60 g of bait per baiting point.

Permanent baiting 60-100 gr of bait per bait point.

Categoria (categoriile) de utilizatori

instruiți profesional

Dimensiunile ambalajului și materialul de ambalare

Bait product of 10 or 20 gr units
Minimum pack size of 1.5 kg up to 25 kg.
Where applicable package is restricted to separately packed bags with a maximum bag size of 10 kg.

Type of packaging
Size/volume of the packaging
Material of the packaging
Type and material of closure(s)
Intended user (e.g. professional, non-professional)
Compatibility of the product with the proposed packaging materials (Yes/No)
Labelled or Printed Bottle
from 1.5 Kg- to 5 kg
Plastic composite (HDPE)
Bottle sealed
Professional, trained professional
Yes
Labelled or Printed Bag/Sack
from 1.5 Kg - to 25 kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled or Printed Bag/Sack - with Inner plastic liner
from 1.5 Kg - to 10 kg - max size inner liner 10 Kg
Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled Plastic Bucket
from 1.5 Kg -to 25 kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes
Labelled Plastic Bucket- with inner plastic liner
from 1.5 - to 10 kg - max size inner liner 10 Kg
Labelled plastic (HDPE) bucket
Rectangular or truncated cone bucket sealed
Professional, trained professional
Yes
Labelled or Printed Carton sachet with plastic inner liner
from 1,5 - to 10 kg single inner plastic bags of up to 10 Kg each

Carton + Plastic composite
Prefabricated bags or serial production bags , both thermal welded
Professional, trained professional
Yes
Labelled or Printed Cardboard with inner plastic liner
1,5-10 kg single inner plastic bags of up to 10 Kg each
Carton + Plastic composite
Prefabricated cardboard
Professional, trained professional
Yes

4.6.1 Instrucțiuni specifice de utilizare

- Protect bait from the atmospheric conditions. Place the bait stations in areas not liable to flooding.
- Replace any bait in baiting points in which bait has been damaged by water or contaminated by dirt.
- Remove the remaining product at the end of treatment period
- Follow any additional instructions provided by the relevant code of best practice.

For permanent baiting :

- Where possible, it is recommended that the treated area is revisited every 4 weeks at the latest in order to avoid any selection of a resistant population.
- Follow any additional instructions provided by the relevant code of best practice.

For application in covered and protected bait points:

- For outdoor use, baiting points must be covered and placed in strategic sites to minimise the exposure to non-target species.
- Follow any additional instructions provided by the relevant code of best practice.

4.6.2 Măsurile specifice de reducere a riscului de utilizare

- Where possible, prior to the treatment inform any possible bystanders (e.g. users of the treated area and their surroundings) about the rodent control campaign
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- To reduce risk of secondary poisoning, search for and remove dead rodents during treatment at frequent intervals, in line with the recommendations provided by the relevant code of best practice.
- Products may only be used in permanent treatments at those sites with a high potential for reinvasion when other methods of control have proven insufficient.
- Do not use the product in pulsed baiting treatments.
- do not apply this product directly in the burrows

In case of permanent baiting:

- Permanent baiting is strictly limited to sites with a high potential for reinvasion when other methods of control have proven insufficient.
- The permanent baiting strategy shall be periodically reviewed in the context of integrated pest management (IPM) and the assessment of the risk for re-infestation.

4.6.3 În funcție de modul de utilizare, particularități ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

When placing bait points close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) or water drainage systems, ensure that bait contact with water is avoided.

4.6.4 În funcție de modul de utilizare instrucțiunile pentru eliminarea în siguranță a produsului și ambalajul acestuia

4.6.5 În funcție de modul de utilizare, condițiile de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

Shelf life: 24 months.

Use specific conditions of storage and shelf-life of the product under normal conditions of storage are not necessary when conditions of storage and shelf-life of the product under normal conditions of storage, as detailed in the section "General directions for use (trained professionals)", are implemented.

5. Direcții generale de utilizare

5.1. Instrucțiuni de utilizare

- Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.
- Carry out a pre-baiting survey of the infested area and an on-site assessment in order to identify the rodent species, their places of activity and determine the likely cause and the extent of the infestation.
- Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.
- The product should only be used as part of an integrated pest management (IPM) system, including, amongst others, hygiene measures and, where possible, physical methods of control.
- Consider preventive control measures (e.g. plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.
- Bait stations should be placed in the immediate vicinity of places where rodent activity has been previously observed (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- The product should be placed in the immediate vicinity of places where rodent activity has been previously explored (e.g. travel paths, nesting sites, feedlots, holes, burrows etc.).
- Where possible, bait stations must be fixed to the ground or other structures.
- Bait stations must be clearly labelled to show they contain rodenticides and that they must not be moved or opened (see section 5.3 for the information to be shown on the label).
- When the product is being used in public areas, the areas treated should be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits.
- Bait should be secured so that it cannot be dragged away from the bait station.
- Place the product out of the reach of children, birds, pets and farm animals and other non-target animals.
- Place the product away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.
- Wear protective chemical resistant gloves (rubber) during product handling phase
- When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.
- The frequency of visits to the treated area should be at the discretion of the operator, in the light of the survey conducted at the outset of the treatment. That frequency should be consistent with the recommendations provided by the relevant code of best practice.
- If bait uptake is low relative to the apparent size of the infestation, consider the replacement of bait points to further places and the possibility to change to another bait formulation.
- If after a treatment period of 35 days baits are continued to be consumed and no decline in rodent activity can be observed, the likely cause has to be determined. Where other elements have been excluded, it is likely that there are resistant rodent so consider the use of a non-anticoagulant rodenticide, where available, or a more potent anticoagulant rodenticide. Also consider the use of traps as an alternative control measure.
- Remove the remaining bait or the bait stations at the end of the treatment period.- Instructions for use that are "bait-specific":- Bait in sachets: Do not open the sachets containing the bait.

5.2. Măsurile de reducere a riscurilor

- Where possible, prior to the treatment inform any possible bystanders about the rodent control campaign
- The product information (i.e. label and/or leaflet) shall clearly show that the product shall only be supplied to trained professional users holding certification demonstrating compliance with the applicable training requirements (e.g. "for trained professionals only").
- Do not use in areas where resistance to the active substance can be suspected.
- Products shall not be used beyond 35 days without an evaluation of the state of the infestation and of the efficacy of the treatment unless authorised for permanent baiting treatments.
- Do not rotate the use of different anticoagulants with comparable or weaker potency for resistance management purposes. For rotational use, consider using a non-anticoagulant rodenticide, if available, or a more potent anticoagulant.
- Do not wash the bait stations or utensils used in covered and protected bait points with water between applications.
- Dispose dead rodents in accordance with local requirements.

5.3. Particularitati ale efectelor posibile directe sau indirecte, instrucțiuni de prim ajutor și măsuri de urgență pentru protecția mediului

- This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.
- Antidote: Vitamin K1 administered by medical/veterinary personnel only.
- In case of:
 - Dermal exposure, wash skin with water and then with water and soap.
 - Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eyes lids open at least 10 minutes.
 - Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet
- Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre
- Hazardous to wildlife.

5.4. Instrucțiuni pentru eliminarea în siguranță a produsului și a ambalajului acestuia

At the end of the treatment, dispose the uneaten bait and the packaging in accordance with local requirements

5.5. Condiții de depozitare și termenul de valabilitate a produsului în condiții normale de depozitare

- Store in a dry, cool and well ventilated place. Keep the container closed and away from direct sunlight.
- Store in places prevented from the access of children, birds, pets and farm animals.
- Shelf life: 24 months

6. Alte informații

- Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after effective consumption of the bait.
- Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.
- This product contains a bittering agent and a dye.