

CLH report

Proposal for Harmonised Classification and Labelling

**Based on Regulation (EC) No 1272/2008 (CLP Regulation),
Annex VI, Part 2**

Substance Name: Chlorophene

EC Number: 204-385-8

CAS Number: 120-32-1

Index Number:

Contact details for dossier submitter:

The Norwegian Environment Agency, Norway

Miljødirektoratet

Postboks 5672 Sluppen,

7485 Trondheim

+47 73 58 05 00

Jorid.frydenlund@miljodir.no

Version number: 4

Date: 26.08.2014

CONTENTS

Part A.

1	PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING	5
1.1	SUBSTANCE.....	5
1.2	HARMONISED CLASSIFICATION AND LABELLING PROPOSAL	5
1.3	PROPOSED HARMONISED CLASSIFICATION AND LABELLING BASED ON CLP REGULATION	7
2	BACKGROUND TO THE CLH PROPOSAL	10
2.1	HISTORY OF THE PREVIOUS CLASSIFICATION AND LABELLING	10
2.2	SHORT SUMMARY OF THE SCIENTIFIC JUSTIFICATION FOR THE CLH PROPOSAL	10
2.3	CURRENT HARMONISED CLASSIFICATION AND LABELLING.....	10
3	JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL.....	10
	SCIENTIFIC EVALUATION OF THE DATA	11
1	IDENTITY OF THE SUBSTANCE	11
1.1	NAME AND OTHER IDENTIFIERS OF THE SUBSTANCE.....	11
1.2	COMPOSITION OF THE SUBSTANCE	12
	PURITY/IMPURITIES, ADDITIVES IN THE ACTIVE SUBSTANCE	12
1.3	PHYSICO-CHEMICAL PROPERTIES	13
2	MANUFACTURE AND USES	14
2.1	MANUFACTURE.....	14
2.2	IDENTIFIED USES	14
3	CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES.....	15
3.1.1	<i>Summary and discussion of physico-chemical properties.....</i>	<i>15</i>
3.1.2	<i>Comparison with criteria.....</i>	<i>15</i>
3.1.3	<i>Conclusions on classification and labelling The substance does not meet the criteria for classification for physico-chemical properties (please also refer to table 3 and 9).....</i>	<i>15</i>
4	HUMAN HEALTH HAZARD ASSESSMENT.....	16
4.1	TOXICOKINETICS (ABSORPTION, METABOLISM, DISTRIBUTION AND ELIMINATION)	16
4.1.1	<i>Summary and discussion of toxicokinetics.....</i>	<i>20</i>
4.2	ACUTE TOXICITY	20
4.2.1	<i>Summary and discussion of acute toxicity</i>	<i>21</i>
4.2.2	<i>Comparison with the CLP criteria.....</i>	<i>21</i>
4.2.3	<i>Conclusions on classification and labelling</i>	<i>21</i>
4.3	SPECIFIC TARGET ORGAN TOXICITY – SINGLE EXPOSURE (STOT SE).....	21
4.3.1	<i>Summary and discussion of specific target organ toxicity – single exposure</i>	<i>21</i>
4.3.2	<i>Comparison with the CLP criteria.....</i>	<i>21</i>
4.3.3	<i>Conclusions on classification and labelling</i>	<i>21</i>
4.4	IRRITATION	22
4.4.1	<i>Skin irritation.....</i>	<i>22</i>
4.4.1.1	<i>Summary and discussion of skin irritation.....</i>	<i>22</i>
4.4.1.2	<i>Comparison with the CLP criteria.....</i>	<i>22</i>
4.4.1.3	<i>Conclusions on classification and labelling</i>	<i>23</i>
4.4.2	<i>Eye irritation.....</i>	<i>23</i>
4.4.2.1	<i>Summary and discussion of eye irritation</i>	<i>24</i>
4.4.2.2	<i>Comparison with the CLP criteria.....</i>	<i>25</i>
4.4.2.3	<i>Conclusions on classification and labelling</i>	<i>25</i>
4.5	SENSITISATION.....	25
4.5.1	<i>Skin sensitisation</i>	<i>25</i>
4.5.1.1	<i>Summary and discussion of skin sensitisation</i>	<i>26</i>
4.5.1.2	<i>Comparison with the CLP criteria.....</i>	<i>26</i>
4.5.1.3	<i>Conclusions on classification and labelling</i>	<i>26</i>

4.5.2	<i>Respiratory tract sensitisation</i>	26
4.5.2.1	Non-human information:.....	26
4.5.2.2	Human information:.....	27
4.5.2.3	Conclusions on classification and labelling	27
4.6	REPEATED DOSE TOXICITY	27
4.6.1	<i>Summary and discussion of repeated dose toxicity</i>	37
4.6.2	<i>Comparison with the CLP criteria</i>	38
4.6.3	<i>Conclusions on classification and labelling:</i>	39
4.7	GERM CELL MUTAGENICITY (MUTAGENICITY).....	39
4.8.3	<i>Summary and discussion of mutagenicity</i>	44
4.8.4	<i>Conclusions on classification and labelling</i>	44
4.8	CARCINOGENICITY	45
4.8.1	<i>Summary and discussions of carcinogenicity</i>	52
4.9.2	<i>Comparison with the CLP criteria</i>	54
4.9.3	<i>Conclusions on classification and labelling</i>	56
4.9	TOXICITY FOR REPRODUCTION	57
4.9.1	<i>Effects on sexual function and fertility</i>	57
4.9.1.1	Summary and discussion of effects on sexual function and fertility	62
4.9.1.2	Comparison with the CLP criteria.....	62
4.9.1.3	Conclusions on classification and labelling	63
4.9.1.4	Summary and discussion of effects on lactation	63
4.9.1.5	Comparison with the CLP criteria and conclusions on classification and labelling	64
4.9.2	<i>Developmental toxicity</i>	65
4.9.2.1	Summary and discussion of developmental toxicity	68
4.9.2.2	Conclusions on classification and labelling	69
4.10	NEUROTOXICITY.....	69
4.11	SPECIFIC INVESTIGATIONS: OTHER STUDIES.....	69
4.12	HUMAN INFORMATION	70
5	ENVIRONMENTAL HAZARD ASSESSMENT	71
5.1	DEGRADATION	71
5.1.1	<i>Stability</i>	72
5.1.1.1	Abiotic degradation.....	72
	Hydrolysis.....	72
	Photolysis in water.....	72
	Photolysis in air and vaporisation behaviour	73
5.1.2	<i>Biodegradation</i>	74
5.1.2.1	Biodegradation estimation	74
5.1.2.2	Screening tests	74
5.1.2.3	Simulation tests.....	76
5.1.3	<i>Summary and discussion of degradation</i>	78
5.1.3.1	Conclusion on abiotic degradation.....	78
5.1.3.2	Conclusion on biodegradation.....	78
5.2	ENVIRONMENTAL DISTRIBUTION.....	79
5.2.1	<i>Adsorption/Desorption</i>	79
5.2.2	<i>Volatilisation</i>	81
	<i>Atmosphere</i>	81
5.2.3	<i>Distribution modelling</i>	81
5.3	AQUATIC BIOACCUMULATION.....	81
5.3.1	<i>Aquatic bioaccumulation</i>	81
5.3.1.1	Bioaccumulation estimation.....	81
5.3.1.2	Measured bioaccumulation data.....	82
5.3.2	<i>Summary and discussion of aquatic bioaccumulation</i>	83
5.4	AQUATIC TOXICITY	83
5.4.1	<i>Fish</i>	84
5.4.1.1	Short-term toxicity to fish	84
5.4.1.2	Long-term toxicity to fish	84
5.4.2	<i>Aquatic invertebrates</i>	85
5.4.2.1	Short-term toxicity to aquatic invertebrates	85
5.4.2.2	Long-term toxicity to aquatic invertebrates	85
5.4.3	<i>Algae and aquatic plants</i>	86
5.4.4	<i>Other aquatic organisms (including sediment)</i>	87
	Inhibition of microbial activity (aquatic)	87

CLH REPORT FOR CHLOROPHENE

	Effects on sediment dwelling organisms.....	88
5.5	COMPARISON WITH CRITERIA FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4).....	88
5.6	CONCLUSIONS ON CLASSIFICATION AND LABELLING FOR ENVIRONMENTAL HAZARDS (SECTIONS 5.1 – 5.4).....	89
6	OTHER INFORMATION.....	90
7	REFERENCES.....	91
8	ANNEXES (EXCERPTS OF DOCUMENT III-A TO THE ACTIVE SUBSTANCE DOSSIER).....	101

Part A.

1 PROPOSAL FOR HARMONISED CLASSIFICATION AND LABELLING

1.1 Substance

Table 1: Substance identity

Substance name:	Chlorophene IUPAC name: 2-Benzyl-4-chlorophenol
EC number:	204-385-8
CAS number:	120-32-1
Annex VI Index number:	Not in Annex VI
Degree of purity:	Minimum degree of purity 96,7% w/w
Impurities:	Impurities are not present at concentrations that affect the Classification and Labelling of this substance. Detailed information about the impurities is presented in the confidential part of IUCLID.

1.2 Harmonised classification and labelling proposal

Table 2: The current Annex VI entry and the proposed harmonised classification

	CLP Regulation
Current entry in Annex VI, CLP Regulation	Not in Annex VI
Current proposal for consideration by RAC	Acute Tox 4 H332: Harmful if inhaled Skin Irrit 2 H315: Causes skin irritation Skin Sens 1A H317: May cause an allergic skin reaction Eye Dam 1 H318: Causes serious eye damage

CLH REPORT FOR CHLOROPHENE

	<p>STOT RE1 H372: Causes damage to kidneys through prolonged or repeated exposure</p> <p>Carc 2 H351: Suspected of causing cancer</p> <p>Repr 2 H361f: Suspected of damaging fertility</p> <p>Aquatic Acute 1 H400: Very toxic to aquatic life M-factor: 1</p> <p>Aquatic Chronic 1 H410: Very toxic to aquatic life with long lasting effects M-factor: 100</p>
<p>Resulting harmonised classification (future entry in Annex VI, CLP Regulation)</p>	

1.3 Proposed harmonised classification and labelling based on CLP Regulation

Table 3: Proposed classification according to the CLP Regulation

CLP Annex I ref	Hazard class	Proposed classification	Proposed SCLs and/or M-factors	Current classification ¹⁾	Reason for no classification ²⁾
2.1.	Explosives	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.2.	Flammable gases	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.3.	Flammable aerosols	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.4.	Oxidising gases	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.5.	Gases under pressure	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.6.	Flammable liquids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.7.	Flammable solids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.8.	Self-reactive substances and mixtures	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.9.	Pyrophoric liquids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.10.	Pyrophoric solids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.11.	Self-heating substances and mixtures	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.12.	Substances and mixtures which in contact with water emit flammable gases	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.13.	Oxidising liquids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.14.	Oxidising solids	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.15.	Organic peroxides	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
2.16.	Substance and mixtures corrosive to metals	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
3.1.	Acute toxicity - oral	Not classified	Not applicable	Not classified	Conclusive but not

CLH REPORT FOR CHLOROPHENE

					sufficient for classification
	Acute toxicity - dermal	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
	Acute toxicity - inhalation	Acute Tox 4 H332 Harmful if inhaled	Not applicable	Not classified	
3.2.	Skin corrosion / irritation	Skin Irrit 2 H315 Causes Skin irritation	Not applicable	Not classified	
3.3.	Serious eye damage / eye irritation	Eye Dam 1 H318 Causes serious eye damage	Not applicable	Not classified	
3.4.	Respiratory sensitisation	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
3.4.	Skin sensitisation	Skin sens 1A H317 May cause an allergic skin reaction	Not applicable	Not classified	
3.5.	Germ cell mutagenicity	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
3.6.	Carcinogenicity	Carc 2 H351 Suspected of causing cancer	Not applicable	Not classified	
3.7.	Reproductive toxicity	Repr 2 H361f Suspected of damaging fertility	Not applicable	Not classified	
3.8.	Specific target organ toxicity –single exposure	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification
3.9.	Specific target organ toxicity – repeated exposure	STOT RE1 H372 Causes damage to kidneys through prolonged or repeated exposure	Not applicable	Not classified	
3.10.	Aspiration hazard	Not classified	Not applicable	Not classified	
4.1.	Hazardous to the aquatic environment	Aquatic Acute 1 H400 Very toxic to aquatic life Aquatic chronic 1 H410 Very toxic to aquatic life with long	M-factor of 1 M-factor of 100	Not classified	

CLH REPORT FOR CHLOROPHENE

		lasting effects			
5.1.	Hazardous to the ozone layer	Not classified	Not applicable	Not classified	Conclusive but not sufficient for classification ³⁾

¹⁾ Including specific concentration limits (SCLs) and M-factors

²⁾ Data lacking, inconclusive, or conclusive but not sufficient for classification

³⁾ Not listed on Annex I of Regulation (EC) No 1005/2009 on substances that deplete the ozone layer

Hazard statements:

H332: Harmful if inhaled

H315: Causes skin irritation

H317: May cause an allergic skin reaction.

H318: Causes serious eye damage

H372: Causes damage to kidneys through prolonged or repeated exposure

H351: Suspected of causing cancer.

H361f: Suspected of damaging fertility

H400: Very toxic to aquatic life

H410: Very toxic to aquatic life with long lasting effects

2 BACKGROUND TO THE CLH PROPOSAL

2.1 History of the previous classification and labelling

Chlorophene is not included in Annex VI of the CLP regulation.

2.2 Short summary of the scientific justification for the CLH proposal

Concerning physico-chemical properties, chlorophene does not fulfil the criteria for a classification according to Regulation (EC) No 1272/2008 (CLP). Therefore no classification is required regarding physico-chemical hazards.

The active substance is harmful by inhalation and irritating to skin. It presents risk of serious damage to eyes and may cause sensitisation by skin contact. It cause damage to kidneys through prolonged or repeated exposure and is suspected of causing cancer. It is suspected of damaging fertility. It is very toxic to aquatic organisms and may cause long-term adverse effects in the aquatic environment.

2.3 Current harmonised classification and labelling

Chlorophene is not included in Annex VI of the CLP regulation.

3 JUSTIFICATION THAT ACTION IS NEEDED AT COMMUNITY LEVEL

No justification is needed

Part B.

SCIENTIFIC EVALUATION OF THE DATA


1 IDENTITY OF THE SUBSTANCE

1.1 Name and other identifiers of the substance

Table 4: Substance identity

EC number:	204-385-8
EC name:	Chlorophene
CAS number (EC inventory):	
CAS number:	120-32-1
CAS name:	
IUPAC name:	2-Benzyl-4-chlorophenol
CLP Annex VI Index number:	Chlorophene is not included in Annex VI of the CLP regulation.
Molecular formula:	$C_{13}H_{11}ClO$
Molecular weight range:	218.7 g/mol

Structural formula:



1.2 Composition of the substance

Purity/impurities, additives in the active substance

Chlorophene is an active substance with a minimal purity of 95% as specified from the commercial producers. However the 5 batch analyses from the reference source gives a purity of minimum 96,7 % (mean conc. - 3*SD) which will be proposed as purity for the approval decision of chlorophene as an active substance according to the Biocide regulation (EU) No 528/2012.

No impurity reported has been found to be of relevance for the CLP proposal.

The identity of impurities and additives in the active substance chlorophene as manufactured is confidential.

Representative production batches of the active substance are analysed for their chlorophene and impurities content.

In all study summaries information on degree of purity and impurities is given when available (doc III documents). Information on the degree of purity of the material used for most of the studies show a higher degree of purity than that proposed as minimum purity.

1.3 Physico-chemical properties

Table 5: Summary of physico - chemical properties

Property	Value	Reference	Comment (e.g. measured or estimated)
State of the substance at 20°C and 101,3 kPa	Solid Purity: 97.9 % White to slightly yellow	Kraus, 2006a	Visual assessment
Melting/freezing point	45.9 °C	Jungheim, 2007a	EC method A.1
Boiling point		Jungheim, 2007a	EC method A.2. Up to the decomposition there is no boiling point of the substance. Exothermal decomposition starts at 110 °C.
Relative density	1.317 at 20 °C	Jungheim, 2007a	EC method A.3
Vapour pressure	< 1.0E-03 Pa at 20 °C < 1.0E-03 Pa at 25 °C 1.66E-02 Pa at 50 °C	Olf, 2006	EC method A.4
Surface tension	57.3 mN/m at 20 °C Chlorophene is surface active.	Jungheim, 2007a	EC method A.5
Water solubility	Results at pH 7: 0.083 g/L at 10°C 0.117 g/L at 20°C 0.199 g/L at 30°C Temperature dependence on water solubility was observed. An effect of pH-value is not expected. Comparison with the tested phenolic substance <i>ortho</i> -phenylphenol does not deliver any indication.	Jungheim, 2006a and Erstling, 2002	EC method A.6
Partition coefficient n-octanol/water	Results at 25 °C: The log Pow is 4.276 for the unionised species. Log Pow = 4.276 at pH 4 Log Pow = 4.275 at pH 7 Log Pow = 4.175 at pH 9 pH dependence on log Pow was observed. An effect of temperature is not expected. Comparison with the phenolic substance <i>ortho</i> -phenylphenol does not deliver any indication.	Greenwood, 2003a; Feldhues, 2006; Erstling, 2002 and Jungheim, 2004	OECD guideline 107
Flash point	-	-	Not applicable as the substance is a solid

CLH REPORT FOR CHLOROPHENE

Property	Value	Reference	Comment (e.g. measured or estimated)
Flammability	The substance is not highly flammable.	Heinz, 2007	EC method A.10
Explosive properties	Not an explosive substance	-	Examination of the chemical structure indicates that the substance does not possess any explosive properties.
Self-ignition temperature	-	Heinz, 2007	EC method A.16. The substance gave no exothermic indication up to its melting point.
Oxidising properties	Not an oxidising substance	-	Examination of the chemical structure indicates that the substance does not possess any oxidising substances
Granulometry	Not conducted	-	-
Stability in organic solvents and identity of relevant degradation products	The solubility of chlorophene in methanol and toluene at 10, 20 and 30 °C is > 250 g/L.	Jungheim, 2007c	CIPAC MT 157 CIPAC MT 181
Dissociation constant	pKa = 9.59	Greenwood, 2003a	OECD guideline 112
Viscosity	-	-	Not applicable as the substance is a solid

2 MANUFACTURE AND USES

2.1 Manufacture

Not necessary for biocides.

2.2 Identified uses

Intended biocide use areas:

Product type 2: Disinfectants and algacides not intended for direct application to humans or animals

The representative product, Remedor with the active substance chlorophene is intended to be used as a heavy duty disinfectant for both professional and private use. Professional use includes heavy duty disinfection of surgery rooms and infectious disease wards as well as small-area use for disinfection of objects as washbasins and toilet facilities in hospitals by professional cleaning personnel. Private use of chlorophene is also restricted to heavy duty disinfection of objects, such as as washbasin and toilet facilities. Professional users may be expected to use chlorophene containing products on a daily basis, while non-professional use occurs more rarely, maybe on a weekly basis.

Product-type 3: Veterinary hygiene

The representative product, Remedor with the active substance chlorophene is intended to be used by professional workers to control pathogenic micro-organisms in industrial poultry barns. Application is performed using powered medium pressure spray equipment which sprays an even

layer across the surfaces to be treated. Disinfection of poultry barns is normally performed once every 6-8 weeks, but the task may be performed by specialised disinfectors who provide cleaning services for animal facilities. These workers may perform this task on a daily basis, and are expected to use personal protective equipment.

3 CLASSIFICATION FOR PHYSICO-CHEMICAL PROPERTIES

Table 6: Summary table for relevant physico-chemical studies

Method	Results	Remarks	Reference
Refer to table 9			

3.1.1 Summary and discussion of physico-chemical properties

The data on physico-chemical properties are conclusive but not sufficient for classification. Refer to table 9.

3.1.2 Comparison with criteria

The substance does not meet the criteria for classification for physico-chemical properties (please also refer to table 3 and 9).

3.1.3 Conclusions on classification and labelling

The substance does not meet the criteria for classification for physico-chemical properties (please also refer to table 3 and 9).

4 HUMAN HEALTH HAZARD ASSESSMENT

4.1 Toxicokinetics (absorption, metabolism, distribution and elimination)

Table 7: Toxicokinetic and metabolism studies with chlorophene

Route	Method Guideline	Species Strain Sex No/group	Label	Dose levels	Reference
Gavage, Dermal, Intravenous injection (i.v).	Absorption, excretion, distribution, metabolism No guideline Non-GLP	Rat, Fischer 344 ♂, ≥ 3 per group	¹⁴ C-o-Benzyl-p-chlorophenol ([¹⁴ C]BCP)	Gavage: 10, 100, 1000 mg/kg bw Dermal: 10 mg/kg bw I.v.: 5, 10, 25 mg/kg bw	Kao & Birnbaum, 1986 A6_2 (1) KEY STUDY
Dermal	US-EPA §85-2 (1982) ≅ OECD 427	Rat, Sprague-Dawley ♂, 16 per group	¹⁴ C-o-Benzyl-p-chlorophenol ([¹⁴ C]BCP) (5% solution)	0.5, 5, 50 mg /mL 5, 50, 500 µg/cm ² ≅ 0.3, 3, 30 mg/kg bw	Confidential, 1994 A6_2(2) KEY STUDY

The **absorption, distribution, metabolism and excretion** of ¹⁴C-labelled chlorophene were studied in male F344 rats (Kao and Brimbaum, 1986). The test substance was administered via gavage, intravenous (i.v.) injection and dermal application. Three animals were used for each time point in each study. The study included examination of excretion of radioactivity into bile in the i.v. experiment. The recovery of labelled compound ranged between 90.0-101.4% (Doc IIIA6_2(1); Table A6_2-2: Tissue distribution of chlorophene in tissues after different routes of exposure). Clinical signs were not reported in any of the experiments.

Following gavage administration of ¹⁴C-o-Benzyl-p-chlorophenol ([¹⁴C]BCP) in corn oil at dosages of 10, 100 and 1000 mg/kg bw (single dose), [¹⁴C]BCP-derived was almost completely excreted (>92%) in urine and faeces after 3 days. The relative amounts excreted in urine and faeces were highly dose-dependent. Three days after gavage administration of 10 mg/kg bw ~ 24% and 76% [¹⁴C]BCP-derived had been excreted in urine and faeces, respectively. The rate of faecal BCP excretion decreased to 41% when the dose was increased to 100 mg/kg bw. However, when the dose given was 1000 mg/kg bw, relative faecal excretion was increased (62%) compared to the mid-dose, but was still lower than in the low dose group (10 mg/kg bw)(Doc IIIA6_2(1); Table A6_2-2: Tissue distribution of chlorophene in tissues after different routes of exposure^a). For the mid-dose group urinary excretion was the major excretion route, while for the lowest and highest dose group faecal excretion was the major excretion route. The lowest measured urine excretion after 72 h was found in the group given 10 mg/kg by gavage and was 24.5%. In the tissues of these animals 0.68% of the total dose was recovered.

Since the levels in bile were not measured after oral administration in the submitted ADME study, an oral absorption could be estimated based on the lowest urine excretion in addition to the chlorophene levels found in the tissues. An estimation of oral absorption for chlorophene based on only urine excretion and tissue levels in animals given 10 mg/kg bw gavage would be 25%. However, in the 417 OECD guideline for the testing of chemicals (Toxicokinetics) adopted 22 July 2010, it is indicated that "oral and IV administration of test substance and measurement of net test

substance present in urine plus expired air plus carcass by each of the two routes" could be used to estimate oral absorption. An estimation of oral absorption for chlorophene based on this assumption would be 70 % $[(0.68 + 24.46) / (1.31 + 34.9)] * 100 \% = 69.4 \%$; figures could be found in the Table 8 below). As the first assumption is assumed to be too conservative and an alternative option is given in the latest version of the OECD 417 guidance, an oral absorption of 70 % was decided for chlorophene.

For the dermal absorption studies, BCP was dissolved in acetone at a concentration of 40 mg/ml, and 10 mg/kg [¹⁴C]BCP (50 µl/animal) was applied to 3.9 cm² intrascapular area (hair was clipped). A perforated tissue capsule was held over the treated area with cyanoacrylated adhesive. Excretion via faeces and urine were 34.4%, 50.5% and 59% of the applied dose after 1, 2 and 3 days, respectively. Approximately 32 % of the total dose was found at the skin site and 3% were in tissues and skin. The recovery of BCP after dermal exposure was approximately 95% (Doc IIIA6_2(1); Table A6_2-2: Tissue distribution of chlorophene in tissues after different routes of exposure). The major excretion route was also for dermal absorption via faeces. No information about stripping of the skin before measuring the BCP level was included. Based on the levels in urine, faeces and tissues, dermal absorption of chlorophene was approximately 62%.

To examine biliary excretion 5, 10, or 25 mg/kg bw BCP was given by injection into the femoral vein. The rats were anesthetized with sodium pentobarbital and the bile duct was cannulated. Bile was collected for 6 h after treatment. After 5, 10 and 25 mg/kg bw, 87%, 72% and 56% of the total radioactivity appeared in the bile, respectively. The excretion was statistical significantly decreased with increasing exposure dose. Also by comparing faecal excretion after 3 d and biliary excretion after 6 h indicated that less BCP-derived radioactivity was excreted in faeces than in bile (54% versus 72%). This is suggested to be caused by a deconjugation of glucuronyl conjugates by beta glucuronidase activity of intestinal microorganisms and the resulting parent BCP and 4 hydroxy-BCP could then be reabsorbed. These findings indicate that BCP is subject to enterohepatic circulation.

To examine distribution, BCP was dissolved in Emulphor:etanol:water (1:1:4) at a concentration of 10 mg/ml and was injected i.v. at 1 ml/kg into a the tail vein. Also for the i.v treatment the major excretion route was via faeces, similar to the excretion after dermal absorption. After 3 days 88% of the dose was excreted. After i.v administration no volatile metabolites or [¹⁴C]CO₂ were detected in exhaled air. Three days after treatment, regardless of dose and exposure route, no tissue contained more than 1% of the total administrated radioactivity. The liver contained the highest percentage of the total administrated BCP-derived radioactivity (0.63%) of all tissues. However, the highest concentration (µg BCP/g tissue) of BCP-derived radioactivity was found in the kidney during the whole measuring period. The BCP concentration in the kidneys was almost 15 times higher than in liver (Doc IIA; Table A6_2-3: Pharmacokinetic parameters for the elimination of chlorophene-derived radioactivity). After 72 h post treatment the highest concentration was found in kidney, liver and spleen 1.7, 1.6, and 0.8 % of the total dose per g tissue, respectively. Muscle, fat and skin contained significant BCP-derived radioactivity 15 min post administration, but little remained after 24 h (< 0.1% total dose/g tissue Doc IIIA6_2(1); Table A6_2-4: Tissue distribution of chlorophene in tissues after i.v. exposure). After giving 10 mg/kg bw the half-life of the total BCP-derived radioactivity in faeces and urine together was found to be 14 h. In blood it was suggested that two BCP-derived components were eliminated with a half-lives of 0.7 and 9.9 h (Doc IIIA6_2(1); Table A6_2-5: Pharmacokinetic parameters for the elimination of chlorophene-derived radioactivity).

The major *in vivo* metabolites detected after BCP exposure were glucuronyl conjugates of chlorophene and hydroxy-chlorophene in faeces and urine. Fifteen minutes after exposure to 10 mg/kg bw i.v. 0.5% and 85% of the total BCP-derived radioactivity were excreted in the bile as UDP-glucuronyl conjugate of OH-BCP and BCP, respectively. After 6 h the proportion of OH-BCP UDP-glucuronyl conjugate were increased to 6%, while the proportion of BCP UDP-glucuronyl conjugate was decreased to 36%. The percent of the conjugates out of the total radioactivity excreted in bile at 15 minutes and 6 hours was 20-25% and 1-2%, respectively. Twenty-four hours after i.v. exposure to 10 mg/kg bw 0.4% and 16% of the total BCP-derived radioactivity were excreted in the urine as UDP-glucuronyl conjugate of OH-BCP and BCP, respectively (Doc IIIA6_2(1) Table A6_2-3 Glucuronide conjugates in bile and urine) . The percent of the conjugates of the total radioactivity excreted in urine was between 25-40% after 24 h. Glutathione conjugates were also found in urine. Based on the study results, a metabolic pathway has been proposed (Figure 1: Proposed metabolic pathway of chlorophene in rats).

Figure 1: Proposed metabolic pathway of chlorophene in rats




Table 8: Distribution of chlorophene in tissues after different routes of exposure^a (figures from Doc IIIA; Table A6_2-2)

Percent of total dose		
Tissue	Intravenous	Oral
	10 mg/kg	10 mg/kg
All tissues	1.31	0.68
Urine	34.9	24.5
Faeces	53.78	76.3
Recovery^b	89.98	101.44
^a at 72 h post administration ^b radioactivity in all tissues +excreta		

Dermal absorption of chlorophene from a commercial 5% disinfectant solution was determined *in vivo* in four male rats (Sprague-Dawley) per group (confidential, 1994 / A6_2(2)). The doses used were 10, 100 and 1000 mg of formulation/ml which corresponds to 0.5, 5 or 50 mg ¹⁴C chlorophene/ml. The solution was diluted in water and applied on a 15 cm² area of shorn and intact dorsal skin. The exposure periods were 1, 4, and 10 h. Before termination the protective cover was removed and the treated skin was wiped with gauze pads. Levels of ¹⁴C chlorophene were determined by liquid scintillation counting. An additional group was treated for 10 h and sacrificed after 168 h to assess the dynamics of excretion and observation of clinical signs. One animal from the mid-dose group showed alopecia on day 7, while an animal from the high-dose group showed increased reactivity on post-treatment days 3 through 5. Since these effects started late during the observation period they were not considered to be treatment related. Absorption of chlorophene was incomplete over the course of a 10 h exposure. The absorption values for all three dose groups sacrificed after 10 h ranged from 41% to 52%, while the values ranged from 42% to 60% for the group sacrificed after 168 h. The dose was calculated based on the levels measured in skin, blood, urine, faeces, remaining carcass and cage wash (Doc IIIA; Table 6 2-2: Distribution of label in the analysed compartments (10-h exposure) [% of applied dose]). Between 3% and 15.5% of the dose remained in the skin after 168 h. About half of the absorbed material was absorbed during the first hour. The lack of strong differences between dose groups indicates that no concentration-dependent limit for the rate of chlorophene absorption exists in the tested dose range. After 10 h the majority of the radiolabelled material was found in the urine, whereas for the animals sacrificed after 168 h a higher portion of labelled material was found in the faeces. Based on the highest measured dermal absorption value in this study the dermal absorption for chlorophene was 60%.

4.1.1 Summary and discussion of toxicokinetics

In an ADME study of chlorophene in rats oral administration of chlorophene resulted in higher relative percentages of chlorophene excreted in the faeces compared compared to i.v. administration. After dermal application, a high percentage of the total dose of chlorophene was present at the application site. These findings indicated that chlorophene was incompletely absorbed through both intestine and skin.

Since the levels in bile were not measured after oral administration in the submitted ADME study, an oral absorption could be estimated based on the lowest urine excretion in addition to the chlorophene levels found in the tissues. As this assumption is assumed to be too conservative; an estimation of oral absorption for chlorophene based on results from an oral and IV administration of test substance (measurement of net test substance present in urine plus expired air plus carcass by each of the two routes) was used. Based on these assumptions an **oral absorption of 70 %** was decided for chlorophene. Based on the levels in urine, faeces and tissues, **dermal absorption of chlorophene was approximately 62%** in the study of Kao & Birnbaum while in the other study (confidential, 1994 / A6_2(2)) the dermal absorption value was approximately **60%**.

Most of the administered chlorophene was excreted and the tissue levels were generally low within 3d post administration (except for the dermal study where 32 % of the total dose was found at the skin site). However, the highest concentration of BCP-derived radioactivity was found in the kidney during the whole measuring period. This affinity of renal tissue for chlorophene is likely to play a role in the suggested nephrotoxicity of this compound. In addition, the studies indicated that enterohepatic circulation was involved in chlorophene disposition.

The major *in vivo* metabolites detected after BCP exposure were glucuronyl conjugates of chlorophene and hydroxy-chlorophene in faeces and urine.

4.2 Acute toxicity

Summary table of relevant acute toxicity studies:

Table 9: Acute toxicity of chlorophene

Route	Method Guideline	Species, Strain, Sex, No/group	Dosage	Value LD ₅₀ /LC ₅₀	Reference
Oral	LD ₅₀ test ≅ OECD 401 Non-GLP	Rat CD, Sprague-Dawley derived ♂+♀, 5/sex/dose	Single dose at 1500, 2500, 3150, 3969, 5000 mg/kg bw. Post exposure period, 14 days	LD ₅₀ = 3852 mg/kg	Confidential,1983a A6_1_1 KEY STUDY
Dermal	Limit test ≅ OECD 402 Non-GLP	Rat CD, Sprague-Dawley derived ♂+♀, 5/sex/dose	Single dose of 2000 mg/kg bw. Exposure duration, 24 hours.	LD ₅₀ > 2000 mg/kg	Confidential,1983b A6_1_2 KEY STUDY
Inhalation	LC ₅₀ test ≅ OECD 403	Rat CD, Sprague-Dawley	Nose-only: 2.07, 2.40, 3.13 mg/L. Duration of exposure 4 h	LC ₅₀ = 2.43 mg/L/4h Acute Tox 4, H332	Confidential, 1983c

	Non-GLP	derived ♂+♀, 5/sex/concentration		Harmful if inhaled	A6_1_3 KEY STUDY
--	---------	--	--	---------------------------	----------------------------

4.2.1 Summary and discussion of acute toxicity

Chlorophene is of low acute toxicity by the oral ($LD_{50} = 3852$ mg/kg bw) and percutaneous route ($LD_{50} > 2000$ mg/kg bw) and of moderate toxicity via inhalation ($LC_{50} = 2.43$ mg/L/4h). Following inhalation increased lung weights were noted in the decedents, indicating pulmonary irritation and respiratory failure caused by oedema. Three cases of hydronephrosis, four cases of enlarged cervical lymph nodes and one case of hepatic nodule were observed in the decedents after exposure via inhalation.

4.2.2 Comparison with the CLP criteria

The criteria for classification with acute toxic by inhalation category 4 is fulfilled. The LC_{50} value of 2,43 mg/L/4h is > 1 but < 5 (dust/mist), and meets the criteria for classification in category 4.

4.2.3 Conclusions on classification and labelling

According to Regulation (EC) No 1272/2008 (CLP) chlorophene should be classified as:

Acute toxic, Category 4 H332: Harmful if inhaled

4.3 Specific target organ toxicity – single exposure (STOT SE)

4.3.1 Summary and discussion of specific target organ toxicity – single exposure

No significant or severe toxicity to a specific organ in the absence of lethality was observed in acute oral, inhalation or dermal toxicity studies in animals. Additionally, no acute organ toxicity was observed in short-term or long-term studies.

4.3.2 Comparison with the CLP criteria

The observed effect in the relevant animal studies of chlorophene does not meet the CLP criteria for classification for specific target organ toxicity after single exposure.

4.3.3 Conclusions on classification and labelling

No classification proposed.

4.4 Irritation

4.4.1 Skin irritation

Summary table of relevant skin irritation studies:

Table 10: Skin irritation by chlorophene

Species	Method	Average score 24, 48, 72 h		Reversibility	Result	Reference
		Erythema	Oedema			
Rabbit	OPPTS 870.2500 \cong OECD 404	2.89	4.00	Yes	Skin Irrit. 2 H315 Causes skin irritation	Confidential, 2000 A6_1_4(1) KEY STUDY
Rabbit	\cong OECD 404 Non-GLP	1.22	0.22	Yes	Moderately irritating, exfoliation, eschar formation	Confidential, 1983d
Rabbit	No Guideline Non-GLP	4.00 (48 h only)	2.25 (48 h only)	Yes	Strongly irritating	Confidential, 1983

Table 11: Skin irritation (individual scores) of chlorophene (confidential, 2000 /A6_1_4(1))

Observation time	Rabbit no.					
	F05790		F05791		F05792	
	Erythema	Oedema	Erythema	Oedema	Erythema	Oedema
4 h	2b	4	2	4	2b	4
24 h	2b	4	2b	4	2b	4
48 h	3b	4	2b	4	2b	4
72 h	4n	4	4n	4	4n	4
96 h	4n	4	4n	4	4n	4
Mean value 24 + 48 + 72 h	3.0	4.0	3.0	4.0	2.7	4.0
Reversibility	Yes	Yes	Yes	Yes	Yes	Yes
Avg. time for reversion	21 d	14 d	14 d	14 d	14 d	14 d

b blanching

n necrotic appearance

4.4.1.1 Summary and discussion of skin irritation

Chlorophene caused strong irritation on the skin of rabbits with strong erythema and oedema. The key study was performed according to OECD guideline 404.

4.4.1.2 Comparison with the CLP criteria

Classification proposal according to Regulation EC 1272/2008: Skin irritant category 2

The criteria for classification with skin irritation category 2 is fulfilled;

Mean value of $\geq 2,3$ - $\leq 4,0$ for erythema/ eschar or for oedema in at least 2 of 3 tested animals from gradings at 24, 48 and 72 hours after patch removal or, if reactions are delayed, from grades on 3 consecutive days after the onset of skin reactions;

4.4.1.3 Conclusions on classification and labelling

Chlorophene was tested on the skin of rabbits. It caused strong irritation on the skin with strong erythema and oedema. The overall results fulfil the criteria of Directive 2001/59/EC and Regulation (EC) No 1272/2008 (CLP) for classification as a skin irritant:

Skin Irrit Cat 2. H315: Causes skin irritation

4.4.2 Eye irritation

Summary table of relevant eye irritation studies:

Table 12: Eye irritation of chlorophene

Species	Method	Average score 24, 48, 72 h				Reversibility	Result	Reference
		Cornea	Iris	Conjunctiva				
				Redness	Chemosis			
Rabbit	≅ OECD 405 Non-GLP	2.78	0.89	2.67	1.78	No	Eye Dam 1 H318 Causes serious eye damage	Confidential, 1983e A6_1_4(2) KEY STUDY
Rabbit	No Guideline Non-GLP	1.50	1.33	2.17	2.00	Yes	Irritating	Confidential, 1983

Table 13: Eye irritation of chlorophene (confidential, 1983e / A6_1_4(2))

	Cornea	Iris	Conjunctiva	
			Redness	Chemosis
Score (average of animals investigated)	0 to 4	0 to 2	0 to 3	0 to 4
1 h	1.67	0.00	1.00	2.00
24 h	2.33	0.67	2.00	1.67
48 h	3.00	1.00	3.00	1.67
72 h	3.00	1.00	3.00	2.00
Average 24 h, 48 h, 72 h	2.78	0.89	2.67	1.78

Table 14: Eye irritation, individual data of chlorophene (confidential, 1983e / A6_1_4(2))

Observation time	Rabbit nr SP 353 M			
			Conjunctiva	
	Cornea	Iris	Redness	Chemosis
1 h	1	0	1	2
24 h	2	1	2	2
48 h	3	1	3	2
72	3	1	3	2

Observation time	Rabbit nr SP 354 M			
			Conjunctiva	
	Cornea	Iris	Redness	Chemosis
1 h	2	0	1	2
24 h	3	0	2	2
48 h	3	1	3	2
72	3	1	3	2

Observation time	Rabbit nr SP 356 M			
			Conjunctiva	
	Cornea	Iris	Redness	Chemosis
1 h	2	0	1	2
24 h	2	1	2	1
48 h	3	1	3	1
72	3	1	3	2

4.4.2.1 Summary and discussion of eye irritation

Chlorophene caused significant irritation of the eye in tests on albino rabbits (confidential, 1983e / A6_1_4(2)). Lesions of cornea and iris as well as conjunctival redness and chemosis, all of which persisted until the end of the observation period, were noted.

The study was terminated 72 h after treatment in light of the deteriorating condition of the treated eyes, especially the corneae. It was considered that significant resolution of the treatment effects was most improbable within the period of extended observation allowed by the OECD test method.

The study was performed according to OECD 405. No formal GLP compliance statement (GLP was not compulsory during the conduct of study) was applied, but there was a signed QAU statement.

4.4.2.2 Comparison with the CLP criteria

Classification proposal according to Regulation EC 1272/2008: Eye dam. Cat 1,

The criteria for classification with eye damage is fulfilled;

If, when applied to the eye of an animal, a substance produces: — at least in one animal effects on the cornea, iris or conjunctiva that are not expected to reverse or have not fully reversed within an observation period of normally 21 days; and/ or — at least in 2 of 3 tested animals, a positive response of: — corneal opacity ≥ 3 and/or — iritis $> 1,5$ calculated as the mean scores following grading at 24, 48 and 72 hours after installation of the test material.

4.4.2.3 Conclusions on classification and labelling

Chlorophene produced ocular irritation characterised by diffuse opacity or translucency of the whole visible corneal surface, injection of the conjunctival blood vessels and eversion of the eyelids due to moderate chemosis. Positive irritation reactions were observed in all animals.

Irritation responses of the treated eyes and other effects of treatment became more marked at each subsequent examination and culminated in a state of severe ocular irritation four days after treatment. The study was terminated 72 h after treatment due to animal welfare and reversibility was not expected.

Chlorophene should be classified as:

Eye dam. Cat1, H318: Causes serious eye damage according to Regulation (EC) No 1272/2008 (CLP).

4.5 Sensitisation

4.5.1 Skin sensitisation

Summary table of relevant skin sensitisation studies:

Table 15: Skin sensitisation data for chlorophene

Species	Method	Number of animals sensitised/ total number of animals	Result	Reference
Guinea pig	Buehler Test OPPTS 870.2600 \cong OECD 406	24 h: 15/20 48 h: 18/20 72 h: 19/20	Skin sensitising Cat 1A H317 May cause an allergic	Confidential, 2001 A6_1_5 KEY STUDY

			reaction	
Guinea pig	Buehler Test OECD 406	9/20	Sensitising	Confidential, 2002
Guinea pig	Klecak Test No guideline	0/24	Not sensitising	Confidential, 1986

4.5.1.1 Summary and discussion of skin sensitisation

Chlorophene was tested for its skin sensitisation potential in Buehler and Klecak tests on Guinea pigs. The first Buehler test (confidential, 2001 / A6_1_5) used 10% and 5% solutions (10 % was the induction dose) of chlorophene in propylene glycol whereas the second Buehler study (confidential, 2002) featured induction and challenge concentrations of both 50% in polyethylene glycol 400.

In the Buehler test by Glaza moderate to strong erythema reactions were observed at the sites treated with the 10% w/v mixture during the induction phase. Subcutaneous haemorrhaging, blanching, and/or necrotic appearing areas were also observed within the induction sites of all test animals.

After the challenge with the 5% test substance mixture, 19 out of 20 of the test compound group showed very faint to moderate redness. Subcutaneous haemorrhaging, desquamation or fissuring was also observed within the challenge sites of 3 test animals. Very faint to faint erythema was observed in 2 of the 10 naïve control animals after challenge.

4.5.1.2 Comparison with the CLP criteria

According to the criteria mentioned in the Regulation No 1272/2008 on CLP' (> 60 % responding at > 0,2 % to 20 % induction dose); the substance should be classified for skin sensitisation, category 1A: H317 May cause an allergic skin reaction (strong potency on basis of the Buehler Occluded Patch Test).

4.5.1.3 Conclusions on classification and labelling

Based on the positive outcome of two Buehler tests, chlorophene should be classified as:

Skin sensitisation, category 1A with H317: May cause an allergic skin reaction according to Regulation (EC) No 1272/2008 (CLP).

4.5.2 Respiratory tract sensitisation

4.5.2.1 Non-human information:

No information available.

4.5.2.2 Human information:

No information available.

4.5.2.3 Conclusions on classification and labelling

There are no human or animal data indicating sensitization following inhalation, thus classification is not warranted.

4.6 Repeated dose toxicity

Summary table of relevant repeated dose toxicity studies:

Table 16: Subacute, subchronic, and chronic toxicity of chlorophene

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
Oral, gavage	16 days ≡ OECD 407 Non-GLP	Rat F344 ♂ + ♀ 5/sex/group	0, 62.5, 125, 250, 500, 1000 mg/kg bw/day Once daily, 5 days/week	Absolute and relative kidney weights were significantly increased in males at ≥ 125 mg/kg bw/day and in females at 500 and 1000 mg/kg bw/day. Absolute and relative liver weights were significantly increased in males at ≥ 250 mg/kg bw/day and in females at 1000 mg/kg bw/day. At ≥ 250 mg/kg bw/day: absolute and relative thymus weight was decreased in females. At ≥ 500 mg/kg bw/day: absolute and relative thymus weight was decreased in males. Mild to moderate nephropathy was observed in all animals of the 1000 mg/kg bw/day group. The incidence and severity increased with dose (incidence: 62.5 mg/kg 1/10; 125 mg/kg 2/10; 250 mg/kg 2/10; 500 mg/kg 7/10). Myocardial degradation was observed in 8/10 rats of the 1000 mg/kg bw/day group. Two females in the 1000 mg/kg bw/day-group died during the study.	LO(A)EL = 125 mg/kg bw/ day NO(A)EL = 62.5 mg/kg bw/ day	Sendelbach, 1982a A6_3_1 KEY STUDY
Oral,	16 days	Mouse	0, 62.5,	Absolute and relative liver	LO(A)EL = 250	Sendelbach,

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
gavage	≅ OECD 407 Non-GLP	B6C3F ₁ ♂ + ♀ 5/sex/group	125, 250, 500, 1000 mg/kg bw/day Once daily, 5 days/week	weights were increased at ≥ 250 mg/kg bw/day in females and at 500 mg/kg bw/day in male mice. At 500 and 1000 mg/kg bw/day nephropathy was observed in some mice. No changes in relative kidney weight were reported. At 1000 mg/kg bw/day: 3 males and 5 females died.	mg/kg bw/ day NO(A)EL = 125 mg/kg bw/ day	1982b
Oral, capsule	21 days No guideline Non-GLP	Dog Beagle ♂+♀ 3/sex/group. In 100 mg/kg bw/day group: 5 males, 1 female	0, 3, 10, 30, 100 mg/kg bw/day Once daily, 7 days/week	At 100 mg/kg bw/day; lower overall body weight gain. Two of three male dogs at 30 mg/kg bw/day and one of five at 100 mg/kg bw/ day died during the study	LO(A)EL = 100 mg/kg bw/ day NO(A)EL = 30 mg/kg bw/ day	Confidential, 1973a
Dermal	5 day dose-finding study No guideline	Rabbit NZW ♂+♀ 1/sex/group	2.4, 60, 125, 500 mg/kg bw/day. Once daily, 6 h/day, 5 days. Group 2.4 mg/kg bw/day and 60 mg/kg bw/day animals sacrificed at day 6 and the remaining animals sacrificed at day 8	No systemic effects. Local reactions comprised erythema, oedema, atonia and discolouration of the skin. Slight to moderate skin effects (erythema) at 1% and 2,5% Chlorophene in 50% v/v ethanol in distilled water. Moderate to severe skin effects (erythema, atonia, oedema) was observed at ≥ 5%.	Systemic effects: NO(A)EL = 500 mg/kg bw/ day (highest dose) Local effects: LO(A)EL = 1 % NO(A)EL = 2,5% Chlorophene in 50% v/v ethanol in distilled water	Confidential, 1984
Dermal	4 weeks US-EPA 82-2 ≅ OECD 410	Rabbit NZW ♂+♀ 7/sex/group	1, 5, 25 mg/ kg bw/day Once daily, 6 h/day 7 days/week	No systemic effects. At 25 mg/kg bw/day local skin effects were observed (erythema, atonia, discoloration). Occasional animals treated with 5 mg/kg bw/day displayed slight to moderate skin effects	Systemic effects: NO(A)EL = 25 mg/kg bw/ day (highest dose) Local effects: LO(A)EL = 5 mg/kg bw/day NO(A)EL = 1	Confidential, 1989

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
					mg/kg bw/day	
Dermal	3 weeks US-EPA 8.22 ≅ OECD 410	Rabbit NZW ♂+♀ 7/sex/group	0, 4, 20, 100 mg/kg bw/day Once daily, 6 h/day 7 days/week	A total of 14 animals died during the treatment period (0 mg/kg 2 (♂)/14; 4 mg/kg 2 (♂)/14; 20 mg/kg 5(3♂+2♀)/14; 100 mg/kg 5(3♂+2♀)/14). Kidney and liver lesions were rather more common among decedents which had been treated with chlorophene. A higher number of female rabbits had kidney lesions at the time of death (dead/killed during treatment period + sacrificed at the end of the study) in the highest dose group, and there were a tendency towards more severe kidney lesions in the 100 mg/kg bw/day group compared to the control group. At 100 mg/kg bw/day Alkaline phosphatase (ALP) was significantly decreased in females. Local reactions to treatment with chlorophene were observed at ≥ 20 mg/kg bw/day.	Systemic effects based on kidney lesions: LO(A)EL= 100 mg/kg bw/day NO(A)EL = 20 mg/kg bw/ day Local effects: LO(A)EL = 20 mg/kg bw/day NO(A)EL = 4 mg/kg bw/day	Confidential, 1985 A6_3_2(1) KEY STUDY
Dermal	3 weeks ≅ OECD 410	Rabbit HC:NZW ♂+♀ 5/sex/group	0, 10, 40, 160 mg/kg bw/day Once daily, 6 h/day 5 days/week	A dose-dependent increase in histopathological changes in the kidneys was observed in females treated with ≥ 40 mg/kg bw/day. The changes included tubular calcinosis, tubular proliferation and cellular infiltration. Absolute and relative liver weights were significantly decreased in females at 160 mg/kg bw/day. At 160 mg/kg ALP was reduced in males and females. Local reactions to treatment were observed in all animals treated with ≥ 40 mg/kg bw/day. One male control + two females (one control and one low-dose) were killed in extremis.	Systemic effects: LO(A)EL = 40 mg/ kg bw/day NO(A)EL = 10 mg/kg/ day STOT RE1 H372: Causes damage to kidneys through prolonged or repeated exposure Local effects: LO(A)EL = 40 mg/kg bw/day NO(A)EL = 10	Confidential, 1985 A6_3_2(2) KEY STUDY

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
					mg/kg bw/day	
Oral, gavage	95 days ≅ OECD 408	Rat F344/N 10/sex/group	0, 30, 60, 120, 240, 480 mg/kg bw/ day Once daily, 5 days/week	<p>Increased absolute and relative kidney weights and reduced absolute and relative thymus weights at ≥ 240 mg/kg bw/day in females and at 480 mg/kg bw/day in males and females.</p> <p>There seems to be a dose-related increase in incidence and severity of nephropathy in male rats at ≥ 30 mg/kg bw/day, although the severity of the nephropathy was minimal to mild at these doses. At higher doses the severity of nephropathy increased (mild to moderate grade) in the 240 mg/kg bw males and in the 480 mg/kg bw males and females. The incidence of nephropathy was significantly increased in male rats at ≥ 120 mg/kg bw/day as stated in Birnbaum et al., 1986 (in the Discussion). These findings could be interpreted to be of limited significance since Fischer 344 male rats spontaneously develop nephropathy (Dixon et al., 1995). However, very few of the concurrent vehicle controls animals in the current study did develop nephropathy (1/10 males) during the study. Historical control data supporting an assumption for non-significance are not provided.</p>	<p>LO(A)EL = 120 mg/kg bw/day based on dose-related significantly increased incidence of nephropathy and dose-related increased severity of nephropathy</p> <p>NO(A)EL = 60</p>	<p>National Toxicology Program Technical Report Series No. 424 (1994) 6_4_1(1) and</p> <p>Birnbaum et al., 1986</p> <p>KEY STUDY</p>
Oral, gavage	95 days ≅ OECD 408	Mouse B6C3F ₁ 10/sex/group	0, 30, 60, 120, 240, 480 mg/kg bw/ day Once daily, 5 days/week	No effects on final body weight or body weight gain were observed. No treatment-related clinical findings or gross lesions.	NO(A)EL = 480 mg/kg bw/day (highest dose)	National Toxicology Program Technical Report Series No. 424 (1994) A6_4_1(1)
Oral, gavage	95 days ≅ OECD 408	Mouse B6C3F ₁ 10/sex/group	0, 500, 650, 800, 1000 mg/kg bw/ day Once daily,	Survival decreased with increasing doses from 500 mg/kg bw/day in females and from 650 mg/kg bw/day in males. Decreased body weight	LO(A)EL = 500 mg/kg bw/day (lowest dose)	National Toxicology Program Technical Report Series

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
			5 days/week	gain at ≥ 500 mg/kg bw/day in males. Absolute and relative kidney weights were reduced at ≥ 500 mg/kg bw/day in males. Absolute and relative liver weights were significantly increased in females at ≥ 500 mg/kg bw/day and in males at 800 mg/kg bw/day.	NO(A)EL = NA	No. 424 (1994) A6_4_1(1)
Oral, capsule	90 days \cong OECD 409 Non-GLP	Dog Beagle $\text{♂}+\text{♀}$ 4/sex	0, 10, 30, 100, (200) mg/kg bw/day Once daily, 7 days/week	At 200 mg/kg bw/day: body weight was decreased (group was discontinued after 14 days). Reduced body weight gain was observed at 100 mg/kg/day. Absolute kidney weights were significantly increased in males at 100 mg/kg bw/day. Relative kidney weights were significantly increased in males at ≥ 30 mg/kg bw/day and in females at 100 mg/kg bw/day. Relative liver weights were significantly increased in males at ≥ 10 mg/kg bw/day and in females at ≥ 30 mg/kg bw/day. Urinalysis showed that specific gravity was significantly reduced at 100 mg/kg bw/ day in both sexes after 90 days.	LO(A)EL = 30 mg/kg bw/ day NO(A)EL = 10 mg/kg bw/ day	Confidential, 1973b A6_4_1(2) KEY STUDY
Oral, diet	16-week dose-finding study No guideline Non-GLP	Rat ♂ 5/group	0, 150, 500, 1500, 3000 ppm <i>ad libitum</i> \cong 0, 10.6, 39.4, 115.4, 254.4 mg/kg bw/day	At 3000 ppm, i.e. 254.4 mg/kg bw/day: body weight \downarrow	LO(A)EL = 254.4 mg/kg bw/ day NO(A)EL = 115 mg/kg bw/ day	Confidential, 2005
Oral, gavage	13 weeks 65 weeks 2 years US-EPA 83-2 \cong OECD 453	Rat F344/N $\text{♂}+\text{♀}$ 50/sex 3- and 15-month interim sacrifice: 10/sex each	0, 30, 60, 120 mg/kg bw/day (♂) 0, 60, 120, 240 mg/kg bw/day (♀) Once daily, 5 days/week	Severe, time- and dose-related nephropathy was observed, occurring as early as after 3 months (females, not possible to evaluate the males due to high spontaneous incidence of nephropathy). The severity of the nephropathy was significantly increased in a time- and dose-dependent manner both in males and females, with males as the most sensitive sex. The severity of	LO(A)EL= 30 / 120 mg/kg bw/day ($\text{♂}/\text{♀}$) based on severity of nephropathy and increased kidney weight NO(A)EL= NA* / 60 mg/kg bw/day ($\text{♂}/\text{♀}$) based on severity	National Toxicology Program Technical Report Series No.424 (1994) A6_5+6_7 KEY STUDY

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
				nephropathy was significantly increased at 30 mg/kg (males) and 120 mg/kg (females) at 65 and 104 weeks. In male rats dosed for 104 weeks secondary hyperparathyroidism developed. Both males and females showed time- and dose-dependent increases in their relative kidney weights, with males as the most sensitive sex (♂ 30 mg/kg bw/day, ♀ 120 mg/kg bw/day, at 2 years). Similarly, time- and dose-dependent increases in heart weight were also observed (♂ 30 mg/kg bw/day, ♀ 120 mg/kg bw/day, at 2 years).	of nephropathy and increased kidney weight *Not applicable due to effect on lowest tested dose	
Oral, gavage	Two-generation study OECD 416	Rat Wistar (HsdCpb: WU conventionally bred) 30/sex/group	0, 60, 180, 540 mg/kg bw/day ♂: 10 weeks pre-mating, 2 wks mating period ♀: 10 weeks pre-mating, through-out gestation and lactation	A significant and dose-related decrease in terminal body weight was observed in P males at ≥ 180 mg/kg bw/day and in F1 males at ≥ 60 mg/kg bw/day. Reduction in body weight gain during gestation was observed in dams at 540 mg/kg bw/day in the P generation and at 180 and 540 mg/kg bw/day in the F1 generation. Treatment-related kidney effects (nephropathy, dilated tubules, basophilic tubules and lymphocytic infiltration) were observed in P and F1 males at ≥ 60 mg/kg bw/day and P and F1 females at 540 mg/kg bw/day.	LO(A)EL: 60 mg/kg bw in males (kidney effects) NO(A)EL: NA due to effect at lowest dose	Confidential, 2008 A6_8_2(3) (KEY STUDY for 3.8 Reproductive toxicity*)
Oral gavage	13 weeks 65 weeks 2 years. US-EPA 83-2 ≅ OECD 451	Mouse B6C3F ₁ ♂ + ♀ 50/sex 3-month and 15-month interim sacrifice: 10/sex	0, 120, 240, 480 mg/kg bw/day Once daily, 5 days/week	At 2 years the absolute kidney weights of high-dose males were lower than those of the controls at the 3-month interim evaluation. Absolute and relative kidney weights of dosed male mice were lower than those of the controls, and absolute and relative kidney weights of female mice were lower than those of the controls at the mid- and high-dose levels, Kidney weights of all dosed males and females of the mid- and high dose group remained significantly reduced	LO(A)EL : 120 mg/kg bw/day based on increased kidney weights, and significantly time- and dose-dependent increased incidence and severity of nephropathy in both sexes. NO(A)EL: NA	National Toxicology Program Technical Report Series No.424 (1994) A6_7 (1) (KEY STUDY for 3.7 Carcinogenicity*)

CLH REPORT FOR CHLOROPHENE

Route	Duration of study, Method, Guideline	Species, Strain, Sex, no/group	Dose levels, Frequency of application	Results	LO(A)EL NO(A)EL	Reference
				<p>at the 2-year sacrifice.</p> <p>The severity of nephropathy was significantly increased in a time- and dose-dependent manner both in males and females, starting at the lowest dose (120 mg/kg bw), with males as the most sensitive sex.</p> <p>The final mean body weights of all dosed males and mid- and high dose females were lower than those of the controls.</p>	due to effects at lowest dose	

*The studies are key studies for other chapters (3.8 Reproductive toxicity and 3.7 Carcinogenicity) and are included in the Tables in the corresponding chapters.

In rats increased kidney weights were observed in a 16-day gavage study (Sendelbach, 1982a, key study, non-GLP, doses: 0; 62.5; 125; 250; 500; 1000 mg/kg bw/day) at ≥ 125 mg/kg bw/day in males and in females at 500 and 1000 mg/kg bw/day. Mild to moderate nephropathy was observed in all animals of the 1000 mg/kg bw/day group. The incidence and severity of the nephropathy increased with dose (incidence: 62.5 mg/kg bw/day 1/10; 125 mg/kg bw/day 2/10; 250 mg/kg bw/day 2/10; 500 mg/kg bw/day 7/10). Information on the ratios between male and female rats with nephropathy in the different dose-groups is not presented in the report from the US National Toxicology Program.

In mice, liver weights were significantly increased at ≥ 250 mg/kg bw/day in females and at 500 mg/kg bw/day in male mice in an analogous subacute 16-day oral gavage study (Sendelbach, 1982b, non-GLP, doses: 0; 62.5; 125; 250; 500; 1000 mg/kg bw/day). Nephropathy characterized by multifocal tubule dilatation and flattening of the proximal convoluted tubule epithelium, tubule regeneration, and minimal focal epithelial cell necrosis occurred in two 500 mg/kg bw/day and six 1000 mg/kg bw/day male and female mice.

In Beagle pups, no organ-specific effects were observed in a 21-day oral dose-range-finding study (confidential, 1973a / A6_4_1(2), doses: 0; 3; 10; 30; 100 mg/kg bw/day). Weight gain was depressed in the highest dose group (100 mg/kg bw/day). This study was performed prior to the enactment of the Good Laboratory Practice (GLP) regulations and no information on the production, purity or stability of the test material Santophen I (chlorophene) was provided in the study report.

In rabbits, three **subacute dermal studies** (all GLP) were conducted according to OECD guideline 410, with only minor deviations. In a **4-week dermal study in rabbit** (confidential, 1989, doses: 1, 5 and 25 mg/kg bw/day) no systemic effects were observed. Moderate to severe skin effects were observed at the highest dose. In addition, some animals treated with 5 mg/kg bw/day showed slight to moderate skin effects.

In a **3-week dermal study in rabbit** (confidential, 1985 / A6_3_2(1), key study; doses: 0, 4, 20, 100 mg/kg bw/day) a higher number of female rabbits of the high dose group had kidney lesions at the time of death (dead/killed during treatment period + sacrificed at the end of the study), and there was a tendency towards an increased severity of the kidney lesions in the high dose group compared to the control group (Table 17). Alkaline phosphatase (ALP) values were reduced in high-dose females. Local skin effects were observed in animals treated with 20 and 100 mg/kg bw/day of chlorophene. The skin lesions in the high dosage group were more frequent and more severe than the ones observed in the 20 mg/kg bw/day group.

Table 17: Results from repeated dermal dose study in rabbits (confidential, 1985 / A6_3_2(1))

Parameter	Control		4 mg/kg		20 mg/kg		100 mg/kg		Dose-response +/-	
	♂	♀	♂	♀	♂	♀	♂	♀	♂	♀
Number of animals examined	7	7	7	7	7	7	7	7		
Mortality	2	0	2	0	3	2	3	2	-	-
Clinical chemistry										
alkaline phosphatase (ALP)	-	-	-	-	-	-	-	↓*	-	+
At application site										
Encrustation	0	0	0	0	4	4	7	7	+	+
Exfoliation	0	0	0	0	1	0	4	2	+	+
Firm	0	0	0	0	1	1	2	1	+	+
Dark	0	0	0	0	1	0	0	1	-	-
Thickened	0	0	0	0	1	1	4	2	+	+
Ulceration	0	0	0	0	0	0	2	1	+	-

In a **3-week dermal study in rabbit** (confidential, 1985 / A6_3_2(2), key study, doses: 0, 10, 40, 160 mg/kg bw/day) a dose-dependent increase in histopathological changes in the kidneys of female rabbits treated with 40 and 160 mg/kg bw/day of chlorophene was observed (Table 18). The changes included tubular calcinosis, tubular proliferation and cellular infiltration. There were five rabbits in each group with one showing histopathological changes in the unexposed group whereas four and five showed histopathological changes in the groups given 40 and 160 mg/kg bw/day, respectively. Moreover, the severity of the histopathological findings increased with dose. Measurements of kidney-associated clinical chemistry and urinalysis parameters were not statistically different between control and treated animals although histopathological changes were observed. It is reported that serum creatinine and urea levels were not changed until there is a 50% reduction in renal function (Price, 2002). Liver weights were reduced in females at the highest dose (160 mg/kg bw/day) and reduced levels of plasma ALP were found in both sexes. Local reactions to treatment were observed in all animals treated with ≥ 40 mg/kg bw/day.

Table 18: Nephropathy in rabbits (confidential, 1985 / A6_3_2(2),)

Animal	0	10	40	160
M=male F=Female	mg/kg bw/day	mg/kg bw/day	mg/kg bw/day	mg/kg bw/day
M1	Iz1 Tp2	NA	0	Tc3
M2	Iz3 Tp2	NA	Tc2	0
M3	Iz1	NA	Iz1	0
M4	0	NA	Iz1	Iz3
M5	0	NA	0	Tu:Di2
F1	0	NA	Tc2	Tc3
F2	0	NA	0	Tp3 Ix1 P11 Va1
F3	0	NA	Iz3 Tp2	Tu:P11
F4	0	NA	Tc2	Tp1 Iz3
F5	Iz1 Tp2	NA	Tc2 Tp2	Tc1

Iz=Infiltration zellulär (cellular infiltration); Tp=Tubuläre proliferation (Tubular proliferation); Tc= Tubuläre calcinose (Tubular calcinosis); Pl=protein in lumen; Va=Vacuolen im zytoplasma (cytoplasmic vacuolisation); Tu:Pl=Tubuli:Preotein in lumen; The numbers indicate grade of severity; NA=not assessed; 0=None detected

The results from these subacute studies indicate that **dermal exposure to chlorophene induces kidney lesions involve histopathological changes in the kidneys of female rabbits leading to a suggested LO(A)EL at 40 mg/kg bw/day** (3-week study, confidential, 1985 / A6_3_2(2)). **The subacute dermal NO(A)EL for systemic effects in rabbits is 25 mg/kg bw/day based on no systemic effects observed** in the 4-week dermal study (confidential , 1989). The skin lesions at the application site observed in all the subacute dermal studies are readily explained by the irritant properties of the test material. The NO(A)EL for local skin effects in rabbits is 1 mg/kg bw/day based on the occasional skin effects observed at 5 mg/kg bw/day in the 4-week study (confidential, 1989).

In Beagle dogs, a subchronic oral study was conducted (administration in gelatine capsules; confidential 1973b / A6_4_1(2), key study, doses: 0, 10, 30, 100, (200) mg/kg bw/day). This study followed a study design that is similar to OECD 409, but with several deviations. No GLP statement was presented as the study was performed prior to the enactment of the GLP regulations. No specification of the test material Santophen I (chlorophene) was provided and information about the purity and stability of the test material was not presented. In this study weight loss was seen in the highest dose group of 200 mg/kg bw/day. This dose group was therefore discontinued after two weeks. Relative liver weights were significantly increased in males at ≥ 10 mg/kg bw/day and in females at ≥ 30 mg/kg bw/day. This increase in liver weight was considered most likely of low toxicological significance. **Relative weights of kidneys were significantly increased in a dose-dependent manner in male dogs at ≥ 30 mg/kg bw/day.** In female dogs relative kidney weights were significantly increased at 100 mg/kg bw/day. Increased kidney-to-brain ratios and

hypostenuria (i.e. inability to concentrate urine; after 90 days) were seen at 100 mg/kg bw/day in both sexes. The observed kidney effects are likely indicators of early nephropathy and **the subchronic oral NO(A)EL in dogs is therefore 10 mg/kg bw/day.**

In rodents, subchronic and chronic gavage studies were performed under the auspices of the US National Toxicology Program (NTP Technical Report 424, 1994; Birnbaum et al., 1986). Rats were more sensitive to chlorophene than mice. The kidney was the main target organ in rats. In the 95-day study, increased absolute and relative kidney weights and reduced absolute and relative thymus weights were observed at ≥ 240 mg/kg bw/day in females and at 480 mg/kg bw/day in male and female rats. There seemed to be a dose-related increase in incidence and severity of nephropathy in male rats at ≥ 30 mg/kg bw/day, although the severity of the nephropathy was minimal to mild at these doses. Nephropathy of increasing severity (mild to moderate severity) was observed in 240 mg/kg bw/day male rats and in both male and female rats at 480 mg/kg bw/day. The incidence of nephropathy was significantly increased in male rats at ≥ 120 mg/kg bw/day (Birnbaum et al., 1986). These findings were suggested by the applicant not to be of significance since Fischer 344 male rats are known to spontaneously develop nephropathy. However, the concurrent vehicle controls of this study did not develop nephropathy during the course of the study and historical reference data were not included. A subchronic LO(A)EL of 120 mg/kg bw/day and NO(A)EL of 60 mg/kg bw/day for rats in this study is concluded. In the 95-day study mice first responded to treatment at 500 mg/kg bw/day with a decreased kidney weight in males and an increase in liver weight in females. At the same dose reduced body weight gain was observed in male mice. Among both rats and mice, males were more prone to chlorophene-induced nephropathy than females. In addition to the above mentioned repeated-dose toxicity studies a two-generation reproductive toxicity study (OECD 416) was recently conducted in Wistar rats (see section 3.8; confidential, 2008 / A6_8_2(3); doses: 60, 180, 540 mg/kg bw/day by oral gavage). This study was conducted according to internationally accepted guidelines and principles for GLP. In this study male Wistar rats of the parent generation (P generation) were exposed to chlorophene for at least 15 weeks corresponding to ~105 days. The LO(A)EL in this study was at the lowest dose tested, 60 mg/kg bw/day, based on increased absolute and relative kidney weights and microscopic kidney lesions (nephropathy, dilated tubules, basophilic tubules and lymphocytic infiltration) in P-males. **A two-year gavage study in mice** (NTP Technical Report 424, 1994, key study for carcinogenesis, 0; 120; 240; 480 mg/kg bw/day), with 3- and 15-month interim sacrifices revealed time- and dose-related increased severity and incidence of nephropathy in male and female mice, at occurring as early as 3 months after the beginning of chemical administration. Absolute kidney weights of high-dose males were lower than those of the controls at the 3-month interim evaluation. At the 15-month interim evaluation the absolute and relative kidney weights of dosed male mice were lower than those of the controls, as were the absolute kidney weights of dosed females. Kidney weights of all dosed males and females of the mid- and high dose group remained significantly reduced at the 2-year sacrifice. The final mean body weights of all dosed males and mid- and high dose females were lower than those of the controls. A chronic LO(A)EL of 120 mg/kg bw/day for mice were thus concluded based on severity and incidence of nephropathy and changes in kidney weights, whereas a conclusion on a NO(A)EL could not be taken due to effects on the lowest dose tested. **A two-year gavage study in rats** (NTP Technical Report 424, 1994, key study, 0; 30; 60; 120 mg/kg bw/d), with 3- and 15-month interim sacrifices revealed severe, time- and dose-related nephropathy in male and female rats, occurring as early as 3 months after the beginning of chemical administration. In male rats dosed for as long as 2 years, secondary hyper-parathyroidism developed, with

parathyroid gland hyperplasia, mineralization of the kidney and glandular stomach, and fibrous osteodystrophy occurring in the high-dose group. At the 3-month interim evaluation performed during this chronic gavage study absolute and relative kidney weights of male rats receiving 120 mg/kg bw/day and female rats receiving 240 mg/kg bw/day of chlorophene were significantly higher than those of the vehicle controls. Kidney damage is a common finding in aging male rats. However, since the kidney effects observed after chlorophene treatment are present in both male (Fischer 344 and Wistar) and female rats as well as in other species (rabbit, dog and mice), the observed kidney effects are not considered male rat specific. A NOAEL of 60 mg/kg bw/day in male Fischer rats in the 95-day study was concluded (kidney effects). However, in male Wistar rats in the 2-generation study the kidney effects was observed at the lowest dose tested (60 mg/kg bw/day), hence a **subchronic oral LO(A)EL of 60 mg/kg bw/day was concluded upon. The chronic oral NO(A)EL was not possible to determine since the LO(A)EL for rodents is proposed to be at the lowest dose tested, 30 mg/kg bw/day**, based on kidney effects at 30 mg/kg bw/day observed in male rats at 15 month interim evaluation and at the end of the two-year gavage study.

4.6.1 Summary and discussion of repeated dose toxicity

In rats, the target organ for chlorophene is the kidney. Increased kidney weights were noted in the **16-day oral gavage study** in male rats at ≥ 125 mg/kg bw/day. Nephropathy was observed in all animals of the highest dose group. The incidence and severity of nephropathy increased with dose. In mice, liver weights, but not kidney weights, were significantly increased at ≥ 250 mg/kg bw/day in an analogous subacute oral gavage study.

Three **subacute dermal studies** were conducted in rabbits. In the second 3-week study (confidential, 1985 / A6_3_2 (2)) a dose-dependent increase in histopathological changes was observed in the kidneys of female rabbits at ≥ 40 mg/kg bw/day of chlorophene. Moreover local reactions to treatment were observed in all animals treated with ≥ 40 mg/kg bw/day. In the first 3-week study (confidential, 1985 / A6_3_2(1)) local skin effects were observed in animals treated with ≥ 20 mg/kg bw/day of chlorophene. A tendency towards increased incidence and severity of kidney lesions was observed in female rabbits at the time of death (dead/killed during study + sacrificed at termination). In an additional 4-week study (confidential, 1989) slight to moderate skin effects were observed at 5 mg/kg bw/day and moderate to severe skin effects were observed at 25 mg/kg bw/day. Based on the results of these studies, **the subacute dermal NO(A)EL for systemic effects in rabbits is 25 mg/kg bw/day based on the kidney effects observed in female rabbit at 40 mg/kg bw/day (LO(A)EL)**. The NO(A)EL for local skin effects in rabbits is 1 mg/kg bw/day based on the slight to moderate skin effects observed at 5 mg/kg bw/day.

A series of **subchronic** and chronic **oral gavage studies in rodents** were performed. Rats were more sensitive to chlorophene than mice. The kidney was the main target organ in rats. Mice first responded to treatment with an increase in liver weight in female mice and a decreased kidney weight in male mice. Among rats and mice, males were more prone to chlorophene-induced nephropathy compared to the females. In Fischer 344 male rats there seemed to be a dose-related increase in incidence and severity of nephropathy at ≥ 30 mg/kg and the incidence of nephropathy was significantly increased at ≥ 120 mg/kg bw/day in the 95-day study, whereas increased absolute and relative kidney weights and microscopic kidney lesions were observed at 60 mg/kg bw/day with exposure duration of equivalent length in male Wistar rats in the 2-generation study. In addition to

the subchronic studies described, a **two-year oral study** was performed in rats (NTP). At the 3-month interim evaluation performed, absolute and relative kidney weights of male rats receiving 120 mg/kg bw/day and female rats receiving 240 mg/kg bw/day of chlorophene were significantly higher than those of the controls. In addition, severe, time- and dose-related nephropathy was observed in male and female rats. Based on the kidney effects observed in the male Wistar rats in the 2-generation study a **subchronic oral LO(A)EL for rodents of 60 mg/kg bw/days was concluded**. Moreover, a **chronic oral LO(A)EL for rodents of 30 mg/kg bw/day was concluded**, based on kidney effects at 30 mg/kg bw/day observed at the end of the two-year gavage study.

In a **subchronic oral study in Beagle dogs** (administration in gelatine capsules), weight loss was seen in the highest dose group (200 mg/kg bw/day). Relative weights of kidneys were significantly increased in a dose-dependent manner in male dogs at ≥ 30 mg/kg bw/day. In female dogs kidney weights were significantly increased at 100 mg/kg bw/day. Hyposthenuria was observed after 90 days in both sexes at 100 mg/kg bw/day. Based on the observed kidney effects, the **subchronic oral NO(A)EL for dogs was 10 mg/kg bw/day**.

4.6.2 Comparison with the CLP criteria

After evaluation of the studies on repeated-dose toxicity, a classification for specific target organ toxicity - repeated exposure (STOT-RE) is suggested for chlorophene based on the kidney effects observed in rats, dogs and rabbits.

According to CLP based on

- Fischer 344 rats in the oral 2-year study with a LO(A)EL of 30 mg/kg bw/day, and
- the 2-generation study in Wistar rats equivalent to a 90-day repeated-dose study with a LO(A)EL at the lowest dose tested of 60 mg/kg bw/day, and
- the 90-day study in Beagle dogs with a LO(A)EL of 30 mg/kg bw/day,

a classification as STOT-RE 2 is warranted. However, due to the results from a 3-week dermal study in rabbits STOT-RE 1 is proposed:

- The 3-week dermal study in rabbits with a LO(A)EL at 40 mg/kg bw/day warrants classification in category 1 (STOT-RE 1), after adjusting the LO(A)EL due to the study length according to CLP Annex 1 3.9.2.9.5.

The classification as STOT-RE 1 is in correspondence with CLP Annex 1, 2.9.2.7.3. stating:

Evidence from appropriate studies in experimental animals can furnish much more detail, in the form of clinical observations, haematology, clinical chemistry, and macroscopic and microscopic pathological examination, and this can often reveal hazards that may not be lifethreatening but could indicate functional impairment. Consequently all available evidence, and relevance to human health, shall be taken into consideration in the classification process, including but not limited to the following toxic effects in humans and/or animals:

Via undernote (d) significant organ damage noted at necropsy and/or subsequently seen or confirmed at microscopic examination.

4.6.3 Conclusions on classification and labelling:

The cut-off values for STOT RE Cat 1 are ≤ 60 mg/kg bw/d in the CLP regulation (in a 28-day dermal repeated dose study). Regarding the dose levels leading to toxicity as well as the quality of the findings it can be concluded that chlorophene is subject to classification for specific target organ toxicity - repeated exposure (STOT-RE) Category 1, based on the kidney effects observed in rabbits. **Regulation (EC) No 1272/2008 (CLP): STOT-RE 1; H372 Causes damage to kidneys through prolonged or repeated exposure**

4.7 Germ cell mutagenicity (Mutagenicity)

***In-vitro* genotoxicity of chlorophene**

Table 19: *In-vitro* genotoxicity of chlorophene

Test system, Method, Guideline	Organism/ strain(s)	Concentrations tested	Results		Remarks	Reference
			- S9	+ S9		
Ames test: Salmonella/ Microsome test No guideline, but \cong OECD 471 Non-GLP	<i>S. typhimurum</i> TA98, TA100, TA1535, TA1537	0.1, 0.3, 1.0, 3.0, 10.0, 33.0, 66.0, 100.0 μ g/plate (-S9 and +S9)	Neg. based on no increased mutation frequencies in the tested strains. The study did not include the recommended test strain (TA102) to reveal mutagenicity at A:T basepairs or of agents with crosslinking potential. The four strains used for testing reveal mutagenesis at G.C basepairs only.	Neg. based on no increased mutation frequencies in the tested strains. The study did not include the recommended test strain (TA102) to reveal mutagenicity at A:T basepairs or of agents with crosslinking potential. The four strains used for testing reveal mutagenesis at G.C basepairs only.	Cytotoxicity at ≥ 33 μ g/plate (-S9) and 100 μ g/plate (+S9)	Mortelmans <i>et al.</i> , 1986 A6_6_1 (1) KEY STUDY
Cytogenicity (Chromosomal aberrations) in Chinese hamster ovary (CHO) cells No guideline, but \cong OECD 473	CHO cells	-S9: 4, 8, 15, 30 and 60 μ g/mL +S9: 1.3, 2.5, 5, 10 and 20 μ g/mL	Neg. based on no increase in chromosome aberrations. Several study shortcomings: Only 100 of the required 200 metaphases were scored, insufficient incubation times with test compound with S9, no study with continuous treatment to	Neg. based on no increase in chromosome aberrations. Several study shortcomings: Only 100 of the required 200 metaphases were scored, insufficient incubation times with test compound with S9, no study with continuous treatment to	Complete inhibition of mitotic index at ≥ 30 μ g/mL without S9, and at 20 μ g/mL with S9	Confidential , 1994 A6_6_2 KEY STUDY

CLH REPORT FOR CHLOROPHENE

Test system, Method, Guideline	Organism/ strain(s)	Concentrations tested	Results		Remarks	Reference
			- S9	+ S9		
			confirm a negative result	confirm a negative result		
Sister chromatid exchanges (SCEs) in Chinese hamster ovary (CHO) cells No guideline Non-GLP	CHO cells	-S9: 0.5, 1.6, 5.0, 16.0 µg/mL +S9: 0.05, 0.16, 0.5, 1.6, 5.0, 16.0 µg/mL	Neg. based on that the SCE frequency did not increase with >= 20% above the concurrent solvent control level which was chosen as a statistically conservative positive response. Fifty second-division metaphase cells were scored at each dose level.	Neg. based on that the SCE frequency did not increase with >= 20% above the concurrent solvent control level which was chosen as a statistically conservative positive response. Fifty second-division metaphase cells were scored at each dose level.	Cytotoxicity: the highest non-lethal dose tested was 16.0 µg/mL	National Toxicology Program (1994) A6_7 (2b)
Mutagenicity test (HPRT) in murine lymphoma cells OECD 476 Exposure time: 3h Cytotoxicity determination: percent relative survival (% RS) adjusted for loss of cells during the 3h exposure period. Mutant frequency (Mf) is expressed as number of mutant cells per number of surviving cells	Mouse L5178Y cells	Exp. 1: 0, 2.5, 5, 10, 20, 25, 30, 35 µg/mL Exp. 2: 0, 5, 10, 20, 22.5, 25, 27.5, 30 µg/mL	Equivocal based on positive findings in on experiment (Exp. 1) that were not confirmed in a second experiment (Exp. 2). A concentration-related increased mutant frequency (Mf) was observed in Exp. 1 at 2.5-10 µg/mL, compared to control values. The Mf at 10 µg/mL was statistically different from the control, at recommended cytotoxicity levels of 10-20%. The tests for linear trend were not significant. The finding was not confirmed in Exp. 2. Exp. 1 -S9 Treatment -Mf - %RS	Neg. based on increased Mfs at concentrations eliciting unacceptably high levels of cytotoxicity. A concentration related increased Mf up to 20 µg/mL with significantly increased Mf at 20 µg/mL were observed in Exp. 2. The relative survival was however <50 % at 20 µg/mL. A test for linear trend was significant in Exp.2. This positive finding was not confirmed in Exp. 1. Exp. 1 +S9 Treatment -Mf - %RS	Cytotoxicity: >= 50% cytotoxicity at ≥ 25 µg/mL (-S9), ≥ 20 µg/mL (+S9)	Confidential , 2005 A6_6_3 (1) KEY STUDY

CLH REPORT FOR CHLOROPHENE

Test system, Method, Guideline	Organism/ strain(s)	Concentrations tested	Results		Remarks	Reference
			- S9	+ S9		
			0-4,84-100 2,5-6,3-96,98 5-7,22-95,64 10-11,41-88.47* 20-6,95-75,01 25-5,19-26,92 30-7,84-3,99 35-(10,16)-0,00 Exp. 2 -S9 Treatment -Mf – %RS 0-2,20-100 5-3,30-100,27 10-2,73-81,42 20-2,68-64,19 22,5-4,54-57,01 25-1,08-36,35 27,5-2,74-14,09 30-6,52-1,53	5-9,53-100,47 10-6,52-57,94 20-11,91-44,00 25-15,07-45,54 30-3,47-17,36 35-9,85-14,83 Exp. 2 -S9 Treatment -Mf – %RS 0-6,05-100 5-2,18-100,08 10-9,76-93,53 20-10,93-40,17* 25-6,29-47,31 27,5-9,99-31,47 30-8,48-8,42 35-5,88-1,48 + Linear trend test significant		
<p>Mutagenicity test (TK^{+/-}) in mouse L5178Y and human TK6 cells No guideline Non-GLP</p> <p>L5178Y: Soft agar Incubation time -S9 : 4 h</p> <p>TK6: Microwell plates Incubation time -S9: 20 h Only fast growing mutants (large colonies), i.e. not chromosomal mutants, were recorded.</p> <p>Cytotoxicity : Relative cell growth/plating efficiency during expression</p>	<p>Mouse L5178Y cells</p> <p>Human TK6 cells</p>	<p>L5178Y: 0, 10, 20, 25, 35, 45 µg/mL</p> <p>TK6 cells: 0, 10, 20, 30, 40 µg/mL</p>	<p>Equivocal in mouse L5178Y cells, An concentration-related increased Mf above control level in mouse L5178Y cells starting at the lowest concentration (10 µg/mL) tested, a conc. with satisfactory relative total growth (RTG; >=50%). L5178Y cells Treatment -Mf – RTG 0-48-100 10-55-ND 20-90-50 25-110-40 35-130-30 45-240-10 According to recommended global evaluation factors the increased Mf at 10 and 20 µg/mL relative to the vehicle/control levels are too low to allow the recommended</p>	/	<p>Reduction in relative total growth (RTG) ≥ 50% at 20 µg/mL for L5178Y, and at 40 µg/mL for TK6.</p>	<p>Caspary <i>et al.</i>, 1988 A6_6_3 (2) KEY STUDY</p>

CLH REPORT FOR CHLOROPHENE

Test system, Method, Guideline	Organism/ strain(s)	Concentrations tested	Results		Remarks	Reference
			- S9	+ S9		
			trend analyses. Uninformative in human TK6 cells. Background Mf too low compared to the recommended assay acceptance criteria. Increased mutant frequencies in human TK6 cells at cons. >= 40 µg/mL with significantly reduced RTG. TK6 cells Treatment -Mf – RTG 0-25-100 10-60-99 20-50-97 30-45-95 40-80-20			

In vivo

Table 20: In vivo genotoxicity of chlorophene

Type of test, Method/ Guideline	Species, Strain, Sex, no/group	Frequency / route of application	Sampling times	Dose levels	Results	Reference
Micronucleus assay OECD 474	Mice CD-1 ♂ + ♀ 5-15/sex Vehicle control: 15♂, 15♀ 250 mg/kg: 5 ♀ 500 mg/kg: 5 ♂, 5 ♀ 1000 mg/kg: 5 ♂, 15 ♀ 2000 mg/kg: 15 ♂ Positive control: 5 ♂, 5 ♀	Single dose, oral	24, 48 and 72 h	500, 1000, 2000 mg/kg (♂) 250, 500, 1000 mg/kg (♀)	Neg. based on that the frequency of micronucleated polychromatic (immature) and mature erythrocytes were generally similar to those in concurrent vehicle controls at any sampling time. There were no indications of bone marrow toxicity. The individual proportions of immature/mature cells were unaffected by treatment. The statistical test used was the Mann-Whitney U-test. Males (2) and females (1) in the highest dose groups showed signs of toxicity with hunched posture, piloerection, closed eyes and reduced motor activity and were sacrificed <i>in extremis</i> .	Confidential, 1990 A6_6_4 (1) KEY STUDY

CLH REPORT FOR CHLOROPHENE

Type of test, Method/ Guideline	Species, Strain, Sex, no/group	Frequency / route of application	Sam-pling times	Dose levels	Results	Reference
					The vehicle control Mfs (mean micronucleated cells/1000) for immature erythrocytes were 0,7 and 0.4 for males and females, respectively and the levels of treated animals never exceeded 1.8 (males and females) compared with the level induced by the positive control (chlorambucil) of 57.8 (males) and 50.7 (females).	
Dominant-lethal test No guideline Non-GLP	Mice ♂ no further details	Single dose, i.p.	-	100, 200 mg/kg	Neg. based on that chlorophene did not induce dominant lethal mutations in germ cells of male mice.	Confidential, 1972 A6_6_4 (2)
<i>In vivo</i> comet assay No guideline “Quasi” guideline in Tice et al., 2000 Single Cell Gel/Comet Assay: Guidelines for In Vitro and In Vivo Genetic Toxicology Testing. <i>Envir. And Mol. Mutagenesis</i> , 35, p206-221. GLP	Mice ICR (CD-1) ♂ Preliminary toxicity test: 2/dose Main test: 7/dose	Two doses: the first at 0h and the second at 20h. Gavage	4h and 24 h after initial dose	Initial preliminary toxicity test using DMSO as solvent: 200, 300, 400, 2000 Preliminary toxicity test using corn oil as solvent: 200, 300, 400, 2000 mg/kg bw Main test using corn oil as solvent: 90, 180, 360 mg/kg bw	Neg. based on no increased DNA damage levels in the organ tested. Mortality was observed with DMSO as solvent at 400 mg/kg. The solvent was changed to corn oil and mortality was observed at 2000 mg/kg. The experiment did not include studies of the target organ (site of contact) for chlorophene, the kidney, as specified to the industry prior to onset of the experiment. There was no evidence of an increase in the percentage tail intensity values in single cell/nuclei suspensions from liver and glandular stomach, or bone marrow from animals dosed with the test item dose groups when compared to the concurrent vehicle control group. The positive control produced an acceptable increase in the percentage tail intensity value in all tissues scored. The presence of clinical signs indicated that systemic absorption had occurred.	Confidential, 2009 A6_6_5 KEY STUDY

4.8.3 Summary and discussion of mutagenicity

Conclusion on the genotoxicity of chlorophene was equivocal, in the *in vitro* tests in bacteria and mammalian cells.

Several of the *in vitro* studies (Mortelmans et al., 1986; confidential, 1994 / A6_6_2) exhibit study insufficiencies that reduce their power to conclude with chlorophene being not genotoxic. In two independent *in vitro* mutagenicity studies in mammalian cells (mouse L5178Y cells) assessing mutagenesis in two different loci (HPRT and TK) there were indications of increased mutation frequencies without metabolic activation (confidential, 2005 / A6_6_3(1); Caspary et al., 1988). The first study is a well-conducted study following OECD Guideline 476 (1997), and the latter study is a non-guideline, non-GLP TK^{+/-} assay conducted with chlorophene of unknown specification (Caspary et al., 1988).

In vivo there were no indications of clastogenicity or aneugenicity in the micronucleus assay in mice. The systemic availability of the test compound to the bone marrow is however in general questioned with negative results in micronucleus test, particularly when no bone marrow toxicity was observed in any of the exposed groups. An agreement with the applicant for conducting a second *in vivo* genotoxicity assay (*in vivo* comet assay) in mice was made (confidential, 2009 / A6_6_5). However assessment of genotoxic effect in the target organ, the kidney, was not performed, leaving the study of low power to conclude on the potential genotoxic properties of chlorophene in relevant tissues. The comet assay was negative for the tested organs. The dominant-lethal test (confidential, 1972 / A6_6_4 (2)) is only available as a summary in which a negative result is reported.

No systematic search for structure-activity relationships to known germ cell mutagens were conducted.

4.8.4 Conclusions on classification and labelling

In summary, several of the key studies exhibit study insufficiencies that hamper establishment of solid conclusions on genotoxicity, but based on an overall evaluation of the available data using a *Weight of Evidence* approach the decision on the genotoxicity is negative.

No classification is proposed.

4.8 Carcinogenicity

Table 21: Carcinogenicity studies

Type of study, Method/ Guideline	Species, Strain, Sex, no/group	Dose levels, frequency of application	Tumours	Reference
Two-year gavage study US-EPA 83-2 ≅ OECD 453	Rat F344/N ♂+♀ 50/sex 3-month and 15-month interim sacrifice: 10/sex	0, 30, 60, 120 mg/kg/day (♂) 0, 60, 120, 240 mg/kg/day (♀)	♂: No evidence of carcinogenic effect in ♂ receiving 30, 60, or 120 mg/kg day. ♀: Equivocal evidence of carcinogenic effect in ♀ based on the occurrence of <u>two very rare renal transitional cell carcinomas; one occurring at 120 mg/kg/day and one at 240 mg/kg/day.</u> No changes in survival or mean body weights in control or exposed groups.	National Toxicology Program Technical Report Series No.424 (1994) A6_5+A6_7 KEY STUDY*
Two-year gavage study US-EPA 83-2 ≅ OECD 451	Mouse B6C3F ₁ ♂ + ♀ 50/sex 3-month and 15-month interim sacrifice: 10/sex	0, 120, 240, 480 mg/kg/day	♂: Some evidence of carcinogenic effect in ♂ based on increased incidences of renal tubule adenoma in high dose and renal tubule adenoma or carcinoma (combined) at mid- and high-dose. ♀: No evidence of carcinogenic activity in ♀. Survival of high-dose male and female mice was lower than that of the controls. The number of mice surviving to the end of the study was considered adequate for evaluation of chronic toxicity and carcinogenicity. Final mean body weights of all dosed males and mid- and high dose females were lower than those of the controls.	National Toxicology Program Technical Report Series No.424 (1994) A6_7 (1) KEY STUDY*
One-year dermal initiation/promotion study No guideline	Mouse Swiss CD-1 ♂ + ♀ 50/sex	As initiator: single application of 10 mg/animal As promoter: 0.1, 1.0, 3.0 mg/animal, 3 times/week As complete carcinogen: 10 mg single initiating dose followed by 0.1, 1.0, 3.0 mg/animal 3 times/week	No skin tumour-initiating potential. Weak tumour-promoting activity. There was a dose-related increased incidence in papillomas in both males and females. Chlorophene did not act as a complete carcinogen.	National Toxicology Program (1995) A6_7 (3)
20-week dermal carcinogenicity study in transgenic	Mouse Tg.AC mice	0.1, 1.0, 3.0 mg/animal	A significant (p<0.01) carcinogenic effect on skin were recorded at 3 mg/animal	Spalding JW, French, Tice,

CLH REPORT FOR CHLOROPHENE

Type of study, Method/ Guideline	Species, Strain, Sex, no/group	Dose levels, frequency of application	Tumours	Reference
Tg.AC mice No guideline	♀ 13-20 animals/group	3 times per week Post-exposure period: 6-10 weeks	measured as incidence of animals with tumours analysed by life-table analysis. Tumour multiplicity was also significantly (p<0.01) increased in the high dose group. The mean latency time to maximum skin tumour yield was the same for the negative and positive control groups and for the highest chlorophene dose group. Survival at 20 weeks decreased with increasing doses (87%, 77% and 68% in the low-, medium and high-dose groups, respectively) No overt skin irritation and ulceration was reported after monitoring by gross examination.	Furedi-Machacek, Haseman and Tennant. Tox. Sci., 49, 241-254, 1999 A6_7 (2) KEY STUDY

*Studies also reported in Table 3.7 (chronic toxicity).

The evaluation of carcinogenicity is based on four studies, see Table above.

The carcinogenic potential of chlorophene was studied in **two-year gavage studies in rats and mice**. The studies were conducted by the National Toxicology Program.

Rats, gavage, two year (NTP):

In rats, very rare renal transitional cell carcinomas occurred in one female rat each of the mid- and high dose group (table 4.14) whereas none of the tumours found in male rats could be ascribed as an effect of the test substance (table 4.15) Due to the low incidence, this finding is considered as **equivocal evidence** for a carcinogenic activity of the test compound in female rats. Survival and mean body weights of dosed animals were similar to those of the controls. Non-neoplastic endpoints comprised severe time- and dose-related nephropathy in male and female rats, occurring as early as 3 months after the beginning of chemical administration (females). In male rats dosed for as long as 2 years, secondary hyper-parathyroidism was observed, with parathyroid gland hyperplasia, mineralization of the kidney and glandular stomach, and fibrous osteodystrophy occurring in the high-dose group. The severity of these lesions was greater in males.

CLH REPORT FOR CHLOROPHENE

Table 22: Nephropathy and pathological findings in female rats

Dose (mg/kg)	0	60	120	240
3-Month Interim Evaluation				
Kidney ^a	10	10	8	9
Nephropathy ^b	1 (0.1) ^c	3 (0.3)	3 (0.4)	7** (1.2)**
15-Month Interim Evaluation				
Kidney	10	10	10	10
Nephropathy	9 (0.9)	10 (1.2)	9 (1.1)	10 (1.8)**
2-Year Evaluation				
Kidney	50	50	51	50
Nephropathy	46 (1.2)	47 (1.2)	50 (1.5)*	50 (2.4)**
Single Sections (Standard Evaluation)				
Kidney	50	50	51	50
Renal Tubule Hyperplasia	0	0	0	1
Renal Tubule Adenoma ^d	0	0	0	1
Transitional Cell Carcinoma ^e	0	0	1	1
Step Sections (Extended Evaluations)				
Kidney	50	- ^f	-	50
Renal Tubule Hyperplasia	2	-	-	2
Single and Step Sections combined				
Kidney	50	-	-	50
Renal Tubule Hyperplasia	2	-	-	3
Renal Tubule Adenoma	0	-	-	1
Transitional Cell Carcinoma	0	-	-	1

* Significantly different (P≤0.05) from control group by Fisher exact test; severity significantly different by Mann-Whitney U test

** P≤0.01

^a Number of animals with kidney examined microscopically

^b Number of animals with lesion

^c Average severity grade of lesion in all animals: 0=none, 1=minimal, 2=mild, 3=moderate, 4=marked

^d Historical incidence for 2-year corn oil gavage studies with vehicle control groups: 2/1,068 (1.9% ± 0.2%); range 0%-2%

^e Historical incidence: 0/1,068

^f Animals not examined in extended or combined evaluations

CLH REPORT FOR CHLOROPHENE

Table 23: Nephropathy and pathological findings in male rats

Dose (mg/kg)	0	30	60	120
3-Month Interim Evaluation				
Kidney ^a	10	10	10	9
Nephropathy ^b	10 (1.0) ^c	8 (0.8)	9 (1.0)	8 (0.9)
15-Month Interim Evaluation				
Kidney	10	10	10	9
Nephropathy	10 (1.7)	10 (2.1) ^o	10 (2.1) ^o	9 (2.2) ^o
2-Year Evaluation				
Kidney	50	49	50	50
Nephropathy	48 (2.3)	48 (2.8) ^o	48 (2.9) ^{oo}	50 (3.3) ^{oo}
Single Sections (Standard Evaluation)				
Kidney	50	49	50	50
Renal Tubule Hyperplasia	0	2	0	2
Renal Tubule Adenoma	1	0	0	0
Renal Tubule Carcinoma	0	0	0	1
Renal Tubule Adenoma or Carcinoma ^d	1	0	0	1
Step Sections (Extended Evaluations)				
Kidney	50	49	50	50
Renal Tubule Hyperplasia	3	7	6	17 ^{oo}
Renal Tubule Adenoma	0	1	2	1
Renal Tubule Carcinoma	0	0	0	1
Renal Tubule Adenoma or Carcinoma	0	1	2	2
Single and Step Sections combined				
Kidney	50	49	50	50
Renal Tubule Hyperplasia	3	9	6	17 ^{oo}
Renal Tubule Adenoma	1	1	2	1
Renal Tubule Carcinoma	0	0	0	1
Renal Tubule Adenoma or Carcinoma	1	1	2	2

^o Significantly different (P≤0.05) from control group by Fisher exact test; severity significantly different by Mann-Whitney U test

^{oo} P≤0.01

^a Number of animals with kidney examined microscopically

^b Number of animals with lesion

^c Average severity grade of lesion in all animals: 0=none, 1=minimal, 2=mild, 3=moderate, 4=marked

^d Historical incidence for 2-year corn oil gavage studies with vehicle control groups: 12/1,069 (1.1% ± 1.4%); range 0%-4%

Mice, gavage, two year (NTP):

In mice following exposure to chlorophene by gavage, there was some evidence of carcinogenic effect in males based on significant increase in renal tubule adenomas in the 480 mg/kg (high-dose) group of males and of renal tubule adenomas or carcinoma in both the 240 (mid-dose) and 480 mg/kg (high-dose) groups of males (table 25).

Table 24: Nephropathy and pathological findings in male mice

Dose (mg/kg)	0	120	240	480
3-Month Interim Evaluation				
Kidney ^a	10	10	10	10
Nephropathy ^b	1 (0.1) ^c	3 (0.3)	10 ^{oo} (1.2) ^{oo}	10 ^{oo} (2.2) ^{oo}
15-Month Interim Evaluation				
Kidney	10	10	10	10
Nephropathy	9 (0.9)	10 (2.7) ^{oo}	10 (2.7) ^{oo}	10 (2.7) ^{oo}
2-Year Evaluation				
Kidney	50	50	50	50
Nephropathy	39 (0.8)	48 ^{oo} (2.0) ^{oo}	50 ^{oo} (2.4) ^{oo}	49 ^{oo} (2.4) ^{oo}
Single Sections (Standard Evaluation)				
Kidney	50	50	50	50
Renal Tubule Hyperplasia	0	0	3	6 ^{oo}
Renal Tubule Adenoma ^d	0	2	2	2
Renal Tubule Carcinoma ^e	0	0	2	1
Renal Tubule Adenoma or Carcinoma ^f	0	2	4 ^o	3
Step Sections (Extended Evaluations)				
Kidney	50	50	50	50
Renal Tubule Hyperplasia	9	16 ^o	13	9
Renal Tubule Adenoma	0	1	2	3
Renal Tubule Carcinoma	0	0	1	0
Renal Tubule Adenoma or Carcinoma	0	1	3	3
Single and Step Sections Combined				
Kidney	50	50	50	50
Renal Tubule Hyperplasia	9	16 ^o	14	13
Renal Tubule Adenoma	0	2	4	5 ^o
Renal Tubule Carcinoma	0	0	2	1
Renal Tubule Adenoma or Carcinoma	0	2	6 ^{oo}	6 ^{oo}

^o Significantly different (P≤0.05) from control group by Fisher exact test; severity significantly different by Mann-Whitney U test

^{oo} P≤0.01

^a Number of animals with kidney examined microscopically

^b Number of animals with lesion

^c Average severity grade of lesion in all animals: 0=none, 1=minimal, 2=mild, 3=moderate, 4=marked

^d Historical incidence for 2-year corn oil gavage studies with vehicle control groups (mean ± standard deviation): 4/949 (0.4% ± 1.0%); range 0%-2%

^e Historical incidence: 0/949

^f Historical incidence: 4/949 (0.4% ± 1.0%); range 0%-2%

Tubular hyperplasia was seen in all treated groups with significant increase in low-dose males but without a dose-response. This study featured gavage administration five days per week of 120, 240, 480 mg/kg bw/day of chlorophene in corn oil to B6C3F1 mice. Treatment was continued for 2 years, with interim sacrifices at 13 and 65 weeks. The 2-year sacrifice comprised 50 animals per sex and dose, whereas each interim sacrifice comprised 10 animals per sex and dose. General toxicity was evident as nephropathy (table 4.16) and reduced body weight development (table 4.17). The

incidence and severity of nephropathy were both age- and dose-related. Nephropathy was first statistically significant in the mid-dose group at the 13-week sacrifice with 10 out of 10 males (8/10 females; 240 mg/kg) all showing minimal nephropathy. At the 65-week sacrifice, nephropathy had progressed to a mild-to-moderate degree and was seen in all treated groups with a dose-related increase in severity of nephropathy and affecting both males and females. Nephropathy was evident as interstitial fibrosis, multifocal dilated tubules with flattening of the renal tubule epithelium, regenerative tubules with basophilic epithelium, thickened basement membranes and hyaline casts.

Table 25: Body weights and absolute and relative organ weights of mice

	Vehicle Control	120 mg/kg	240 mg/kg	480 mg/kg
Male				
n	45	32	38	30
Necropsy body weight	48.0 ± 1.0	39.1 ± 0.9 ^{oo}	35.5 ± 0.7 ^{oo}	32.6 ± 0.7 ^{oo}
Brain				
Absolute	0.453 ± 0.004	0.465 ± 0.005	0.473 ± 0.008 ^o	0.469 ± 0.005 ^{ob}
Relative	9.66 ± 0.24	12.08 ± 0.27 ^{oo}	13.48 ± 0.32 ^{oo}	14.51 ± 0.29 ^{oo} ^b
L. Kidney				
Absolute	0.399 ± 0.008	0.263 ± 0.005 ^{oo}	0.253 ± 0.006 ^{oo}	0.267 ± 0.007 ^{oo}
Relative	8.45 ± 0.22	6.80 ± 0.15 ^{oo}	7.20 ± 0.21 ^{oo}	8.26 ± 0.25
R. Kidney				
Absolute	0.415 ± 0.006	0.280 ± 0.005 ^{oo}	0.266 ± 0.006 ^{oo}	0.281 ± 0.005 ^{oo}
Relative	8.80 ± 0.20	7.24 ± 0.15 ^{oo}	7.57 ± 0.21 ^{oo}	8.71 ± 0.23
Liver				
Absolute	2.424 ± 0.117	2.398 ± 0.202	1.949 ± 0.078 ^o	2.113 ± 0.056 ^o
Relative	51.25 ± 2.73	64.56 ± 7.14	55.66 ± 2.80	65.35 ± 1.95 ^o
Female				
n	36	40	33	25
Necropsy body weight	47.2 ± 1.3	44.9 ± 1.2	40.4 ± 1.2 ^{oo}	33.7 ± 0.9 ^{oo}
Brain				
Absolute	0.477 ± 0.004	0.473 ± 0.003	0.465 ± 0.003 ^o	0.462 ± 0.005 ^o
Relative	10.39 ± 0.32	10.85 ± 0.31	11.88 ± 0.39 ^{oo}	13.94 ± 0.44 ^{oo}
L. Kidney				
Absolute	0.277 ± 0.004	0.261 ± 0.005	0.231 ± 0.004 ^{oo}	0.206 ± 0.014 ^{oo}
Relative	5.99 ± 0.17	5.96 ± 0.20	5.90 ± 0.21	6.35 ± 0.72
R. Kidney				
Absolute	0.291 ± 0.005	0.276 ± 0.005 ^o	0.250 ± 0.005 ^{oo}	0.214 ± 0.006 ^{oo}
Relative	6.29 ± 0.16	6.29 ± 0.20	6.38 ± 0.22	6.43 ± 0.22
Liver				
Absolute	1.970 ± 0.077	2.223 ± 0.123	2.208 ± 0.090	2.121 ± 0.089
Relative	42.38 ± 1.66	52.14 ± 4.24 ^o	56.67 ± 3.53 ^{oo}	63.13 ± 2.34 ^{oo}

^o Significantly different (P≤0.05) from the control group by Williams' or Dunnett's test

^{oo} P≤0.01

^a Organ weights and body weights are given in grams; organ-weight-to-body-weight ratios are given as mg organ weight/g body weight (mean ± standard error).

^b n=29

The survival of chlorophene-treated animals decreased following chlorophene exposure (table 4.18). Survival of dosed male mice was lower than that of controls (90%) with 69% (p=0.014), 81% (p=0.222) and 64% (p=0.002) in the low-, mid- and high dose groups, respectively. Survival of female mice was 74% for controls and 85%, 69% (p=0.665) and 51% (p=0.007) in the low-, mid- and high-dose groups, respectively. The number of mice surviving to the end of the study was

considered adequate for evaluation of chronic toxicity and carcinogenicity. The decreased survival was associated in part with dose-related increases in incidence and severity of nephropathy. The mean body weights at necropsy of all dosed males and mid- and high-dose females were lower than those of the controls. Liver and kidney weights were dose-dependently increased in both sexes.

Table 26: Survival of mice

	Vehicle Control	120 mg/kg	240 mg/kg	480 mg/kg
Male				
Animals initially in study	70	70	70	70
3-Month interim evaluation ^a	10	10	10	10
15-Month interim evaluation ^a	10	10	10	10
Moribund	2	12	7	8
Natural deaths	3	4	5	10
Accidental deaths ^a		2		2
Animals surviving to study termination	45	32	38	30
Percent probability of survival at end of study ^b	90	69	81	64
Mean survival (days) ^c	591	551	572	530
Survival analysis ^d	P=0.007	P=0.014	P=0.222	P=0.002
Female				
Animals initially in study	70	70	70	70
3-Month interim evaluation ^a	10	10	10	9
15-Month interim evaluation ^a	10	10	10	9
Moribund	9	4	10	12
Natural deaths	5	3	5	14
Accidental deaths ^a		3	2	1
Animals surviving to study termination	36	40	33	25
Percent probability of survival at end of study ^b	74	85	69	51
Mean survival (days) ^c	583	554	551	520
Survival analysis ^d	P<0.001	P=0.314N	P=0.665	P=0.007

^a Censored from survival analyses

^b Kaplan-Meier determinations

^c Mean of all deaths (uncensored, censored, and terminal sacrifice)

^d The result of the life table trend test (Tarone, 1975) is in the control column, and the results of the life table pairwise comparisons (Cox, 1972) with the controls are in the dosed columns. A lower mortality in a dose group is indicated by N.

Mice, dermal:

Female transgenic Tg.AC mice carrying an inducible v-Ha-ras oncogene were exposed to 0.1, 1.0, 3.0 mg/animal by the dermal route 3 times per week for 20 weeks followed by a 6 week post-exposure period (Spalding et al., 1999; A6_7 (2)), (table 4.19) They used 13-20 animals/group at the age of 7-18 weeks. Doses of 14.3, 143, 429 µg/cm² BPA solved in acetone (0.5, 5, 15 mg/mL concentrations, total volume of 0.2 mL/animal) were applied to a dorsal interscapular skin area of 5-7 cm². Positive control used was 12-O-tetradecanoylphorbol-13-acetate (TPA). Body weights were recorded weekly and cage observations were made twice a day for mortality and morbidity. Detailed clinical observations 2-5 times weekly. Overt skin irritation and ulceration was monitored by gross examination. Numbers of skin tumours were recorded weekly. Evaluations of gross

appearance of epidermal papillomas were done, and in order to be counted as a positive response, skin tumours had to reach 1 mm in size and persist for at least 3 weeks. Once an equivocally positive skin response to either the test agent or positive control (TPA) had been established, further observations became unnecessary. Survival decreased in a dose-dependent manner at 20 weeks with 87%, 77% and 68% in the low-, medium and high-dose groups, respectively. **A significant ($p < 0.01$) carcinogenic effect was recorded** at 3 mg/animal measured as the incidence of animals with tumours, analysed by life-table analysis. **Tumour multiplicity was also significantly ($p < 0.01$) increased** in the high-dose group. The mean latency time to maximum tumour yield was the same for the negative and positive control groups and for the high-dose group.

Table 27: Skin Tumour Incidence in homozygous female Tg.AC mice

Treatment	Incidence of animals with tumors (%)	Mean weeks to first tumor (\pm SD)	Mean tumors/animals at risk (\pm SD) ^a	Mean weeks to maximal tumor yield (\pm SD)	Survival at 20 weeks (%)
Acetone	4/14 (29%)	13.3 \pm 2.1	0.3 \pm 0.0	13.3 \pm 2.1	14/14 (100%)
BCP ^b					
0.1 mg	5/15 (33%)	8.6 \pm 6.1	0.7 \pm 1.2	16.0 \pm 1.9	13/15 (87%)
1.0 mg	1/13 (8%)	16.0	0.1	16.0	10/13 (77%)
3.0 mg	16/19 (84%) ^c	10.9 \pm 3.6	2.3 \pm 1.9 ^d	13.5 \pm 3.6	13/19 (68%)
TPA ^e					
1.25 μ g	19/20 (95%) ^c	7.4 \pm 2.4	19.5 \pm 12.4 ^d	13.6 \pm 3.5	9/20 (45%)

Note. Mice were 18 weeks old at start of treatment.

^a Animals were considered to be at risk after 10 weeks of dosing.

^b o-Benzyl-p-chlorophenol was dissolved in acetone and applied 3 times per week

^c $p < 0.01$ vs. acetone controls (Life Table Test).

^d $p < 0.01$ vs. acetone controls (Mann-Whitney U-Test).

^e TPA (1.25 μ g) in 200 μ L acetone was applied twice per week.

In a study with male and female CD-1 mice the potential to **initiate or promote skin tumours** was investigated in a 20 week dermal initiation/promotion study using phorbol ester (TPA) or dimethyl-benz-anthracene (DMBA) as model promoter and initiator, respectively (NTP, 1995, A6_7 (3)). **Chlorophene acted as a weak skin tumour promoter**, and did not demonstrate activity as a skin tumour initiator or a complete carcinogen.

4.8.1 Summary and discussions of carcinogenicity

Rats, gavage, two year (NTP):

In the 2-year study (table 4.14 and table 4.15), F344 rats received chlorophene by gavage administrations of 30, 60, 120 (males) or 60, 120, 240 (females) mg/kg bw/day in corn oil. Treatment was continued for 2 years, with interim sacrifices at 13 and 65 weeks. The 2-year sacrifice comprised 50 animals per sex and dose, whereas each interim sacrifice comprised 10 animals per sex and dose. General toxicity was restricted to kidney effects (nephropathy and effects on urine parameters).

The findings in female rats included very rare carcinomas of the renal transitional epithelium (TCC). One female each of the mid- and high-dose group were found to carry this type of tumour. This type of tumour has not been recorded in NTP historical control data (0/1068). The rarity of this tumour type raises concern, since the tumour occurred twice in this study, which reduces the possibility that the tumours occurred by chance. The likelihood that the two TCCs observed in this study are of spontaneous origin is very low. Considerations of tumour type and incidence of tumour

type relative to historical control incidence is highlighted in the R7 Guidance to CLP (3.6.2.2.6): “*Historical control data can also be useful to judge the biological significance of marginal increases in uncommon tumours. If there is a small increase in a particular tumour type which historical data shows to be very uncommon and unlikely to have occurred by chance then this may support a conclusion of carcinogenicity without the requirement for a statistically significant increase.*”

Another consideration is the evaluation of whether there is any indication that the observed tumours are substance related, such as increased incidences of pre-neoplastic lesions. A review was conducted within the NTP study to specifically evaluate the transitional cell hyperplasia. This review of high-dose and vehicle control rats from the 15-month interim evaluation and 2-year study was limited to the transitional epithelium lining, the renal pelvis, and papilla. An increased incidence of transitional cell hyperplasia was detected in both high-dose females and males after 15 months and 2 years (the two time points not differentiated in the report). Thus, the observed tumours were accompanied by increases in pre-neoplastic lesions, but the existence of hyperplasias *per se* is not on its own predictive of the development of tumours. Sex-differences seem to exist.

Moreover, since the tumour type (Transitional cell carcinoma) is relevant for humans, their occurrence should be included in the overall evaluation of the carcinogenicity of chlorophene.

Mice, gavage, two year NTP: It is logical to speculate that the renal hyperplasias and neoplasms observed in male mice following orally administered chlorophene are secondary to the nephrotoxicity exerted by the test compound. As discussed in Marsman et al., 1995, previous studies in mice with other chemicals, both genotoxic and non-genotoxic

(bromochloromethane CAS number 74-27-5; nitrilotriacetic acid CAS number 139-13-9; tri(2,3-dibromopropyl)phosphate CAS number 126-72-7; 2,4-diaminophenol dihydrochloride CAS number 137-09-7) showed poor association between nephropathy and renal carcinogenicity. Thus, although nephropathy may be a permissive factor, other primary and secondary mechanisms may be operative in the induction of the mouse renal neoplasms.

In lifetime bioassays compounds are routinely tested using at least three dose levels to enable hazard identification and hazard characterisation as part of risk assessment. Of these doses, the highest dose is meant to provoke minimal toxicity, characterised by overt toxicity, toxicity believed to reduce the life span or an approximately 10% reduction in body weight gain (maximal tolerated dose, MTD dose). The MTD is the highest dose of the test agent during the bioassay that can be predicted not to alter the animal’s normal longevity from effects other than carcinogenicity. In the current study the low-dose males express sufficient toxicity to qualify as an actual high-dose group, and in the low-dose animals there is an increase in renal neoplasms, and in the mid- and high-dose the increase is statistically significant. Thus there appears to be a dose-response relationship between the dose of chlorophene and the development of renal neoplasms, starting at the lowest dose. The increase in renal tubule hyperplasia was also significantly increased in the low-dose males. Altogether these findings support a classification of the test compound in Category 2 (According to **Regulation (EC) No 1272/2008 (CLP)**).

Other compounds have been classified as Category 2 carcinogens on the basis of tumours occurring at doses at and exceeding MTD, in which the incidence of tumours showed a dose-response relationship (*e.g.*, Polyhexamethylene biguanide– PHMB, CAS number 27083-27-8). The data on genotoxicity were limited, and PHMB was concluded as non-genotoxic.

Mice, dermal: Transgenic female Tg.AC mice

The study in the transgenic Tg.AC mice suggests that chlorophene is carcinogenic, although dose-dependent decreased survival was observed at 20 weeks. Dermal dosing of female Tg.AC mice with 3 mg chlorophene in acetone over 20 weeks led to a significant increase of skin tumours (84% of animals had tumours) compared to the acetone control (29%).

Considerations of the relevance of the model used, for studies with regulatory purposes:

As stated in R7.7.10 of the document “Guidance on information requirements and chemical safety assessment R.7a: Endpoint specific guidance“ related to the REACH Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006, data from transgenic rodent models can be used for assessing the carcinogenicity of a chemical: **“Genetically engineered (transgenic) rodent models** (e.g., *Xpa*^{-/-}, *p53*^{+/-}, *rasH2* or Tg.AC): *animals can be genetically engineered such that one or more of the molecular changes required for the multi-step process of carcinogenesis has been accomplished (Tennant et al., 1999). This can increase the sensitivity of the animals to carcinogens and/or decrease the latency with which spontaneous or induced tumours are observed. The genetic changes in a given strain of engineered animals can increase sensitivity to carcinogenesis in a broad range of tissues or can be specific to the changes requisite for neoplastic development in one or only a limited number of tissues (Jacobson-Kram, 2004; Pritchard et al., 2003; ILSI/HESI 2001). Data from these models may be used in a Weight of Evidence analysis of a chemical’s carcinogenicity.*”

Although studies of carcinogenicity using the Tg.AC model are not yet fully validated for regulatory purposes, extensive investigations have already been conducted. In the study with chlorophene included here (Spalding et al., 1999) several compounds were tested and the results showed good correlation with results from 2-year bioassays, with 3/6 compounds including chlorophene showing clear carcinogenic activity, and 3/6 that were negative in Tg.AC mice, of which one non-genotoxic compound have shown to induce liver tumours in female mice in the 2-year bioassay. A more extensive array of compounds have been tested in the Tg.AC model and the results have been compared with results from the 2-year bioassay (Eastin et al., 1998; Pritchard et al., 2003), showing good concordance, and that the model is able to detect also non-genotoxic carcinogens.

4.9.2 Comparison with the CLP criteria

The “Guidance on the Application of Regulation (EC) No 1272/2008” (Guidance to CLP) 3.6.2.2.6 lists important factors, which may be taken into consideration when assessing the overall level of concern of possible carcinogenic compounds. In the following sections chlorophene have been evaluated in relation to these factors:

(a) tumour type and background incidence; The observed tumour types (renal transitional cell carcinomas in female rats, the renal tubular adenomas and carcinomas in male mice and skin tumours in mice) are relevant for humans, and they arose from histologically unrelated tissues, although vicinal, suggesting that chlorophene may lead to more than one type of tumour.

With respect to the two TCCs observed in female rats, the Guidance to CLP states “If there is a small increase in a particular tumour type which historical data shows to be very uncommon and unlikely to have occurred by chance then this may support a conclusion of carcinogenicity without the requirement for a statistically significant increase”.

(b) multi-site responses; It is stated in the Guidance to CLP: “Evidence shows that substances which cause tumours in either multiple sites and/or multiple species tend to be more potent carcinogens than those causing tumours at only one site in one species (Dybing et al., 1997). This is often true for substances which are mutagenic. Also, where human carcinogens have been tested in two or more species, the majority have caused cancer in several species (Tennant, 1993). Thus, if a substance causes tumours at multiple sites and/or in more than one species then this usually provides strong evidence of carcinogenicity.”

The kidney tumours appeared at two different, but vicinal sites, and they arose from histologically unrelated tissues and occurred in two species: TCCs in female rats and renal tubular adenomas and carcinomas in male mice. Besides these skin tumours occurred in mice. The occurrence of tumours at different sites suggests that chlorophene may be carcinogenic.

(c) progression of lesions to malignancy; In male mice there seems to be a low but treatment-related increase in benign renal adenomas, but not in malignant renal carcinomas. The number of tumours may be too low for progression from adenomas to carcinomas to be evident.

(d) reduced tumour latency; The latency of tumour development, i.e. how quickly a substance induces tumours, often reflects the potency of a carcinogen. No tumours were found at the interim sacrifices in the studies in mice and rats and the mean latency time to maximum tumour yield was not decreased in Tg.AC mice exposed to chlorophene. In summary, there is no indication for reduced tumour latency due to chlorophene.

As mentioned in the Guidance to CLP, the latency of tumour formation does not materially affect the classification and hazard category. Any substance causing cancer will attract classification regardless of the latency for tumour development. This also includes tumour responses at late treatment/life periods if substance-related.

(e) whether responses are in single or both sexes; Tumours appeared in both sexes, male and female mice and in female rats. The renal tubular adenomas and carcinomas are restricted to male mice, the skin tumours appeared in female mice, whereas the TCCs were only seen in female rats. There are no data on possible differences in toxicokinetics between the sexes. Female mice did not develop adenomas despite almost a similar degree of nephropathy at the highest dose.

(f) whether responses are in a single species or several species; The renal tubular adenomas and carcinomas were observed in mice (oral, males), and the skin tumours were observed in mice (dermal, females), whereas the TCCs were observed in rats (oral, females). Thus, these findings are to be discussed as a several-species response.

It is mentioned in the Guidance to CLP that “a single study in one species and sex in combination with positive in-vivo mutagenicity data would be considered to provide sufficient evidence of carcinogenicity. Positive responses in several species add to the weight of evidence, that a chemical is a carcinogen.”

(g) structural similarity to a substance(s) for which there is good evidence of carcinogenicity; No comment added.

(h) routes of exposure; The oral exposure route chosen in the NTP studies is *per se* not relevant to the foreseeable human exposure pattern towards chlorophene. The dermal studies in CD-1 and transgenic mice cover the most relevant human exposure patterns, both test systems showing that chlorophene as a weak carcinogen.

(i) *comparison of absorption, distribution, metabolism and excretion between test animals and humans*; This comparison cannot be made due to a lack of relevant human data.

(j) *the possibility of a confounding effect of excessive toxicity at test doses*; See discussion above, under Mice, gavage, two-years NTP.

Despite evidence of nephropathy in mice and rats of both sexes, no increased incidences of renal neoplasms were observed in female mice or in male rats. This suggests that while nephrotoxicity may have been a necessary component, factors other than the marked nephrotoxicity of chlorophene were critical to the development of renal carcinogenesis in male mice.

(k) *mode of action and its relevance for humans, such as cytotoxicity with growth stimulation, mitogenesis, immunosuppression, mutagenicity*. The genotoxicity of chlorophene was concluded as negative, although some uncertainties with respect to assay insufficiencies existed. The mode of action underlying the development of renal tubular adenomas and carcinomas may involve regenerative hyperplasia. On the other hand as previously discussed studies in mice with both genotoxic and non-genotoxic chemicals, (bromochloromethane CAS number 74-27-5; nitrilotriacetic acid CAS number 139-13-9; tri(2,3-dibromopropyl)phosphate CAS number 126-72-7; 2,4-diaminophenol dihydrochloride CAS number 137-09-7) showed poor association between nephropathy and renal carcinogenicity. Thus, although nephropathy may be a permissive factor, other primary and secondary mechanisms are likely to be operative in the induction of the mouse renal neoplasms.

The aetiology of the TCCs is not clear. The low incidence may not be reflected in the occurrence of precursor lesions in the urothelium, even if there was transitional epithelium hyperplasia in the urinary bladder of one control animal and one high dose animal.

4.9.3 Conclusions on classification and labelling

In view of the induced renal neoplasms in male mice (2-year study) along with the two cases of renal transitional cell carcinoma, not observed in historical animals, in female rats (2-year study), and supported by the carcinogenic effect of chlorophene following dermal exposure of female CD-1 mice or female Tg:AC mice, chlorophene is considered a weak carcinogen. The genotoxic property of chlorophene was concluded as negative and no clear mode of action was established.

A classification for carcinogenicity is proposed:

According to Regulation (EC) No 1272/2008 (CLP): Carcinogen category 2, H351 suspected of causing cancer.

This is based on induction of two different types of tumours, in two species, in two sexes and following two routes:

Rat female (oral) + mice males (oral) + mice (dermal)

Mode of action: No clear mode of action was established

Threshold or non-threshold: The totality of the data available indicates no genotoxicity and threshold

4.9 Toxicity for reproduction

4.9.1 Effects on sexual function and fertility

Table 28: Effects of chlorophene on sexual function and fertility

Route of exposure	Test type, Method, Guideline	Species, Strain, Sex, no/group	Exposure Period	Doses	Critical effect	NOAEL/ LOAEL Parental	NOAEL/ LOAEL F1	NOAEL F2	Reference
Oral, gavage	One-generation study No guideline, but \cong OECD 415 Non-GLP	Rat Charles River albino strain 10 ♂+ 20 ♀ per group	♂: 60 days pre-mating, 106 days total ♀: 14 days pre-mating, through-out gestation and lactation	0, 50, 150 mg/kg bw/day	P males in the 150 mg/kg bw/day group gained less body weight during the pre-mating period than the control males. The body weight reduction continued until the end of the study. Females (P) were not affected. Both 4-day survival index and lactation index were reduced in litters at 150 mg/kg bw/day. At \geq 50 mg/kg bw/ day the body weight of male weanlings were significantly reduced on day 21 post partum.	Males: NOAEL = 50 mg/kg bw/day LOAEL = 150 mg/kg bw/day Females: not affected NOAEL \geq 150 mg/kg bw/day (highest dose)	LOAEL = 50 mg/kg bw/day (lowest dose)	n.a.	Confidential, 1973a* A_6_8_2(1)
Oral, gavage	Perinatal+ lactation study No guideline Non-GLP	Rat Albino 17-19 ♀ per group	From Day 15 of gestation through-out gestation and lactation	0, 50, 150 mg/kg bw/day	At \geq 50 mg/kg bw/day both the 12-day survival index and the 21-day lactation index were reduced.	NOAEL = 150 mg/kg/day (highest dose)	LOAEL = 50 mg/kg bw/day (lowest dose)	n.a.	Confidential, 1973b* A6_8_2(2)

* The studies were found to be of limited quality and not fulfilling the data requirement for biocides.

CLH REPORT FOR CHLOROPHENE

Route of exposure	Test type, Method, Guideline	Species, Strain, Sex, no/group	Exposure Period	Doses	Critical effect	NOAEL/ LOAEL Parental	NOAEL/ LOAEL F1	NOAEL F2	Reference
Oral, gavage	Two-generation study OECD 416	Rat Wistar (HsdCpb: WU convention ally bred) 30/ sex/group	♂: 10 weeks pre-mating, 2 wks mating period ♀: 10 weeks pre-mating, through-out gestation and lactation	0, 60, 180, 540 mg/kg bw/day	A significant and dose-related decrease in terminal body weight was observed in P males at ≥ 180 mg/kg bw/day and in F1 males at ≥ 60 mg/kg bw/day. Reduction in body weight gain during gestation was observed in dams at 540 mg/kg bw/day in the P generation and at 180 and 540 mg/kg bw/day in the F1 generation. Treatment-related kidney effects (nephropathy, dilated tubules, basophilic tubules and lymphocytic infiltration) were observed in P and F1 males at ≥ 60 mg/kg bw/day and P and F1 females at 540 mg/kg bw/day. A significantly lower female fertility index was observed at 540 mg/kg bw/day in the P generation and at 180 and 540 mg/kg bw/day in the F1 generation. A significantly longer oestrous cycle length and reduced fecundity was observed at 540 mg/kg bw/day in the F1 dams compared to control animals. (Repr Cat 2 H361f: Suspected of damaging fertility) Pups: Decreased terminal body weights at ≥ 180 mg/kg bw/day in the F1 litter and at 540 mg/kg bw/day in the F2 litter. Lower percentage of ear and eye opening at 540 mg/kg bw/day and incisor eruption at 180 mg/kg bw/day was found in both generations.	Males: LOAEL = 60 mg/kg bw/day (lowest dose) Females: NOAEL = 180 mg/kg bw/day LOAEL = 540 mg/kg bw/day	Parental F1: Males: LOAEL = 60 mg/kg bw/day (lowest dose) Females: NOAEL = 60 mg/kg bw/day LOAEL = 180 mg/kg bw/day Pups F1: NOAEL = 60 mg/kg bw/day LOAEL = 180 mg/kg bw/day	NOAEL = 60 mg/kg bw/day LOAEL = 180 mg/kg bw/day	Confidential, 2008 A6_8_2(3) KEY STUDY

There are submitted two studies examining **fertility and sexual function** and one follow up study on **lactation, all in rats**. The one-generation study and the perinatal and lactation study are both non-GLP and performed before guideline. Individual data are missing and the purity of the test compound is not given. The studies were found to be of limited quality and not fulfilling the data requirement for biocides (reproductive toxicity), hence a two-generation study was required in the completeness check phase of the evaluation. The two-generation study submitted by the applicant follows the OECD 416 guideline, are GLP and the purity and stability of the test compound are given.

The **one-generation** study was performed in albino **rats**. In this study male body weight gain was reduced in the high-dose group (150 mg/kg/bw/day) during the pre-mating period. The body weight reduction continued until the end of the study. Females of either dose group were not affected. There were no effects on the number of implantation sites, resorption sites and corpora lutea. Reproductive performance was not affected in any treatment groups. All delivered pups were normal in appearance. A dose dependent reduced 4-day survival and lactation index (79,1 and 61,1 % in the 50- and 150-mg/kg bw/day groups, compared with the control value of 80,4 mg/kg bw/day) was observed. Body weights of male weanlings in the 50- and 150-mg/kg bw/day groups were significantly lower than in control weanlings. Hence, the parental NOAEL in this study is set to 50 mg/kg bw/day for the males based on the reduced body weight. Since there were no effects observed in the dams the maternal NOAEL is ≥ 150 mg/kg bw/day The LOAEL for the F1 generation pups is 50 mg/kg bw/day based on the reduced body weight in the male weanlings. The maternal NOAEL in this study is higher than the NOAEL for the weanlings and it can not be excluded that the effect observed are treatment related. However, since there are shortcomings in the study design and report like; the reduced body weight in the male weanlings are not dose-dependent and individual data are not provided, the findings can not be used for classification of developmental toxicity.

A **two-generation reproductive toxicity study** (OECD 416) was recently conducted in **Wistar rats**. This oral gavage study (doses: 60, 180, 540 mg/kg/day) confirmed that the kidney is the target organ for general chlorophene toxicity in rats (please refer table 29 below for further details). Increases in absolute and relative kidney weights were observed in all treated males of both generations (please refer to table 29 for details). These findings were associated with macro- and microscopic renal lesions. A reduction in body weight gain during gestation was observed in dams at 540 mg/kg bw/day in the P generation and at 180 and 540 mg/kg bw/day in the F1 generation (please refer to table 4.13 for details). Body weights of the F1 pups were reduced at 180 and 540 mg/kg bw/day at lactation day 7, 14 and 21 (please refer to table 4.13 for details). The maternal body weight was unaffected at 60 and 180 mg/kg bw. No treatment-related changes in food intake were recorded in the dams. In the F2 pups body weights were reduced during lactation in the 540-mg/kg bw/day group. At the same dose level some effects on maternal bodyweight were recorded on lactation day 1. Lower percentage of ear and eye opening at 540 mg/kg bw/day and incisor eruption at 180 mg/kg bw/day was observed in both F1 and F2 pups.

In this study effects on fertility, fecundity and oestrus cycle length were reported. A significantly lower female fertility index was observed in the P generation at 540 mg/kg bw/day and at 180 and 540 mg/kg bw/day in the F1 generation, and a reduced fecundity was observed at 540 mg/kg bw/day in the F1 generation. A significant increase in oestrus cycle length (4.5 days) was observed in the F1 females after treatment with 540 mg/kg bw/day. In the study report this effect is suggested

to be incidental since the observed oestrus cycle length is within the historical control range (4.3-4.5). However, the oestrus cycle length of the concurrent control animals as well as the animals in the two lowest dose groups are lower than the historical control range (4.1, 4.0 and 4.1, respectively), and hence the effect should be compared to the experimental control group rather than the historical control range. Based on these findings chlorophene has an effect on the oestrus cycle length.

The overall NOAEL for adverse effect on development of the offspring is **60 mg/kg bw/day** based on the reduction in body weight in the F1 pups at 180 mg/kg bw/day. The NOAEL for adverse effects on sexual function and fertility is **60 mg/kg bw/day** based on the reduced female fertility index observed in the F1 generation. The **parental LOAEL** is **60 mg/kg bw/day** based on the kidney effects of male rats of both generations.

Table 29: Table for two-generation reproductive toxicity study in rats (confidential, 2008 / A6_8_2(3))

Parameter		Genera tion	Dose [mg/kg bw/day]								Dose respon se +/-	
			0		60		180		540		m	f
			m	f	m	f	m	f	m	f		
Food consumption		P	-	-	-	-	-	-	-	↑	-	+
		F1	-	-	-	-	-	-	-	↑	-	+
Body weight gain males: pre-mating females: gestation	% of control	P	100	100	96	101	91	94	72*	71*	+	+
		F1	100	100	94*	99	92*	86*	82*	81*	+	+
Organ weights												
<u>Kidney</u>	abs [g]	P	2.531	1.808	2.764*	1.817	2.735*	1.900	5.076*	2.401*	+	+
		F1	2.397	1.807	2.534*	1.888*	2.567*	1.832	3.020*	2.302*	+	+
	rel [% bw]	P	0.553	0.699	0.613*	0.706	0.640*	0.727	1.342*	0.911*	+	+
		F1	0.548	0.667	0.609*	0.693*	0.647*	0.688	0.839*	0.870*	+	+
<u>Adrenals</u>	rel [% bw]	P	0.013	0.033	0.014*	0.034	0.014*	0.034	0.018*	0.035	+	-
		F1	0.014	0.031	0.014	0.033*	0.016*	0.032	0.017*	0.033	+	-
Pathology												
<u>Kidney, surface</u> uneven or rough	Inci- dence	P	0	0	0	0	0	1	30	3	+	+
		F1	0	0	0	0	0	0	17	5	+	+
Histopathologic examination												
<u>Kidney,</u> lymphocyte infiltration	Inci- dence	P	0	0	0	2	2	1	0	8	-	+
		F1	2	0	0	0	0	0	0	2	-	-
<u>Kidney, dilated tubules</u> or nephropathy	Inci- dence	P	0	0	1	0	14	1	30	22	+	+
		F1	0	0	2	0	11	1	30	18	+	+
<u>Adrenals, hypertrophy</u> zona glomerulosa	Inci- dence	P	0	0	/	1	/	0	3	2	±	-
		F1	0	2	0	0	0	0	12	4	+	-
<u>Oestrus cycle length,</u> prior to co-habitation	[d]	P	-	4.5	-	4.2	-	4.5	-	4.5	./.	-
		F1	-	4.1	-	4.0	-	4.1	-	4.5*	./.	-
Reproductive Performance												
Fertility index	P	93.3	86.7	86.7	76.7*	+						
	F1	100.0	100.0	90.0*	83.3*	+						
Fecundity index	P	96.4	96.2	96.2	95.7	-						
	F1	100.0	100.0	96.3	96.0*	+						
Pup effects												
Pup weight PND1	[g]	F1	5.9	6.1	5.9	5.4*	+					
		F2	5.9	5.9	5.8	5.5*	+					
Pup weight PND21	[g]	F1	46.1	45.3	43.0*	39.1*	+					
		F2	44.3	44.5	43.6	40.5*						
Incisors erupted	% on PND11	F1	84.9	75.8	67.0*	45.5*	+					
		F2	89.5	87.0	72.4*	62.3*	+					
Ears open	% on PND14	F1	78.6	72.4	84.8	28.6*	+					
		F2	96.5	96.7	97.9	34.9*	+					

Parameter		Generation	Dose [mg/kg bw/day]								Dose response +/-	
			0		60		180		540		m	f
			m	f	m	f	m	f	m	f		
Eyes open	% on PND16	F1	100.0		98.4		100.0		91.6*		+	
		F2	99.6		98.6		99.0		97.7		-	

* Statistically significant difference to controls, p<0.05

4.9.1.1 Summary and discussion of effects on sexual function and fertility

In the one-generation study in rat, male body weight gains were reduced in the high-dose group during the pre-mating period. Females of either dose group were not affected. There were no effects on the number of implantation sites, resorption sites and corpora lutea. Reproductive performance was not affected in any treatment groups. All delivered pups were normal in appearance. Based on reduced body weight the **paternal NOAEL is 50 mg/kg bw/day**. Since there were not reported any maternal treatments related effects, **maternal NOAEL is 150 mg/kg bw/day** (highest dose). Based on reduced 4-day survival and 12-day survival and lactation index the **offspring LOAEL is 50 mg/kg bw/day** (lowest dose). Due to insufficiencies in the study design of both the one-generation and lactation study, the two-generation study is put as the key study for **fertility**. The **two-generation reproduction** oral gavage study in rats is recently performed (2008) and it confirmed that the kidneys are the target organ of chlorophene in rats. A reduction of body weight gain during gestation was observed in dams of both generations at 540 mg/kg bw/day and at 180 mg/kg bw/day. Pup weights of the F1 and F2 generation were reduced in the 180- and 540-mg/kg groups, respectively. A significantly lower female fertility index was observed in the P generation at 540 mg/kg bw/day and at 180 and 540 mg/kg bw/day in the F1 generation. A significantly increased oestrous cycle length and reduced fecundity was observed at 540 mg/kg bw/day in the F1 dams. It is likely that the reduced female fertility index is treatment-related as it was observed in both generations. Based on this the **NOAEL for adverse effects on sexual function and fertility** should be **60 mg/kg bw/day**. The overall **NOAEL for adverse effect on development of the offspring** is **60 mg/kg/day** based on the reduction of F1 pup weights in the two-generation study at 180 mg/kg bw/day. The **parental LOAEL is 60 mg/kg bw/day** based on the kidney effects of male rats of both generations.

4.9.1.2 Comparison with the CLP criteria

The effect on fertility, fecundity and oestrus cycle length were found at dose levels where some effects (reduced bw gain) of chlorophene were observed. In the Guidance on the application of the CLP Criteria (2009) it is stated in "3.7.2.2.1.1. *Effects to be considered in the presence of marked systemic effects*" that *adverse effect on fertility and reproductive performance seen only at dose levels causing marked systemic toxicity (e.g. lethality, dramatic reduction in absolute body weight, coma) are not relevant for classification purposes*". In this study the reduced maternal bw gain at the highest dose in P and F1 generation were lower than 10% compared to the control group. No lethality or coma related to treatment was observed at any dose levels. Further in the guidance it is stated "There is no established relationship between fertility effects and less marked systemic

toxicity. Therefore it should be assumed that effects on fertility seen at dose levels causing less marked systemic toxicity are not a secondary consequence of this toxicity.” Since the female fertility index was significantly reduced in both generations the effect is likely to be treatment-related. In addition, the fecundity and oestrous cycle was affected in the F1 dams of the highest dose-group. Based on the effects on fertility, fecundity and oestrus cycle length, chlorophene should be classified for fertility.

4.9.1.3 Conclusions on classification and labelling

Based on the reduced female fertility index observed in both generations and reduced fecundity and increased oestrous cycle length in the two-generation reproductive study, chlorophene should be classified for fertility effects. **A classification for fertility is proposed:**

Repr Cat 2 H361f: Suspected of damaging fertility according to Regulation (EC) No 1272/2008 (CLP).

4.9.1.4 Summary and discussion of effects on lactation

A dose dependent reduced 4-day survival and lactation index was observed in the one generation study performed in **albino rats**. A dose-dependent reduction in lactation index was observed, but no cross fostering was performed. No information on the purity and stability of the test material was provided.

A follow-up study (confidential, 1973b / A6_8_2(2)) was performed (table 30) to examine the effects of chlorophene on prenatal and postnatal performance of albino rats. Female rats were treated with 50 or 150 mg/kg bw/day from day 15 of gestation throughout the lactation period. Body weights of test and control females revealed no differences which could be attributed to the exposure to chlorophene. There were no deaths or abnormal behavioural reactions observed during the investigation. The number of pups delivered viable and retained through lactation day 4 were similar for all groups. Dose-dependent reduction in pups retained by females exposed to chlorophene compared to control females was observed on lactation days 12 and 21 (weaning) in the 50 and 150 mg/kg bw/day groups.

Table 30: Perinatal and lactation study in rats

Parameter	Controls	50 mg/kg	150 mg/kg	Dose-response
4-day survival index $\frac{\text{Number of Pups Viable at Lactation Day 4}}{\text{Number of Viable Pups Born}} \times 100$	98.6	96.7	97.9	–
12-day survival index $\frac{\text{Number of Pups Viable at Lactation Day 12}}{\text{Number of Pups Retained at Lactation Day 4}} \times 100$	97.9	88.9	86.9	–
Lactation index $\frac{\text{Number of Pups Viable at Lactation Day 21}}{\text{Number of Pups Retained on Lactation Day 4}} \times 100$	90.5	77.2	77.0	–

4.9.1.5 Comparison with the CLP criteria and conclusions on classification and labelling

Based on insufficiencies in the first study design (confidential, 1973a / A6_8_2(1)) these finding does not support classification for lactation.

It can not be excluded that the reductions in survival indexes a in the second study (confidential, 1973b / A6_8_2(2)) are treatment-related, but since no individual animal data was provided, no cross fostering test was performed and the effects observed were not dose dependent, no classification for lactation is recommended.

4.9.2 Developmental toxicity

Table 31: Teratogenicity of chlorophene

Route of exposure	Test type, Method, Guideline	Species, Strain, no/group	Exposure Period	Doses	Critical effects 1) dams 2) foetuses	NOAEL / LOAEL Maternal toxicity	NOAEL / LOAEL Teratogenicity, Embryo-toxicity,	Reference
Oral, gavage	Developmental toxicity No guideline, but \cong OECD 414	Rat CD (Sprague-Dawley derived) ♀ 20 /group	Day 6-15 of gestation	0, 100, 300, 900 mg/kg bw/day	1) At 300 mg/kg bw/day and 900 mg/kg bw/day animals showed significantly reduced body weight and food intake. 2) No treatment-related effects on litter size or foetal survival, weight and morphological development.	NOAEL = 100 mg/kg bw/day LOAEL = 300 mg/kg bw/day	NOAEL = 900 mg/kg bw/day (highest dose)	Confidential, 1985a A6_8_1(1)
Oral, gavage	Developmental toxicity Dose-range finding study Non-GLP	Rat CD (Sprague-Dawley derived) ♀, 6 /group	Day 6-15 of gestation	0, 100, 300, 1000 mg/kg bw/day	1) At \geq 100 mg/kg bw/day the overall weight gain was significantly reduced (day 6-15). One animal at 1000 mg/kg bw died after rapid weight and condition loss. 2) At 1000 mg/kg bw/day: foetal weight was reduced	LOAEL = 100 mg/kg bw/day (lowest dose)	NOAEL = 300 mg/kg bw/day LOAEL = 1000 mg/kg bw/day	Confidential, 1985b A6_8_1(4)

CLH REPORT FOR CHLOROPHENE

Route of exposure	Test type, Method, Guideline	Species, Strain, no/group	Exposure Period	Doses	Critical effects 1) dams 2) fetuses	NOAEL / LOAEL Maternal toxicity	NOAEL / LOAEL Teratogenicity, Embryo-toxicity,	Reference
Oral, gavage	Developmental Toxicity No guideline, but ≅ OECD 414	Rat Wistar (KFM-HAN) ♀ 25 /group	Day 6-15 of gestation	0, 15, 75, 375 mg/kg bw/day	1) At 375 mg/kg bw/day reduced body weight gain and food intake was observed. Three females of the 375 mg/kg bw/day group died during the treatment period. 2) The body weight was slightly but significantly reduced in the fetuses of the 375 mg/kg bw/day group and an increased incidence of non-ossified phalangeal nuclei was observed.	NOAEL = 75 mg/kg bw/day LOAEL = 375 mg/kg bw/day	NOAEL = 75 mg/kg bw/day LOAEL = 375 mg/kg bw/day	Confidential, 1984 A6_8_1(3) KEY STUDY
Oral, gavage	Developmental toxicity No guideline, but ≅ OECD 414	Rabbit NZW ♀ 14-16 /group	Day 6-19 of gestation	0, 10, 30, 100 mg/kg bw/ day	1) No effects on treated dams. 2) No adverse effects on fetuses.	NOAEL =100 mg/kg bw/day (highest dose)	NOAEL =100 mg/kg bw/day (highest dose)	Confidential, 1985c A6_8_1(2) KEY STUDY
Oral, gavage	Developmental toxicity Dose range finding study Non-GLP	Rabbit NZW ♀, 4/group	Day 6-19 of gestation	0, 30, 100, 150, 200, 300 mg/kg bw/ day	1) High mortality at 200 and 300 mg/kg bw. At 150 mg/kg/bw two animals was killed in extremis and abortions related to reduced bw was observed. 2) No compound related effects were observed at any dose-level.	NOAEL=100 mg/kg bw LOAEL=150 mg/kg bw	NOAEL = 150 mg/kg bw/day (highest dose)	Confidential, 1985d A6_8_1 (4)
Oral, gavage	Developmental toxicity No guideline, but ≅ OECD 414	Rabbit NZW ♀ 24 /group	Days 7-19 of gestation	0, 40, 80, 160 mg/kg bw/day	1) At 160 mg/kg bw/day body weights of dams were reduced during treatment, in addition a	NOAEL = 80 mg/kg bw/day (mortality, impaired bw gain)	NOAEL = 80 mg/kg bw/day. LOAEL = 160	Confidential, 1979 A6_8_1(5)

CLH REPORT FOR CHLOROPHENE

Route of exposure	Test type, Method, Guideline	Species, Strain, no/group	Exposure Period	Doses	Critical effects 1) dams 2) fetuses	NOAEL / LOAEL Maternal toxicity	NOAEL / LOAEL Teratogenicity, Embryo-toxicity,	Reference
	Non-GLP				<p>total of 10 of 24 (approximately 42 %) females died.</p> <p>2) None of the examined effects observed in the fetuses were considered to be treatment-related. Due to high maternal mortality in the highest dose group a proper embryotoxic/teratogenic evaluation could not be performed.</p>	LOAEL = 160 mg/kg bw/day.	mg/kg bw/day; no clear teratogenic or embryotoxic effects, but the maternal lethality was too high that one could conclude upon possible effect on the fetuses at this dose.	

Three oral **developmental toxicity** studies were performed in the **rat**. In these studies pregnant female rats were exposed to chlorophene by gavage from day 6 to 15 of gestation. In the key study (confidential, 1984 / A6_8_1(3)) reduced body weight gain and food intake was observed in the dams at 375 mg/kg bw/day. The body weight was reduced in the foetuses of the 375 mg/kg bw/day group and an increased incidence of non-ossified phalangeal nuclei was observed. This effect was probably secondary to the weight depression. The **NOAELs for maternal toxicity and developmental toxicity in rats** are both **75 mg/kg bw/day**. A limitation to these teratogenic studies in rats is that the dams have only been exposed to chlorophene during organogenesis and not from implantation and all the way through the gestation as required in the current version of OECD guideline 414 (2001).

The three **developmental toxicity** studies in **rabbits** did not reveal any clear adverse effects on foetal development. In the key study (confidential, 1985c / A6_8-1(2)) no adverse effect on dams or the foetuses were observed at highest dose level (100 mg/kg bw/day). In a dose range finding study (confidential, 1985d / A6_8_1 (4)) no treatments related effects were found at 100 mg/kg bw/day (NOAEL) in dams or at 150 mg/kg bw/day in foetuses (NOAEL). High mortality (2 of 4 animals died) of the dams was observed at 150 mg/kg bw/day, hence 100 mg/kg bw/day was used as highest dose level in the main study. Since the MTD seems to be between 100 and 150 mg/kg bw/day, a slightly higher top dose in the main study would have been preferable in the main study (confidential, 1985c / A6_8-1(2)).

In an older study (confidential, 1979 A6_8-1(5)) the mortality rate was significantly higher than control at the high-dose level (160 mg/kg bw/day) where 10 of 24 high-dose females died. According to OECD guideline 414, each test and control group should contain a sufficient number of females to result in approximately 20 female animals with implantation sites at necropsy. Groups with fewer than approximately 16 animals with implantation sites may be inappropriate. Maternal mortality does not necessarily invalidate the study providing it does not exceed 10 %. In the study the mortality was approximately 42 %. Therefore, the LOAEL of embryotoxic/teratogenic effects of 160 mg/kg bw/day could not be justified due to high maternal mortality. In addition, due to several shortcomings in the study design (uncertainties concerning purity and stability of the test substance, no information whether females inseminated by the same male were evenly distributed across the groups, maternal organ weights like the gravid uteri including the cervix were omitted and the heads of one-half of the foetuses examined should have been removed and processed for evaluation of soft tissue alterations in the brain, and this was also omitted) the assessment of maternal and offspring toxicity was incomplete. However based on the results given in the study a NOAEL of 80 mg/kg bw/day for maternal and embryotoxic/teratogenic effects was concluded..

Similarly to the studies on rats, the rabbits were only exposed to chlorophene during the period of organogenesis (days 6/7-19) and not all the way from implantation and through gestation. Based on the key study (confidential, 1985c / A6_8-1(2)) **NOAELs for maternal toxicity and developmental toxicity in rabbits** are both **100 mg/kg bw/day**.

4.9.2.1 Summary and discussion of developmental toxicity

Summary and discussion on developmental toxicity

The **developmental toxicity** studies in **rabbits** did not reveal any adverse effects on foetal development. Maternal and foetal body weight gain was the only affected parameter in the

developmental toxicity studies in rats. The **NOAEL rabbits for maternal effects is 100 mg/kg bw/day** and the **NOAEL rabbits for developmental toxicity is 100 mg/ kg bw/day (highest dose tested)**. The **NOAELs for maternal toxicity and developmental toxicity in rats are 75 mg/kg bw/day**.

4.9.2.2 Conclusions on classification and labelling

No classification is proposed.

4.10 Neurotoxicity

Not applicable

Chlorophene bears no structural similarity to organophosphates, carbamates or other known inducers of delayed neurotoxicity. Acute and repeated-dose studies in several species did not indicate the occurrence of neurotoxic effects and the rapid excretion of chlorophene precludes an accumulation of the compound.

4.11 Specific investigations: other studies

Supplementary studies on the active substance

A study on the induction of drug-metabolising enzymes by chlorophene was published by Kao *et al.* (1986). In this study, male F344 rats were treated orally with 500 mg/kg bw/day chlorophene (purity >99%) in corn oil on three consecutive days. Phenobarbital (PB), 3-methylcholanthrene (3-MC) and pregnelone-16 α -carbonitrile (PCN) were used as model substances. Livers and kidneys were harvested 24 h after the last treatment. Arylhydrocarbon hydroxylase (AHH), glutathione S-transferase (GST), N-demethylase, O-demethylase, and UDP-glucuronyl transferase (UGT) activities were determined. Cytochrome P450b (now known as CYP2B1) protein levels were determined by immunoblotting and radioimmunoassay. Other cytochrome P450 isozymes were separated and quantified using HPLC.

Treatment with chlorophene resulted in an increase in cytochrome P-450 content and an accompanying decrease in aryl hydrocarbon hydroxylase (AHH) activity in both liver and kidney microsomes. Several other drug-metabolizing enzymes were not affected by chlorophene treatment. However, in kidney, chlorophene induced NADPH-cytochrome-c reductase and UGT activities and caused a small increase in total cytochrome P-450 content and glutathione concentration.

The cytochrome P-450 isozymes induced by chlorophene were fractionated by HPLC. The HPLC profile following chlorophene treatment most closely resembled that seen after phenobarbital. Using an immunoblotting procedure and a radioimmunoassay, it was shown that the increase in cytochrome P-450 content in the liver after chlorophene treatment was, in part, due to an increase in the phenobarbital-inducible isozymes, CYP2B1 and 2B2.

In the kidney, the increase in total cytochrome P 450 content after chlorophene exposure was not due to an increase in CYP2B1 and 2B2. The decrease in AHH activity appeared to be caused by

non-competitive inhibition of constitutive AHH activity by chlorophene. Chlorophene also inhibited benzphetamine demethylation, although to a lesser extent. The failure to observe an increase in benzphetamine demethylase activity *in vivo*, despite the induction of CYP2B1, was probably due to the concomitant induction and inhibition of drug-metabolizing enzymes by chlorophene.

4.12 Human information

Medical surveillance of manufacturing plant personnel involved in chlorophene production revealed no health complaints associated with potential exposure to chlorophene (confidential, 2007 / A6_12_1).

A single report of contact dermatitis from chlorophene exposure is reported in the literature (Sonnex and Rycroft, 1986). A 49-year old bar manager developed contact dermatitis against chlorophene from a glass cleaning product.

5 ENVIRONMENTAL HAZARD ASSESSMENT

5.1 Degradation

Table 32: Summary of relevant information on degradation

Method	Results	Remarks	Reference Doc III
OECD 301B, ready biodegradability test (CO ₂ evolution)	Over 60 % degradation was observed during the test duration (28 days), but not within 10 days.	The two tests on ready biodegradability gave ambiguous results. The chlorophene concentration in the manometric respirometry test was high and inhibition might have taken place.	A7.1.1.2.1_01 (Bealing and Watson, 2002)
OECD 301 F, ready biodegradability test (manometric respirometry)	Chlorophene is considered not readily biodegradable under the conditions of this test.	The CO ₂ evolution test is considered more relevant as the chlorophene concentrations were below the EC ₅₀ value. This study indicates that chlorophene is readily biodegradable, since the pass level of 60 % degradation was reached. However, the pass level was not reached within the 10 day window as required by OECD 301B/CLP.	A7.1.1.2.1_02 (Reis, 2007a)
OECD 302B, inherent biodegradability (Zahn/Wellens test)	Inherently biodegradable	Despite the issues of adsorption (or dissipation) at the beginning of the test, the remaining DOC is eliminated to a large degree (92 %) and the overall conclusion "inherently biodegradable" is considered acceptable.	A7.1.1.2.2 (Reis, 2007c)
Test procedure taken from Loehr and Matthews, 1992	DT ₅₀ (dissipation): 21.4 d. A normalisation of the DT ₅₀ (23 °C) to 12 °C results in a DT ₅₀ (12 °C) of 51.6 d.	The study examines the primary biodegradation of chlorophene in a sandy silt loam soil under aerobic conditions and a temperature of 22-23 °C. There is no information about degradation products, bound residues, mineralisation and degradation pathways in the study.	A7.2.1 (Nitsche, 2011)
OECD proposal for a new test guideline 311, anaerobic biodegradation of chlorophene in anaerobically digesting sewage sludge.	No net carbon-production (as methane and carbon dioxide) was found. Chlorophene is considered not anaerobically biodegradable under the conditions of the test.	A study on the anaerobic degradation of chlorophene in pork liquid manure was also submitted as supporting information (Gerhardz, 2011). The study supports the result of the OECD 311 test.	A7.1.2.1.2 (Reis, 2007b)

5.1.1 Stability

5.1.1.1 Abiotic degradation

Hydrolysis

The hydrolysis of chlorophene in sterile aqueous buffered solutions at pH 4, 7 and 9 was studied according to test method EC C.7, degradation – abiotic degradation hydrolysis as a function of pH, Annex V, 92/69/EEC (Greenwood, 2003b). The concentrations of chlorophene were measured via HPLC-UV.

The preliminary test run at 50 °C indicated less than 10% hydrolysis after 5 days, therefore in accordance with the guideline no further testing is required, and chlorophene is considered hydrolytically stable at the tested pH values. The highest loss was observed at pH 7 (6.5% in 5 days). The correlation coefficients of the hydrolysis slopes at pH 4, 7 and 9 (concentration as a function of time) range from 0.0369 to 0.1159, therefore any further extrapolation is difficult. The estimated half-lives of chlorophene at pH 7 and 9 were 44.4 and 37.4 days, but as they are based on the slopes with weak correlation coefficients, they must only be viewed as indicative.

Results of the test are summarised in the following table.

Please see Document III-A7.1.1.1.1 for further details on this study.

Table 33: Hydrolysis of chlorophene

Guideline / Test method	pH	Temp. [°C]	Initial TS concentration, C ₀ [mg/L]	Reaction rate constant, K _h [d]	Half-life, DT ₅₀ [d]	Coefficient of correlation, r ²	Reference Doc III
EU method Annex V, C7 (92/69/EEC)	4, 7, 9	50	62.5 mg a.s./L	pH 4: no dissipation pH 7: 0.0156 day ⁻¹ pH 9: 0.0186 day ⁻¹	pH 4: no dissipation pH 7: 44.4 days pH 9: 37.4 days	pH 4: 0.0369 pH 7: 0.0862 pH 9: 0.1159	A7.1.1.1 (Greenwood, 2003b)

Photolysis in water


A study on photolysis of chlorophene in water was conducted according to the OECD proposal for a test guideline on phototransformation of chemicals in water – direct and indirect photolysis, draft August 2000 (Meinerling and Herrmann, 2007). The structure of the major photolysis product was elucidated in a follow-up study, by the combination of NMR, MS and UV/VIS spectroscopy (Freudenberger and Wesener, 2011), and the maximum concentration of this major photolysis product was estimated on the basis of the response of the test item in the UV chromatogram (Meinerling, 2011).

Chlorophene was incubated in aqueous buffer solution (pH 7) at a concentration of 5 mg/L over a time interval of 5 days (first experiment) and 4 hours (second experiment). The temperature during the study was maintained within the range of 20 to 30 °C. For technical reasons, the temperature during the first experiment was > 30 °C. Samples were either kept in the dark or exposed to a filtered xenon arc lamp. Residues of chlorophene were analysed by HPLC.

The chlorophene concentration rapidly dropped from an initial value of 99.4 % to 0 % of applied dose after 48 hours in the 5 days experiment and from an initial value of 99.7 % to a low value of 4.3 % of applied dose at the end of the 4 hours experiment. The aqueous photolysis half-life of chlorophene was calculated to be 0.7 hours.

The major photolysis product of chlorophene was identified as 2-hydroxy-xanthene (9H-xanthen-2-ol), and its maximum concentration was estimated to be 2.7 mg/L after 3 hours of irradiation. This corresponds to a relative concentration of 52.9 % of the parent substance. Furthermore, investigation by LC-MS/MS showed that two components with masses 184 and 212 were present in an additional sample containing a higher concentration of chlorophene (50 mg/L). However, as this sample was irradiated for analytical investigations only and since the high concentration of chlorophene is of no environmental relevance, a further in-depth investigation of these metabolites was not carried out.

Figure 2: The major photolysis product of chlorophene, 2-hydroxy-xanthene (9H-xanthen-2-ol)



Molecular Weight = 198.22
Molecular Formula = C₁₃H₁₀O₂

Under non-irradiated conditions, the mean parent concentration is not significantly reduced. Results are summarised in the following table.

Please see Document III-A7.1.1.1.2 for further details in this study.

Table 34: Photodecomposition of chlorophene in water

Guide-line / Test method	Initial TS concentration	pH value of the medium	Photolysis rate constant (k ^c _p)	Direct photolysis sunlight rate constant (k _{pE})	Reaction quantum yield (φ ^c _E)	Half-life (DT ₅₀)	Degradation products	Reference Doc III
OECD, 2000	5 mg a.s./L	7	0.98 [1/h]	11.97 [1/day]	Not available	0.7 h (experimental)	2-hydroxy-xanthene (9H-xanthen-2-ol), max. conc.: 52.9% of parent	A7.1.1.1.2 (Meinerling and Herrmann, 2007; Meinerling, 2011; Freudenberger and Wesener, 2011)

Photolysis in air and vaporisation behaviour

The half-life of chlorophene in air due to indirect photodegradation, i.e. oxidation with photochemically produced hydroxyl radicals, was calculated using the software programme AOPWIN, v. 1.91, 2000 by U.S.-EPA (Fàbregas, 2006). AOPWIN is a QSAR model which

requires only a chemical structure to make these predictions. The half-life of chlorophene in the troposphere was calculated to be 21.7 hours with a degradation rate of $18E^{-12} \text{ cm}^3 \times \text{molecule}^{-1} \times \text{s}^{-1}$. These estimations were carried out with respect to the OH radical reaction only, using a 24-hour-day with 5×10^5 OH radicals/cm³. According to these results ($DT_{50} < 2$ days), Chlorophene is rapidly degraded by photochemical processes and no accumulation of chlorophene in the air is to be expected.

The vaporisation of chlorophene from an inert surface (glass) was studied according to an internal test method (Nitsche, 2011). The test item was dissolved in isopropanol at a concentration of 1.043 mg/L, and 6.2 mL of this solution was put in petri dishes (three replicates). In addition, two petri dishes were filled with pure isopropanol and two were empty. All the dishes were stored on a desk surface in the working laboratory for 125 days, and were weighed at certain time intervals (15 times in total) throughout the test period. Temperature and humidity were also recorded.

After 125 days, the chlorophene quantity left on the glass surface was approximately 40 % of the amount originally applied. The vaporisation rate was linear ($0.020 \text{ g chlorophene} / \text{m}^2 \cdot \text{day}$). This study thus clearly indicates that chlorophene evaporates from an inert surface, though slowly.

Please see Document III-A7.3.1 and III-A7.3.2 for further details on this study.

5.1.2 Biodegradation

5.1.2.1 Biodegradation estimation

No data.

5.1.2.2 Screening tests

Ready biodegradability

Chlorophene was investigated for its ready biodegradability in two different tests, a CO₂ evolution test according to OECD test guideline 301 B (Bealing and Watson, 2002) and a manometric respirometry test according to OECD test guideline 301 F (Reis, 2007a).

The CO₂ evolution test was performed with an initial test substance concentration of 10 mg a.s./L and activated sludge with a suspended solids concentration of 30 mg/L. The pass level of 60 % ThCO₂ was reached within the test duration of 28 days. However, CO₂ production failed to satisfy the 10 day window requirement. After a lag phase of less than 4 days, there was a rapid degradation of chlorophene corresponding to a ThCO₂ evolution of just below 60 % (mean). Further degradation was inhibited for around 14 days, where after the ThCO₂ evolution reached 68 % (mean) of the theoretical maximum at the applied concentration over the course of the 28 day incubation. A similar behaviour was observed in the toxicity control, where the rate of CO₂ evolution initially kept pace with that observed in the reference control, but declined and stopped during a plateau phase between days 14 and 18 before restarting and continuing gradually until termination of the study. The inhibition occurred only after chlorophene was extensively degraded and may be caused by an unidentified degradation product of chlorophene. Alternatively, this could be explained by a second lag-phase required for the adaptation to degradation of a metabolite. In conclusion, over 60 % degradation of chlorophene was observed, but the rate of degradation did not

meet the criterion of 60 % degradation within 10 days from the onset of biodegradation (i.e. when 10 % degradation is reached) as required by OECD 301B/CLP.

Please see Document III-A7.1.1.2.1_01 for further details on the CO₂ evolution study.

In the manometric respirometry test the ready biodegradation of chlorophene was tested at an initial test substance concentration of 102 mg a.s./L and activated sludge with a suspended solids concentration of 30 mg/L. The degradation rate of chlorophene based on BOD was found to be 4 % (mean) after 14 days of incubation and 9 % (mean) after 28 days, at the end of the test. The degradation rate of chlorophene did thus not reach the pass level for ready biodegradability of 60 % based on BOD, and chlorophene is considered not readily biodegradable under the conditions of this test.

The level of test substance in the solution at the end of the test was high (81 % of the initial concentration), therefore adsorption to the activated sludge was not supposed to be of major influence. The possible degradation products 4-chlorophenol and 4-chlorocatechol were not found on day 14 sampling, but on day 28, minor amounts of 4-chlorophenol (about 20 µg/L) were identified. In the toxicity control, the degradation of chlorophene was similarly low, but degradation of the reference substance was not inhibited. The test substance concentration was above the EC₅₀ value for microorganisms (59.6 mg/L), therefore it is likely that the chlorophene inhibited the microorganisms in the activated sludge.

Please see Document III-A7.1.1.2.1_02 for details on the manometric respirometry study.

In conclusion, the two tests on ready biodegradability gave ambiguous results. The chlorophene concentration in the manometric respirometry test was high and inhibition might have taken place. Therefore, also assuming lower concentrations of chlorophene in STPs, the results of the CO₂ evolution test are considered more realistic. Here, 60 % degradation of chlorophene was observed during the 28 day test, but the level of 60 % degradation was not reached within the 10 day window.

Inherent biodegradability

The inherent biodegradability of chlorophene was tested in a Zahn/Wellens test according to OECD test guideline 302B (Reis, 2007c). The initial concentration of chlorophene was 87.4 mg a.s./L and the concentration of the activated sludge suspended solids was 200 mg/L. Based on DOC measurement, and after an initial rapid decline, the chlorophene level after 28 days was 7.8 % of the 3-hour concentration. This corresponds to a 92 % biodegradation. Chlorophene was thus found to be inherently biodegradable under the conditions of this test.

The DOC of the test flasks at 0 hours was 37-49 % of the initial concentration. Such a decline of chlorophene in the solution was also observed in the toxicity control. The recovery of chlorophene in the abiotic control without inoculum, however, was 100 %. This indicates that chlorophene adsorbed to the inoculum. Extraction of the activated sludge with acetonitrile resulted in very low recoveries; therefore it is likely that there is a strong, non-extractable binding of chlorophene (or primary degradation products) to the inoculum.

Despite these issues of adsorption (or dissipation) at the beginning of the test, the remaining DOC is eliminated to a large degree (92 %) and the overall conclusion "inherently biodegradable" is considered acceptable. The test item concentration was above the EC₅₀ for microorganisms, but the

high level of inoculum might have facilitated degradation, even if the inoculum might have been inhibited to some degree.

Please see Document III-A7.1.1.2.2 for further details on the inherent biodegradation study.

The results of the ready and inherent biodegradation studies are summarized in the table below.

Table 35: Ready and inherent biodegradation of chlorophene in laboratory tests

Guide-line/ Test method	Test type ¹	Test parameter	Inoculum			Add. substrate	Test substance conc.	Degradation		Reference Doc III
			Type	Concentration	Adaptation			Incubation period	Degree [%]	
OECD 301B	Ready	CO ₂	domestic sewage	0.03 g/L susp. solids	no	no	10 mg/L (nominal)	28 d	66-69%	A7.1.1.2.1_01 (Bealing and Watson, 2002)
OECD 301F	Ready	DOC or BOD	domestic sewage	0.03 g/L susp. solids	no	no	102 mg/L	28 d	9%	A7.1.1.2.1_02 (Reis, 2007a)
OECD 302B	Inherent	DOC and HPLC	industrial sewage	0.2 g/L susp. solids	Yes	no	87.4 mg/L	28 d	≥ 97%	A7.1.1.2.2 (Reis, 2007c)

¹ Test on *inherent* or *ready* biodegradability according to OECD criteria

An article from open literature (Swischer and Gledhill, 1973; Doc III A7.1.1.2.1_03, non-key study) is also available as supplementary information on the biodegradation of chlorophene. In this publication, the biodegradation of chlorophene (initial concentrations ranging from 0.1 to 20 mg/L) was studied in river water and three types of sludge systems over four weeks. A CO₂ evolution to the extent of approximately 60 % within the four week test period was observed. The study has shortcomings as several parameters were not reported. Nevertheless, it provides supporting information that biodegradation is expected under natural conditions.

5.1.2.3 Simulation tests

Aerobic degradation

There are no surface water simulation test data available.

The primary aerobic biodegradation of chlorophene in a single soil was studied according to a test procedure taken from Loehr and Matthews, 1992 (Nitsche, 2011).

The primary biodegradation of chlorophene in a sandy silt loam soil under aerobic conditions and a temperature of 22-23 °C was tested. The nominal concentration of the test substance was 10 mg/kg soil (analytically determined to be 8.6 mg/kg wet soil). The concentration of chlorophene and a reference substance (4-chloro-3-methylphenol) in the soil was monitored for 68 days using a substance specific HPLC analytical method. At the end of the test period, the test substance was dissipated for more than 85 % and a dissipation half-life of 21.4 days was derived. Normalisation of the DT₅₀ (23°C) to 12°C results in a DT₅₀ (12 °C) of 51.6 days.

There is no information about degradation products, bound residues, mineralisation and degradation pathways in this study. However, it shows the primary degradation of chlorophene and is considered acceptable for deriving a dissipation DT_{50} (12 °C) in soil of 51.6 days for the compound.

The results are summarized in the table below.

Please see Document III-A7.2.1 for further details on this study.

Table 36: Aerobic biodegradation of chlorophene in soil

Guideline / Test method	Test parameter	Test duration	Soil type	Test conditions	Initial test substance conc.	Results	Reference Doc III
Internal method	Concentration (HPLC measurements)	68 days	Sandy silt loam Total carbon: 1.1 % Humus: 1.9 % Clay: 12.9 % Sand: 53.9 % pH: 6.9 Moisture content: 80 % of field capacity	Temp: 22-23°C Light conditions: Darkness	10 mg/kg wwt (nominal) 8.6 mg/kg wwt (measured)	DT_{50} = 21.4 d (primary degradation/dissipation)	A7.2.1 (Nitsche, 2011)

Anaerobic biodegradation

The anaerobic biodegradation of chlorophene in anaerobically digesting sewage sludge was assessed according to the OECD proposal for a new test guideline 311 (Reis, 2007b).

A manometric test was conducted with a nominal chlorophene concentration of 140 mg/L over a period of 60 days at 30-37 °C in darkness. No net carbon-production (as methane and carbon dioxide) was found. The reference item sodium benzoate was found to be anaerobically biodegradable, but in the toxicity control containing both chlorophene and the reference substance, no biodegradation was observed. Chlorophene can thus be assumed to inhibit the digesting sludge inoculum. Chlorophene is considered not anaerobically biodegradable under the conditions of this test. The results are summarized in the table below. Please see Document III-A7.1.2.1.2 for further details on this study.

An anaerobic study with Preventol BP[®] (chlorophene) in pork liquid manure was also submitted as supporting information (Gerhardz, 2011; no study summary is provided, non-key study). Chlorophene was found to be degraded moderately during the first 20 days of the anaerobic test period, to a level of approximately 76 % of the initial concentration at day 20. Thereafter, the decrease of the chlorophene concentration was decelerated to an insignificant extent reaching a level of approximately 70 % of the initial concentration after 64 days. The study indicates that anaerobic biodegradation of chlorophene cannot be expected under conditions of the test system, and it thus supports the results of the study on anaerobically digesting sewage sludge.

Table 37: Anaerobic degradation of chlorophene in sewage sludge

Guideline /Test method	Test parameter	Inoculum			Test conditions	Initial test substance conc.	Degradation		Reference Doc III
		Type	Concentration	Adaptation			Incubation period	Degree [%]	
OECD 311 (proposal, anaerobic biodegradation)	CH ₄ and CO ₂ evolution	Anaerobically digesting sludge from a domestic STP in Darmstadt, Germany	2.1 g/L susp. solids	no	Temp: 30-37°C pH: 7.3-7.4 Light conditions: Darkness	140 mg/L (nominal)	60 d	0 %	A7.1.2.1.2 (Reis, 2007b)

5.1.3 Summary and discussion of degradation

5.1.3.1 Conclusion on abiotic degradation

It is not expected that hydrolytic processes will contribute to the degradation of chlorophene in the aquatic systems. Hydrolysis is not a major degradation pathway for chlorophene.

The aqueous photolysis half-life of chlorophene was calculated to be 0.7 hours. Therefore, photolysis will significantly contribute to the overall degradation of chlorophene in aquatic systems. The major degradation product is 2-hydroxy-xanthene (9H-xanthen-2-ol).

The calculated half-life of chlorophene in the troposphere is < 2 days (21.664 hours), and therefore no accumulation in the air is to be expected. Based on the vapour pressure and the Henry's Law constant (calculated, 3.7×10^{-3} Pa m³/mol), no significant volatilisation of chlorophene from water is to be expected. However, the study on vaporisation of chlorophene from an inert surface shows that evaporation does occur, but with a slow rate.

5.1.3.2 Conclusion on biodegradation

The two tests on ready biodegradability (Bealing and Watson, 2002 and Reis, 2007a) show dissimilar results. One test showed not ready biodegradability. This test was performed with high concentrations that are not considered environmentally relevant and which are above the EC₅₀ for microorganisms. The other test, which is considered more relevant, indicated that chlorophene is readily biodegradable, since the pass level of 60 % degradation was reached. However, the pass level was not reached within the 10 day window as required by OECD 301B/CLP. In this test, the chlorophene concentrations were below the EC₅₀ value.

Also in the inherent biodegradability test (Reis, 2007c), the chlorophene concentration was above the EC₅₀ for microorganisms. However, the inoculum concentration was very high, and this might have compensated for a partial inhibition. The results from this study indicate that > 50 % of the initially applied chlorophene was rapidly and strongly bound to the inoculum, and could not be extracted with acetonitrile. After the initial rapid decline, 92 % of the available DOC was biodegraded at the end of the test. Chlorophene is thus considered inherently biodegradable.

The study on aerobic biodegradation in soil (Nitsche, 2011) indicates a primary degradation/dissipation of more than 85 % of the initially applied chlorophene at the end of the test period of 68 days. A half-life in soil of 51.6 days (12 °C) was derived.

In the study on anaerobic biodegradation in sewage sludge (Reis, 2007b), no net carbon-production (as methane and carbon dioxide) was found. Chlorophene is therefore considered as not anaerobically biodegradable.

Based on the available data, the substance cannot be considered rapidly biodegradable.

5.2 Environmental distribution

5.2.1 Adsorption/Desorption

The adsorption/desorption behaviour of chlorophene in soil was investigated in two tests; a screening test according to OECD test guideline 121 (Jungheim, 2006b) and a test using four different soil types according to OECD test guideline 106 (Meinerling, 2007b).

In the screening study, the adsorption coefficient K_{oc} of chlorophene on soil was estimated using High Performance Liquid Chromatography (HPLC). Six reference standards of known K_{oc} were analysed on a HPLC system to determine an average capacity factor k' . Sodium nitrate was used to determine the HPLC system dead time (t_0). A regression curve with determined k' values and the known K_{oc} values ($\log k'$ versus $\log K_{oc}$) was plotted.

Chlorophene was analysed on the same HPLC system during the same sample sequence as the reference substances. The K_{oc} value for chlorophene was estimated by interpolation from the reference substance regression line. The estimated K_{oc} value for chlorophene is $\log 3.43$ ($K_{oc} = 2511$), further data is given in the table below. According to OECD, the K_{oc} estimated according to OECD test guideline 106 has a higher reliability. However, the output from an OECD 121 test is acceptable as an estimation of a chemical's partitioning behaviour.

Please see Document III-A7.1.3 for further details on the screening study.

Table 38: HPLC retention times and K_{oc} values for chlorophene and reference substances, OECD 121

Calibration substance	Retention time [tr] = min*	Capacity factor k' *	Log k' *	Log K_{oc}	Reference Doc III
2-Nitrobenzamide	5.535	0.678	- 0.169	1.45	A7.1.3 (Jungheim, 2006b)
3-Nitrobenzamide	5.680	0.722	- 0.141	1.95	
3,5-Dinitrobenzamide	6.018	0.824	- 0.084	2.31	
Triazoxide	7.389	1.240	0.093	2.44	
Naphthalene	8.276	1.509	0.179	2.75	
Phenanthrene	9.550	1.896	0.278	4.09	
Chlorophene (Test item)	8.532	1.587	0.201	3.43	

* mean value from 3 single values

In the OECD 106 test on four soil types (Meinerling, 2007b), the soils comprised of a sand, a loamy sand, a clay loam and a loam, respectively (corresponding to the soils Lufa 2.1, 2.2, 6S and Eurosoil 3). Aqueous solutions of the test item were equilibrated with the four soil types and the adsorption and desorption coefficients and constants were determined. The concentration of the test item was determined using HPLC methods.

According to the guideline, the OECD 106 study consists of a preliminary study, screening tests for adsorption and desorption, and the determination of Freundlich adsorption isotherms.

From results of the preliminary tests, a soil/solution ratio of 1:25 (2 g soil and 50 mL aqueous 0.01 M CaCl₂ solution of the test item) and a 48 h incubation time (to reach adsorption equilibrium) were chosen.

In the adsorption and desorption screening tests, desorption was estimated at one concentration of the test item using 4 different soil types. At desorption equilibrium, desorption of the adsorbed test item from the different soil types ranged from 18% up to 45% desorption. Adsorption was measured and the distribution coefficients K_d were in the range of 19 to 115 mL/g.

The Freundlich adsorption coefficients (K_F^{ads} values) varied between 31.8 and 145 μg^{1-1/n}(cm³)^{1/n}g⁻¹. This relates to average distribution coefficients between 19 (Lufa 2.1) and 115 mL/g (Eurosoil3) which correspond to the values determined in the screening test.

Freundlich coefficients determined for the desorption (K_F^{des} values) were higher than those determined for the adsorption (range of K_F^{des}: 66.0-190.3 μg^{1-1/n}(cm³)^{1/n}g⁻¹).

New Distribution coefficients, K_d, may be derived for the lowest test concentration for soil and water using the Freundlich equation, C_{soil} = K_d^F * C_{water}^{1/n} with K_d^F being the Freundlich distribution coefficient and 1/n the Freundlich exponent. Since c_{soil} is equal to K_d^F * c_{water}, the new K_d is K_d^F * C_{water}^{1/n-1}. From the new K_d, new K_{oc} values can be calculated by using the equation K_{oc} = K_d/f_{oc} with f_{oc} being the fraction organic carbon. The mean value of the four soil types for K_{oc} is used for PEC calculations as the tested concentrations are closest to the estimated environmental concentrations.

Table 39: Koc values at the lowest tested soil concentrations, OECD 106

Soils	Test conc. in water	Testconc. in soil	Kd	Koc
Eurosoil3	0.73	111	156	4726
Lufa 6S	1.44	93	64	3490
Lufa 2.2	1.51	103	75	3165
Lufa 2.1	2.51	63	25	2210
Mean value for Koc				3398

In conclusion, the resulting K_{oc} value derived from this study is 3398.

Please see Document III-A7.2.3.1 for further details on this study.

5.2.2 Volatilisation

Atmosphere

No atmospheric effect study is available as it is not required for biocides used as disinfectant (product types 2 and 3). Furthermore, based on the Henry's Law constant (calculated, $3.7 \times 10^{-3} \text{ Pa} \times \text{m}^3/\text{mol}$; Fàbregas, 2006), no significant volatilisation of chlorophene from water is to be expected. The vaporisation of chlorophene from an inert surface (glass) was studied according to an internal test method. Results show that evaporation of chlorophene occurs; however quite slowly. Calculations of the chemical lifetime in the troposphere resulted in a half-life of 21.7 hours. According to these results ($DT_{50} < 2$ days), chlorophene is rapidly degraded by photochemical processes and no accumulation of chlorophene in the air is to be expected.

Please see Document III-A7.3.1 for further details on the calculation.

5.2.3 Distribution modelling

No data.

5.3 Aquatic bioaccumulation

5.3.1 Aquatic bioaccumulation

Table 40: Summary of relevant information on aquatic bioaccumulation

Method	Results	Remarks	Reference Doc III
OECD 305, bioconcentration factor in fish (<i>Danio rerio</i>)	107-110 (whole body) 1401-1130 (lipid normalised)	The lipid content of the fish increased by more than 25 % from the start to the end of the study. A higher lipid content gives a lower BCF value (lipid normalised), but on the other hand, a higher lipid content may lead to a higher concentration (wet weight) at steady state and thus a higher BCF based on whole body wet weight. In conclusion, the measured steady-state whole body wet weight BCF values of around 100 L/kg are considered valid. 24 h after initiation of the depuration phase, no chlorophene was detected in any of the fish samples.	A4.3.3.1 (Confidential, 2009)

5.3.1.1 Bioaccumulation estimation

The aquatic BCF of chlorophene was estimated using the QSAR-approach as recommended in the Technical Guidance Document on Risk Assessment in support of Commission Directive 93/67/EEC on Risk Assessment for new notified substances; Commission Regulation (EC) No 1488/94 on Risk Assessment for existing substances; Directive 98/8/EC of the European Parliament and of the Council concerning the placing of biocidal products on the market (European Commission, 2003).

Based on a measured log Kow value of 4.275, the BCF-value of chlorophene was calculated to 858.5.

However, since a study on bioconcentration in fish is available (see below), this result is only included for information purposes.

Please see Document III-A7.4.2 (Fàbregas, 2007) for further details on the calculation.

5.3.1.2 Measured bioaccumulation data

The bioconcentration factor (BCF) of chlorophene in fish was determined in a study conducted according to OECD test guideline 305 (Confidential, 2009).

Groups of 37 and 47 individuals of *Danio rerio* were exposed to chlorophene for 8 days (uptake phase) at nominal concentrations of 3 and 15 µg/L, respectively. On day 9, the depuration phase was initiated, i.e. the fish were transferred to water with no chlorophene. The depuration phase lasted for 7 days. As pre-tests indicated that the bioconcentration of the test item might be fast and low, the study was carried out over only 15 days and with a reduced number of fish and water samples for analysis. The number of control fish taken for analysis was also reduced. Samples of the water and of fish were taken on days 1, 4, 6, 8, 9, 11 and 13, and the steady-state BCF was calculated as the ratio of the chlorophene concentrations in fish and water. In the fish samples during the uptake phase, the percentage of applied chlorophene in the treatment groups of 3 and 15 µg/L were 0.329-0.395 % and 1.433-2.046 %, respectively (HPLC determinations). 24 h after initiation of the depuration phase, no chlorophene was detected in any of the fish samples. For the treatment groups of 3 and 15 µg chlorophene/L, the steady-state BCF in terms of total bodyweight (wet weight) was 107 and 110 L/kg, respectively. In terms of lipid content, the steady-state BCF was 1401 and 1130 L/kg, respectively. Because of the fast uptake and elimination kinetics, no kinetic BCF could be determined.

The lipid content of the fish increased by more than 25 % from the start to the end of the study. A higher lipid content gives a lower BCF value (lipid normalised), but on the other hand, a higher lipid content may lead to a higher concentration (wet weight) at steady state and thus a higher BCF based on whole body wet weight. In conclusion, it is assumed that the increase in lipid content did not impair the test and that the measured steady-state whole body wet weight BCF values of around 100 L/kg are considered valid. The results are summarised in the table below. Please see Document III-A7.4.3.3.1 for further details on this study.

Table 41: Bioconcentration factors obtained for chlorophene

Test method	Test species	Exposure time	Log K _{ow}	Concentration of a.s. [µg/L]	BCF (L/kg)	Reference Doc III
Measured steady-state BCF according to OECD 305	<i>Danio rerio</i>	15 days	4.275	3	107	A7.4.3.3.1 (Confidential, 2009)
				15	110 (whole body)	
					1401 1130 (lipid normalised)	

5.3.2 Summary and discussion of aquatic bioaccumulation

The log K_{ow} value for chlorophene is 4.27. According to the TGD (2003), values greater than or equal to 3 indicate that the substance may bioaccumulate. However, the available fish bioconcentration study according to OECD 305 results in a measured steady-state BCF_{fish} based on whole body and lipid content of 110 L/kg and 1401 L/kg, respectively. However, no chlorophene could be detected in the fish samples 24 h after initiation of the depuration phase.

Based on this information, chlorophene is not expected to bioaccumulate in the environment.

5.4 Aquatic toxicity

Studies on the acute effects of chlorophene on fish and aquatic invertebrates are available. However, these are not considered valid due to insufficient reporting and the fact that only two test concentrations were used, from which no reliable LC_{50} or EC_{50} could be established. Please see sections 5.4.1.1 and 5.4.2.1 and Document III-A7.4.1.1 (non-key study) and III-A7.4.1.2 (non-key study) for further details.

However, chronic studies are available and the results of these tests are valid. The chronic toxicity towards fish was studied according to OECD 210. **This study resulted in a NOEC (mortality) of 0.58 µg a.i./L, the lowest NOEC among the chronic aquatic toxicity test indicators and the value used for the classification with H410: Very toxic to aquatic life with long lasting effects.**

The chronic toxicity towards daphnids was tested according to OECD 211 and a NOEC (reproduction) of 6.7 µg a.i./L was determined.

Growth inhibition of algae was tested according to OECD 201. The study resulted in an E_rC_{50} value of 197.2 µg a.i./L and a $NOErC$ value of 103.6 µg a.i./L. **The E_rC_{50} value of 197.2 µg a.i./L = 0.197 mg a.i./L gives a classification with H400: Very toxic to aquatic life.**

The inhibition of aquatic microbial activity was investigated according to OECD 209. In this study, an EC_{50} for aquatic microorganisms of 59.6 mg a.i./L was derived.

Table 42: Summary of relevant information on aquatic toxicity

Guideline / Test method	Results	Remarks	Reference Doc III
OECD 210, early life stage test with zebrafish (<i>Danio rerio</i>)	NOEC (mortality) = 0.58 µg/L = 0.00058 mg/L	Criteria for classification as H410, aquatic chronic 1, fulfilled by NOEC _{fish} and NOEC _{daphnia} . Criteria for classification as H400, aquatic acute 1, fulfilled by E _r C ₅₀ algae.	A7.4.3.2 (Confidential, 2008)
OECD 211, <i>Daphnia magna</i> reproduction test	NOEC (reproduction) = 6.7 µg/L = 0.0067 mg/L		A7.4.3.4 (Weyers, 2007)
OECD 201, growth inhibition of <i>Pseudokirchneriella subcapitata</i>	E _r C ₅₀ = 197.2 mg/L = 0.197 mg/L NOEC (growth) = 0.104 mg/L		A7.4.1.3 (Egeler et al., 2006)

5.4.1 Fish

5.4.1.1 Short-term toxicity to fish

A study on the acute toxicity of chlorophene towards fish (*Danio rerio*) in a static water system was submitted, but was not considered valid. The study was insufficiently reported, i.e. there was a lack of reporting of several test system details and test conditions (e.g. dissolved oxygen in the water and replicates). Furthermore, only two chlorophene concentrations were tested, 1.3 mg/L and 1.8 mg/L. No effects occurred in the lower treatment group, but at 1.8 mg/L, 100 % of the fish died. This considerable difference in effects at such close concentrations seems questionable. The derivation of a reliable LC₅₀ value would require a better and more thoroughly reported test system, and also more than two test concentrations. No statistical analyses could be carried out with only two test concentrations. There were also no measurements of the actual chlorophene concentrations. Please see Document III-A7.4.1.1 (non-key study) for further details.

However, a valid chronic toxicity study was submitted and the results from this tests is used for the purpose of the CLH proposal. Therefore, no new acute toxicity studies were requested.

5.4.1.2 Long-term toxicity to fish

The long-term toxicity of chlorophene has been tested in an early-life stage test with the Zebra-fish (*Danio rerio*) according to OECD test guideline 210 (Confidential, 2008).

Fertilised eggs were exposed under flow-through conditions to nominal concentrations of 0 (control), 0.95, 3.05, 9.77, 31.3 and 100 µg chlorophene/L. The test duration was 30 days after hatching of the larvae. The chlorophene concentrations were analytically determined prior to the initiation of the study, and at days 0, 6, 13, 20, 27, 34 and 37 for all test concentrations and the control. Mean measured concentrations ranged from 54 % to 81 % of the nominal concentrations, and therefore, analytically determined mean concentrations were used for the calculation and reporting of results. The daily recorded effects were mortality, hatching success, growth and symptoms of intoxication. The effect concentrations were statistically determined. The resulting lowest NOEC, based on mortality, was 0.58 µg test item/L (mean measured concentration). The LOEC was determined to be 1.65 µg test item/L (mean measured concentration).

Please see Document III-A.7.4.3.2 for further details on this study.

Table 43: Long-term toxicity of chlorophene to fish

Guideline/ Test method	Species	Endpoint/ Type of test	Exposure		Results, mean measured conc. [µg a.i./L]		Reference Doc III
			Design	Duration	NOEC	LOEC	
OECD 210 (1992)	<i>Danio rerio</i> (Zebra-fish)	Mortality, hatching success, symptoms of intoxication, growth parameters body weight and length of surviving fish	Flow- through	Until 30 days after hatching of larvae	Mortality: 0.58 Hatching: 73.0 Growth: 22.9 Criteria for classification as H410, aquatic chronic 1 fulfilled by NOEC (mortality).	1.65	A.7.4.3.2 (Confidential, 2008)

5.4.2 Aquatic invertebrates

5.4.2.1 Short-term toxicity to aquatic invertebrates

A study on the acute toxicity of chlorophene towards invertebrates (*Daphnia magna*) in a static water system was submitted, but as the acute fish study, it was not considered valid. This was mainly due to insufficient reporting (no information on holding and dilution water given, also no reported information on test conditions, e.g. pH, aeration of water, dissolved oxygen and temperature, and no information given on replicates) and the fact that only two concentrations were tested, 1.0 and 2.0 mg/L. The immobilisation in the two treatment groups were 0 % and 100 %, respectively, and these are not considered reliable results. The derivation of a reliable EC₅₀ value would require a better and adequately reported test system, as well as more than two test concentrations. No statistical analyses could be carried out with only two test concentrations. There were also no measurements of the actual chlorophene concentrations. Please see Document III-A7.4.1.2 (non-key study) for further details.

However, a valid chronic toxicity study was submitted and the results from this tests is used for the purpose of the CLH proposal. Therefore, no new acute toxicity studies were requested.

5.4.2.2 Long-term toxicity to aquatic invertebrates

A *Daphnia magna* reproduction study was conducted according to EEC Methods for Determination of Ecotoxicity Annex to Directive 97/548/EEC Part C, Method 20 'Daphnia magna Reproduction Test', corresponding to OECD test guideline 211 (Weyers, 2007).

The influence of chlorophene on the reproduction and survival rate of daphnids was investigated during 21 days in a semi-static system, with nominal chlorophene concentrations of 0.0032, 0.01, 0.032, 0.1 and 0.32 mg/L. The chlorophene concentrations were measured analytically on day 0, 7 and 14, and were in the range of 60-101 % and 30-95 % of the nominal values in the freshly

prepared and aged test, respectively. The results were therefore based on mean measured concentrations. The test parameters were survival of parent animals and number of offspring per adult.

A dose response relationship was observed for the number of offspring per adult, with no effects in the two lowest treatments. The NOEC and LOEC with regards to reproduction were determined to be 6.7 µg a.i./L 25.4 µg a.i./L. The NOEC and LOEC for mortality were ≥ 29.6 µg a.i./L.

Please see Document III-A7.4.3.4 for further details on this study.

Table 44: Reproduction test of chlorophene with *Daphnia magna*

Guideline / Test method	Endpoint/ Type of test	Exposure		Results, mean measured conc., [µg a.i./L]		Reference Doc III
		Design	Duration	NOEC/ LOEC		
EEC Method 20 (2001) (equal to OECD 211, 1998)	reproduction	Semi-static	21 d	6.7/ 25.4	reproduction	A7.4.3.4 (Weyers, 2007)
				≥ 29.6 / >29.6	mortality Criteria for classification as H410, aquatic chronic 1 fulfilled by NOEC (reproduction)	

5.4.3 Algae and aquatic plants

The influence of chlorophene on the growth of the green alga *Pseudokirchneriella subcapitata* (formerly *Selenastrum capricornutum*) was investigated in a 72 hours static test according to OECD test guideline 201 (Egeler et al. 2006).

P. subcapitata were exposed to nominal concentrations of 0.0048, 0.0153, 0.0488, 0.1563 and 0.5000 mg chlorophene/L. The chlorophene concentrations were measured analytically at 0 and 72 hours. The mean measured concentrations ranged from 82 % to 107 % of the nominal test concentrations at the beginning of the test and from 35 % to 39 % at the end. Therefore, all the results were based on mean measured concentrations. The calculated average growth rates decreased in a dose dependent manner. Effect concentrations were determined statistically. A clear dose-response relationship could be derived from the algae concentration data. The 72-hour E_rC_{50} value (growth rate inhibition) was 197.2 µg a.i./L and the 72 h NOE_rC value was 103.6 µg a.i./L. A deviation from the OECD 201 guideline is that the nutrient concentrations in the test medium were higher than recommended. However, there are no reasons to expect interactions between the test material and constituents of the test medium, and the test is therefore still regarded as reliable.

Please see Document III-A7.4.1.3_01 for further details on this study.

A second algae growth inhibition study is also available, which was not considered valid due to major deviations from the guideline and poor documentation of the study. A guideline which reportedly resembles the OECD 201 guideline was used. However, there were several deviations from the latter guideline. The information on test conditions and culture medium was very limited

or lacking, and not sufficient to assess the validity criteria of OECD 201. There were not enough test concentrations, and no monitoring of the chlorophene concentrations during the test. No information on replicates is given. The studied test parameter was cell number (algal biomass), and the effect concentration, E_bC_{50} , was determined graphically to be 0.038 mg/L, based on nominal concentrations. The initial cell concentration was not determined, the cell numbers were only counted at the end of the test (72 hours). No statistical analysis was carried out. Due to these deficiencies, only the first alga growth inhibition study (Egeler et al., 2006) is considered for CLH purposes and is presented in the table below.

Please see Document III-A7.4.1.3_02 for details.

Table 45: Toxicity of chlorophene to algae

Guideline / Test method	Species	Endpoint / Type of test	Exposure		Results, mean measured conc. [mg a.i./L]			Reference Doc III
			Design	Duration	NOEC	E_bC_{50} ¹	E_rC_{50} ²	
OECD 201 (2002)	<i>Pseudokirchneriella subcapitata</i>	Growth inhibition	Static	72 h	0.0323 (biomass) 0.1036 (growth)	0.1314	0.1972	A7.4.1.3 (Egeler et al., 2006)
					Criteria for classification as H400, aquatic acute 1 fulfilled by E_rC_{50}.			

¹ calculated from the area under the growth curve; ² calculated from growth rate

5.4.4 Other aquatic organisms (including sediment)

Inhibition of microbial activity (aquatic)

The effect of chlorophene on aquatic micro-organisms was investigated according to ISO Guideline 8192 (1986), which equals OECD test guideline 209 (Caspers and Müller, 1991).

Activated sludge was exposed to chlorophene concentrations of 10, 18, 32, 56 and 100 mg/L. The endpoint investigated was inhibition of the respiratory rate. Chlorophene showed inhibitory effects on the microbiological activity of activated sludge, and the EC_{50} was determined to be 59.6 mg a.i./L.

The results are based on nominal chlorophene concentrations, as there were no concentration measurements during the study. The EC_{50} value was not confirmed by statistical analysis and the results of the controls and reference substance are not reported. The test is still considered valid as inhibitory effects on microorganisms have been seen in biodegradation studies at concentrations in the same order of magnitude as the EC_{50} from this study.

Please see Document III-A7.4.1.4 for further details on this study.

Table 46: Influence of chlorophene on inhibition of microbial activity (aquatic)

Guideline/ Test method	Species/ Inoculum	Endpoint/ Type of test	Exposure		Results [mg a.i./L]			Reference Doc III
			Design	Duration	EC ₀	EC ₁₀	EC ₅₀	
ISO 8192 (1986) (equal to OECD 209)	Activated sludge	Oxygen consump- tion	Respira- tion inhibition	3 hours	< 8.22	20	59.6	III-A7.4.1.4 (Caspers and Mueller 1991)

Effects on sediment dwelling organisms

Testing on sediment-dwelling organisms is not a product-type specific requirement for PTs 2 and 3. The log K_{ow} for chlorophene is above the trigger value for a sediment assessment, and indirect emissions to water bodies can occur via effluent water from STPs and due to run-off from soil. An assessment using the equilibrium partitioning method was considered sufficient.

5.5 Comparison with criteria for environmental hazards (sections 5.1 – 5.4)

The criteria are taken from the Guidance on the Application of the CLP Criteria (ECHA, 2012).

- Rapid biodegradation:
 - The result from the manometric respiration test was: BOD 4 % (mean) after 14 days of incubation and 9 % (mean) after 28 days. This means that chlorophene does not meet the criteria. However, the concentration of chlorophene was above the EC₅₀ value for microorganisms (59.6 mg/L), therefore it is likely that the substance inhibited the microorganisms in the activated sludge.
 - The result from the CO₂ evolution test was: CO₂ reached 68 % of the theoretical maximum at the applied concentration over the course of the 28 day incubation. CO₂ production failed to reach the pass level within the ten-day window. Chlorophene did not degrade biotically in the aquatic environment by > 70 % in 28 days, and the substance can therefore **not** be classified as rapidly biodegradable.
 - Regarding abiotic degradation, hydrolysis is not an important degradation way for chlorophene. Photolysis can be an important way of primary degradation. However the test is not an ultimate biodegradation test and cannot be used for classification purposes.
 - Simulation tests: There are no surface water simulation test data available. The soil simulation test shows primary degradation, but not ultimate degradation.

The main conclusion is that the substance **cannot** be classified as rapidly biodegradable.

- Potential of bioaccumulation:
 - log K_{ow} > 4: Chlorophene meets the criterion.
 - BCF_{fish} based on whole body (110 L/kg) < 500: Chlorophene does not meet the criterion

Chlorophene has a high lipophilicity and has therefore a potential to bioconcentrate. However, experimentally derived BCF_{fish} (whole body) is preferred for classification purposes as such data overrides surrogate data such as a Kow value alone. In addition, in the fish BCF test no chlorophene could be detected in the fish samples 24 h after initiation of the depuration phase. Based on this, chlorophene does not meet the criterion for bioaccumulation.

- Aquatic toxicity:
 - Acute toxicity: The studies on acute toxicity towards fish and daphnids were not considered valid (please see sections 5.4.1.1 and 5.4.2.1 for more information), but there is a valid acute endpoint available for algae. The E_rC_{50} from the growth inhibition study on algae is < 1 mg/L (0.1972 mg/L) and gives the acute classification Category Acute 1.
 - Chronic toxicity: There are adequate chronic toxicity data available for all three trophic levels. The lowest chronic NOEC from the early life stage fish test is < 0.1 mg/L (0.00058 mg/l) and gives the chronic classification Category Chronic 1.

According to Annex 1: table 4.1.0 in the Guidance on the Application of the CLP Criteria, chlorophene meets the criterion for: (b) **Long-term aquatic hazard. (i) Non-rapidly degradable substances for which there are adequate chronic toxicity data available. Category Acute 1. Category Chronic 1.**

Substances with chronic toxicities below 0.1 mg/l (if non-rapidly degradable) contributes as components of a mixture to the toxicity of the mixture even at a low concentration and shall normally be given increased weight in applying the summation of classification approach. Therefore a multiplying factor (M-factor) has to be assigned. The M-factor can be taken from table 4.1.3 In Annex 1 to the Guidance on the Application of the CLP Criteria. The chronic M-factor for chlorophene is found in the table to be 100.

Regarding the acute toxicity classification, the acute M-factor is 1, according to table 4.1.3 in Annex I to the Guidance on the Application of the CLP criteria.

5.6 Conclusions on classification and labelling for environmental hazards (sections 5.1 – 5.4)

Symbol: GHS09

Signal word: WARNING

Aquatic Acute 1

H400: Very toxic to aquatic life

M-factor: 1

Aquatic Chronic 1

H410: Very toxic to aquatic life with long lasting effects

M-factor: 100

6 OTHER INFORMATION

7 REFERENCES

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
--	Eastin W. C. <i>et al.</i>	1998	The National Toxicology Program evaluation of genetically altered mice as predictive models for identifying carcinogens. <i>Toxicol Pathol.</i> 1998 Jul-Aug;26(4):461-73.	--	--	Yes	No
--	European Chemicals Agency (ECHA)	2012	Guidance on the Application of the CLP Criteria. <i>Version 3.0 November 2012</i>	--	--	Yes	No
--	European Commission, editor Joint Research Center, Institute for Health and Consumer Protection	2003	TGD for Risk Assessment (2003): Technical Guidance Document in Support of Commission Directive 93/67/EEC on Risk Assessment for New Notified Substances, Commission Regulation (EC) No 1488/94 on Risk Assessment for Existing Substances and Commission Directive 98/8/EEC concerning the Placing of Biocidal Products on the market.	--	No	Yes	No
--	Marsman D. S. <i>et al.</i>	1995	Chronic nephropathy and renal carcinogenicity of o-benzyl-p-chlorophenol in F344/N rats and B6C3F1 mice. <i>Fundam Appl Toxicol.</i> 1995 Sep;27(2):252-62.	National Institute of Environmental Health Services, Research Triangle Park, North Carolina, USA	--	Yes	No
--	Pritchard J. B. <i>et al.</i>	2003	The Role of Transgenic Mouse Models in Carcinogen Identification. <i>Environ Health Perspect</i> 2003 111(4): <i>doi:10.1289/ehp.5778</i>	--	--	Yes	No
A3.1(01) A3.10(01) A3.13(01)	Jungheim, R.	2007a	Physicochemical properties of chlorophene.	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen,	Yes	No	Yes

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
				Germany			
A3.2(01)	Olf, G.	2006	Vapor pressure, physical-chemical properties.	Bayer AG, BTS-PT-RPT-KPM, Leverkusen, Germany	Yes	No	Yes
A3.2(02)	Beiell, U.	2007	Calculation of Henry's Law Constant of Chlorophen (2-benzyl-4-chlorophenol).	Dr. Knoell Consult GmbH, Leverkusen, Germany	No	No	Yes
A3.3(01)	Kraus, H.	2006a	2-Benzyl-4-chlorophenol / Appearance.	LANXESS Deutschland GmbH, Leverkusen, Germany	No	No	Yes
A3.4(01)	Jungheim, R.	2007b	Spectral data of chlorophene.	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	Yes	No	Yes
A3.5(01)	Jungheim, R.	2006a	Determination of the water solubility (flask method) of chlorophene at 10 °C, 20 °C and 30 °C.	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	Yes	No	Yes
A3.5(02) A3.9(03)	Erstling, K.	2002	Water solubility, Preventol O extra in Schuppen.	Bayer AG, ZF-Zentrale Analytik, Leverkusen, Germany	Yes	No	Yes
A3.6(01), A3.9(01)	Greenwood, J.	2003a	BCP: Determination of the partition coefficient.	Covance LaboratoriesLtd., North Yorkshire, England	Yes	No	Yes
A3.7(01)	Jungheim, R.	2007c	Solubility of chlorophene in methanol and toluene at 10 °C, 20 °C and 30 °C.	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen,	Yes	No	Yes

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
				Germany			
A3.9(04)	Jungheim, R.	2004	Solubility of Preventol O extra in organic solvents.	Bayer Industry Services, Leverkusen, Germany	Yes	No	Yes
A3.9(02)	Feldhues, E.	2006	Statement Partition coefficient n-octanol / water of Preventol O extra, Temperature and pH dependence.	Bayer Industry Services, BIS-SUA-PUA I, Leverkusen, Germany	No	No	Yes
A3.11(01)	Heinz, U.	2007	Determination of safety-relevant data of Preventol BP.	Bayer Industry Services GmbH & Co. OHG, Safety / Environment / Analytics, Process and Plant Safety, Leverkusen, Germany	Yes	No	Yes
A3.17(01)	Kraus, H.	2006b	2-Benzyl-4-chlorophenol (chlorophene) / reactivity towards container material.	LANXESS Deutschland GmbH, Leverkusen, Germany	No	No	Yes
A4.1(01)	Erstling, K.	2007	Validation of a HPLC method for the determination of the relevant main and minor components in Preventol BP. CONFIDENTIAL	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	Yes	No	Yes
A4.2(02)	Meinerling, M.	2007a	Validation of an analytical method for the determination of Preventol BP (chlorophene) in water.	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany	Yes	No	Yes
A5.3.1(01)	Kugler, M.	2003	Determination of the antimicrobial effects of Preventol BP against bacteria and fungi.	Bayer Chemicals AG, Leverkusen, Germany	No	No	Yes
A5.3.1(02)	Bomblies, L. and	2000	Preventol BP (active substance). Determination	Labor L+S, Bad-	No	No	Yes

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
	Wedde, A.		of the "Minimal Inhibitory Concentration (MIC) against various test microorganisms.	Bocklet-Großenbrach, Germany			
A6_1_1	Confidential	1983a	Ortho-Benzyl Parachlorophenol, (Chlorophen): Acute Oral Toxicity in the Rat.	Confidential	No	No	Yes
A6_1_2	Confidential	1983b	Ortho-Benzyl Parachlorophenol, (Chlorophen): Acute Percutaneous Toxicity in the Rat.	Confidential	No	No	Yes
A6_1_3	Confidential	1983c	Ortho-Benzyl Parachlorophenol, (Chlorophen): Acute Inhalation Toxicity in the Rat.	Confidential	No	No	Yes
(A6_1_4)	Confidential	1983	Preventol BP - Examination of its Irritative Effects on Skin and Mucosa.	Confidential	No	No	Yes
(A6_1_4)	Confidential	1983d	Ortho-Benzyl Parachlorophenol, (Chlorophen): Acute Dermal Irritation/Corrosivity in Rabbits.	Confidential	No	No	Yes
A6.1.4(1)	Confidential	2000	Primary Dermal Irritation Study in Rabbits with Preventol BP (EPA/OECD/MAFF Guidelines).	Confidential	Yes	No	Yes
A6_1_4(2)	Confidential	1983e	Ortho-Benzyl Parachlorophenol (Chlorophen): Acute Eye Irritation/Corrosion Test in Rabbits.	Confidential	No	No	Yes
A6.1.5	Confidential	2001	Dermal Sensitization Study in Guinea Pigs – Closed Patch Test Technique with Preventol BP (EPA/OECD/MAFF Guidelines)	Confidential	Yes	No	Yes
(A6_1_5)	Confidential	1986	Preventol BP - Test for sensitizing effect on guinea pig skin ("Open Epicutaneous Test" according to Klecak)	Confidential	Yes	No	Yes
(A6_1_5)	Confidential	2002	Preventol BP Schuppen - Study for the skin sensitization effect in guinea pigs (Buehler Patch Test)	Confidential	Yes	No	Yes
A6_2_(1)	Kao, L. R. and Birnbaum, L. S.	1986	Disposition of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol in Male Rats	Systemic Toxicology Branch, NIEHS, Research Triangle Park, NC, USA	No	Yes	No

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
A6_2 (2)	Confidential	1994	Dermal Absorption of ¹⁴ C- <i>o</i> -Benzyl- <i>p</i> -Chlorophenol From a 5% Formulation.	Confidential	Yes	No	Yes
A6_2_6	Sonnex, T. S. and Rycroft R. J. G.	1986	Allergic Contact Dermatitis from Orthobenzyl P Chlorophenol in a Drinking Glass Cleaner	St, John's Hospital for Diseases of the Skin, London, England	No	Yes	No
A6_3_1	Confidential	1973a	21-Day Subacute Oral Toxicity Study with Santophen I in Beagle Dogs.	Confidential	No	No	No
A6_3_1	Sendelbach, L. E.	1982a	Repeated Oral Dose Study of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol in F344/N Rats.	Battelle, Columbus, OH, USA	No	Yes	No
A6_3_1	Sendelbach, L. E.	1982b	Repeated Oral Dose Study of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol in B6C3F ₁ Mice.	Battelle, Columbus, OH, USA	No	Yes	No
A6_3_2	Confidential	1984	Ortho-Benzyl Parachlorophenol (Chlorophen): Preliminary Dermal Toxicity Study in the Rabbit.	Confidential	Yes	No	Yes
(A6_3_2)	Confidential	1989	Ortho-Benzyl Parachlorophenol (Chlorophen): 21-Day Percutaneous Toxicity Study in the Rabbit.	Confidential	Yes	No	Yes
A6_3_2(1)	Confidential	1985	Ortho-Benzyl Parachlorophenol (Chlorophen): 21-Day Dermal Toxicity Study in the Rabbit.	Confidential	Yes	No	Yes
A6_3_2(2)	Confidential	1985	Preventol BP-Subacute toxicological study in rabbits (3-week trial with cutaneous application)	Confidential	Yes	No	Yes
A6_4_1(1)	Birnbaum, L. S. <i>et al.</i>	1986	Prechronic toxicity of <i>o</i> -benzyl- <i>p</i> -chlorophenol in rats and mice. <i>Fundam Appl Toxicol. 1986 Nov;7(4):615-25.</i>	--	--	Yes	No
A6_5+6_7 A6_7	National Toxicology Program	1994	NTP Technical Report on the Toxicology and Carcinogenesis Studies of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol (CAS No. 120-32-1) in F344/N Rats and B6C3F ₁ Mice (Gavage Studies).	National Toxicology Program, Research Triangle Park, NC, USA	No	Yes	No
A6_4_1(2)	Confidential	1973b	90-Day Subacute Oral Toxicity Study with	Confidential	No	No	No

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
			Santophen I in Beagle Dogs.				
A6_5	Confidential	2005	2-Benzyl-4-chlorphenol (Preventol BP) – Exploratory Subchronic Toxicity Study in Male Rats (16-Weeks Administration via Diet)	Confidential	No	No	Yes
A6_5+A6_7	Hejtmancik, M. <i>et al.</i>	1988a	The Chronic Gavage Study of o-Benzyl-p-Chlorophenol (CAS No. 120-32-1) in Fischer 344 Rats.	Battelle, Columbus, OH, USA	No	Yes	No
A6_6_1 (1)	Mortelmans, K. <i>et al.</i>	1986	<i>Salmonella</i> mutagenicity tests: II. Results from the testing of 270 chemicals.	EG&G Mason Research Institute & SRI International	No	Yes	No
A6_6_2	Confidential	1994	Chromosome Aberrations in Chinese Hamster Ovary (CHO) Cells.	Confidential	Yes	No	Yes
A6_6_3 (1)	Confidential	2005	BCP: Mutation at the <i>hprt</i> locus of L5178Y Mouse Lymphoma Cells using the Microtitre® Fluctuation Technique.	Confidential	Yes	No	Yes
A6_6_3 (2)	Caspary, W. J. <i>et al.</i>	1988	The mutagenic activity of selected compounds at the TK locus: rodent vs. human cells.	–	No	Yes	No
A6_6_4 (1)	Confidential	1990	Nipacide BCP: Assessment of Clastogenic Action on Bone Marrow Erythrocytes in the Micronucleus Test.	Confidential	Yes	No	Yes
A6_6_4 (2)	Confidential	1972	Mutagenic Study with Santophen I in Albino Mice.	Confidential	No	No	No
A6_6_5	Confidential	2009	Single Cell Gel Electrophoreses (Comet) Assay in the Male Mouse: In Vivo.	Confidential	Yes	No	Yes
A6_7 (1)	Hejtmancik, M. <i>et al.</i>	1988b	The Chronic Gavage Study of o-Benzyl-p-Chlorophenol (CAS No. 120-32-1) in B6C3F ₁ Mice.	Battelle, Columbus, OH, USA	No	Yes	No
A6_7_ (2)	Spalding, J. W. <i>et al.</i>	1999	Development of a transgenic mouse model for carcinogenesis bioassays: evaluation of	Research Triangle Park, North Carolina,	No	Yes	No

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
			chemically induced skin tumors in Tg.AC mice	US			
A6_7 (3)	National Toxicology Program	1995	One-year initiation/promotion study of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol (CAS No. 120-32-1) in Swiss (CD-1 [®]) Mice (Mouse Skin Study)	National Toxicology Program, Research Triangle Park, NC, USA	Yes	Yes	No
A6_8_1	Confidential	1984	Embryotoxicity (Including Teratogenicity) Study with Preventol BP Technical in the Rat.	Confidential	Yes	No	Yes
A6_8_1	Confidential	1979	A Segment II Teratology Study with Santophen I in Rabbits	Confidential	No	No	Yes
A6_8_1_(1)	Confidential	1985a	Chlorophen: Teratology Study in the Rat.	Confidential	Yes	No	Yes
A6_8_1_(2)	Confidential	1985c	Chlorophen: Effects of Oral Administration upon Pregnancy in the Rabbit.	Confidential	Yes	No	Yes
A6_8_1_(4)	Confidential	1985b	Chlorophen: Effects of Oral Administration upon Pregnancy in the Rat. 1. Dosage Range-Finding Study.	Confidential	No	No	Yes
A6_8_1_(4)	Confidential	1985d	Chlorophen: Effects of Oral Administration upon Pregnancy in the Rabbit. Dosage range-finding study.	Confidential	No	No	Yes
A6.8.2(1)	Confidential	1973a	Reproduction Study with Santophen I in Albino Rats.	Confidential	No	No	Yes
A6.8.2(2)	Confidential	1973b	Perinatal and Lactation Study with Santophen I in Albino Rats.	Confidential	No	No	Yes
A_6_8_2_(3)	Confidential	2008	Chlorophene: Two Generation Reproduction Toxicity Study by Gavage in Wistar Rats.	Confidential	Yes	No	Yes
(A6_10)	Kao, L. R., <i>et al.</i>	1986	Effect of <i>o</i> -Benzyl- <i>p</i> -Chlorophenol on Drug-Metabolizing Enzymes in Rats	Systemic Toxicology Branch, NIEHS, Research Triangle	No	Yes	No

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
				Park, NC, USA			
A6_12_1	Confidential	2007	Medical statement – 2-benzyl-4-chlorophenol (BP)	Confidential	No	No	Yes
A7.1.1.1.1(01)	Greenwood, J.	2003b	BCP: Evaluation of hydrolysis as a function of pH (HPLC screen).	Covance LaboratoriesLtd., North Yorkshire, England	Yes	No	Yes
A7.1.1.1.2	Meinerling, M. and Herrmann, S.	2007	Phototransformation of Preventol BP (Chlorophene) in Water.	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany	Yes	No	Yes
A7.1.1.1.2	Meinerling, M.	2011	Non-GLP Statement on IBACON Project 33341176, Photolytic degradation of Preventol BP	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany	--	No	Yes
A7.1.1.1.2	Freudenberg, Ch. and Wesener, J. R.	2011	Structure elucidation of the major photolysis product of Preventol BP (chlorophene)	Currenta GmbH & Co. OHG, Leverkusen, Germany	Yes	No	Yes
A7.1.1.2.1(01)	Bealing, D. J. and Watson, S.	2002	BCP: Assessment of ready biodegradability by measurement of carbon dioxide evolution.	Covance Laboratories Ltd., Harrogate, England.	Yes	No	Yes
A7.1.1.2.1(02)	Reis, K. H.	2007a	Ready biodegradability of chlorophene in a manometric test	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany.	Yes	No	Yes
A7.1.1.2.1(03)	Swisher, R. D. and Gledhill, W. E.	1973	Microbial degradation of O-Benzyl-p-Chloro-phenol <i>CSMA – Proceedings of the 60th Annual Meeting</i>	Published by Chemical Specialities Manufacturers	--	Yes	No

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
				Association Inc.			
A7.1.1.2.2(01)	Reis, K. H.	2007c	Inherent Biodegradability of Chlorophene in a Zahn-Wellens/EMPA Test.	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany.	Yes	No	Yes
A7.1.2.1.2	Reis, K. H.	2007b	Anaerobic biodegradability of Chlorophene in digested sludge: Measurement of gas production.	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany.	Yes	No	Yes
A7.1.2.1.2	Gerhardz, T.	2011	Biodegradation of 5 mg/kg Preventol BP® (2-benzyl-4-chlorophenol) in pork liquid manure under anaerobic conditions.	Lanxess Deutschland GmbH, Leverkusen.	No	No	Yes
A7.1.3(01)	Jungheim, R.	2006b	Determination of the Adsorption Coefficient (K_{oc}) by High Performance Liquid Chromatography (HPLC) Method of Chlorophene.	Bayer Industry Services GmbH & Co. OHG, BIS-SUA-Analytics, Leverkusen, Germany	Yes	No	Yes
A7.2.1	Nitsche, M.	2011	Biodegradation of Preventol® BP (Chlorophen) in soil under aerobic conditions	LANXESS Deutschland GmbH, Leverkusen, Germany	No	No	Yes
A7.2.3.1(01)	Meinerling, M.	2007b	Determination of the Adsorption / Desorption Behaviour of 2-Benzyl-4-chlorophenol (Preventol BP).	Institut für Biologische Analytik und Consulting IBACON GmbH, Rossdorf, Germany	Yes	No	Yes
A7.3.1(01)	Fàbregas, E.	2006	Calculation of indirect photodegradation of Chlorophene.	DR. KNOELL CONSULT GmbH, Mannheim, Germany	No	No	Yes
A7.4.1.1(01)	Confidential	1986	Preventol BP (2-benzyl-4-chlorophenol): Fish	Confidential	No	No	Yes

CLH REPORT FOR CHLOROPHENE

Section No. in in CA report on chlorophene	Author(s)	Year	Title	Testing Company	GLP Study (Yes/No)	Published (Yes/No)	Data Protection Claimed (Yes/No)
			Toxicity, <i>Brachydanio rerio</i> .				
A7.4.1.2(01)	Caspers, N.	1986a	Preventol BP (2-benzyl-4-chlorophenol): Toxicity, <i>Daphnia magna</i> .	Bayer AG, WV Umweltschutz, Leverkusen, Germany	No	No	Yes
A7.4.1.3(02)	Caspers, N.	1986b	Preventol BP (2-benzyl-4-chlorophenol): Growth inhibition test Algae.	Bayer AG, WV Umweltschutz, Leverkusen, Germany	No	No	Yes
A7.4.1.3	Egeler, Ph. <i>et al.</i>	2006	Preventol BP: A study on the toxicity to algae (<i>Pseudokirchnerella subcapitata</i>)	ECT Ökotoxikologie GmbH, Flörsheim, Germany and Batelle, Geneva, CH	Yes	No	Yes
A7.4.1.4	Caspers, N. and Müller, G.	1991	Untersuchungen zur Bakterientoxizität von Preventol BP Schuppen.	Bayer AG, Institut für Umweltanalyse und Bewertungen, Leverkusen, Germany	Yes	No	Yes
A7.4.2(01)	Fàbregas, E.	2007	Calculation of the Bioconcentration Factor (BCF) of Chlorophene.	DR. KNOELL CONSULT GmbH, Mannheim, Germany	No	No	Yes
A7.4.3.2(01)	Confidential	2008	Toxicity of 2-Benzyl-4-chlorophenol (Preventol BP) to Zebra-fish (<i>Danio rerio</i>) in an Early-Life Stage Test	Confidential	Yes	No	Yes
A7.4.3.3.1	Confidential	2009	Bioconcentration: Flow-through Fish Test with Chlorophene (Preventol BP)	Confidential	Yes	No	Yes
A7.4.3.4(01)	Weyers, A.	2007	Chlorophene, <i>Daphnia magna</i> Reproduction Test.	Bayer Industry Services GmbH & Co: OHG, Leverkusen Germany	Yes	No	Yes

8 ANNEXES (EXCERPTS OF DOCUMENT III-A TO THE ACTIVE SUBSTANCE DOSSIER)

Annexed to the CLH report are study summaries of relevant health and environmental studies in the form of excerpts of Document III-A from the active substance dossier submitted under the Biocidal Products Directive (98/8/EC). The annexes are confidential.

List of annexes

- | | |
|---------|--|
| Annex 1 | Extract from Competent Authority Report – Directive 98/8/EC on the placing of biocidal products on the market. Document III-A (A6.1-A6.12), study summaries – active substance, section A6: human health |
| Annex 2 | Extract from Competent Authority Report – Directive 98/8/EC on the placing of biocidal products on the market. Document III-A (A7.1-A7.4), study summaries – active substance, section A7: environment |