

Summary of product characteristics for a biocidal product

Product name: Magnum Gel Cucarachas IGR PLUS

Product type(s): PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Authorisation number: ES/APP(NA)-2021-18-00746

R4BP 3 asset reference number: ES-0015195-0000

Table Of Contents

Administrative information	1
1.1. Trade names of the product	1
1.2. Authorisation holder	1
1.3. Manufacturer(s) of the biocidal products	1
1.4. Manufacturer(s) of the active substance(s)	2
2. Product composition and formulation	2
2.1. Qualitative and quantitative information on the composition of the biocidal product	2
2.2. Type of formulation	2
3. Hazard and precautionary statements	3
4. Authorised use(s)	3
5. General directions for use	14
5.1. Instructions for use	14
5.2. Risk mitigation measures	14
5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment	14
5.4. Instructions for safe disposal of the product and its packaging	15
5.5. Conditions of storage and shelf-life of the product under normal conditions of storage	15
6. Other information	15

Administrative information

1.1. Trade names of the product

Magnum Gel Cucarachas IGR PLUS
MAGNUM GEL CUCARACHAS IGR
ATOMIX PRO GEL
ULTIMATUM ROACHES IGR

1.2. Authorisation holder

Name and address of the authorisation holder	Name	MYLVA S.A.
	Address	Via Augusta, 48 08006 Barcelona Spain
Authorisation number	ES/APP(NA)-2021-18-00746	
R4BP 3 asset reference number	ES-0015195-0000	
Date of the authorisation	23/03/2021	
Expiry date of the authorisation	23/03/2026	

1.3. Manufacturer(s) of the biocidal products

Name of the manufacturer	Mylva S.A.
Address of the manufacturer	Via Augusta 48 08006 Barcelona (Catalonia) Spain
Location of manufacturing sites	Sant Galderic 23, poligon industrial ponent , Sant pol de Mar 08395 Barcelona (Catalonia) Spain

1.4. Manufacturer(s) of the active substance(s)

Active substance	37 - imidacloprid
Name of the manufacturer	ADAMA Agriculture España S.A.
Address of the manufacturer	C/ Príncipe de Vergara 110, planta 5 28002 Madrid Spain
Location of manufacturing sites	Adama Makhteshim Ltd. - Neot-Hovav Eco-Industrial Park 84100 Beer Sheva Israel Jiangsu Yangnong Chemicals Group Co. Ltd - 39 Wenfeng Road 225009 Yangzhou China
Active substance	1386 - S-Methoprene
Name of the manufacturer	Bábolna Bioenvironmental Centre Ltd.
Address of the manufacturer	H-1107 Budapest, Szállás u. 6. 1107 Szállás, Budapest Hungary
Location of manufacturing sites	H-1107 Budapest, Szállás u. 6. 1107 Szállás, Budapest Hungary

2. Product composition and formulation

2.1. Qualitative and quantitative information on the composition of the biocidal product

Common name	IUPAC name	Function	CAS number	EC number	Content (%)
imidacloprid	(2E)-1-[(6-chloropyridin-3-yl) methyl]-N-nitroimidazolidin-2-imine	Active Substance	138261-41-3	428-040-8	2,15
S-Methoprene	Isopropyl-(2E,4E,7S)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate	Active Substance	65733-16-6		0,5
OIT		Non-active substance	26530-20-1	247-761-7	0,04

2.2. Type of formulation

RB - Bait (ready for use)

3. Hazard and precautionary statements

Hazard statements

Very toxic to aquatic life with long lasting effects.

Contains 2-octyl-2H-isothiazol-3-one (OIT) and 1,2-benzisothiazol-3-(2H)-one (BIT).
May produce an allergic reaction.

Precautionary statements

Keep out of reach of children.

Read carefully and follow all instructions.

Avoid release to the environment.

Collect spillage.

Dispose of contents to container as hazardous waste to a registered establishment or undertaking, in accordance with current regulations (Trained Professionals).

Dispose of contents to or its container as hazardous waste according to the regulations in force. (Professionals and Non-professionals).

4. Authorised use(s)

4.1 Use description

Use 1 - Use # 1 – Indoors, cracks and crevices - Gel bait applied as drops - General public (non-professional users).

Product type

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Where relevant, an exact description of the authorised use

Insecticide and growth regulator against cockroaches.

Target organism(s) (including development stage)

Scientific name: *Periplaneta americana*
Common name: American cockroaches
Development stage: Adults and nymphs

Scientific name: *Blatta orientalis*
Common name: Oriental cockroaches
Development stage: Adults and nymphs

Scientific name: *Blattella germanica*
Common name: German cockroaches
Development stage: Adults and nymphs

Field(s) of use

Indoor

Indoors, crack and crevices.
Private houses.

Application method(s)	Method: Bait application Detailed description: Open application of a gel bait applied as drops from a syringe.
Application rate(s) and frequencies	Application Rate: 1 aplicación cada 2 semanas. Volver a aplicar una vez más si la infestación persiste. Dilution (%): 0 Number and timing of application: Dose: depends on the level of infestations and species of cockroaches (1 drop = 0.04 g). German cockroaches (<i>Blatella germanica</i>): 0.12-0.16 g/m ² (3-4 drops/m ²). Oriental cockroaches (<i>Blatta orientalis</i>): 0.16-0.24 g/m ² (4-6 drops/m ²) American cockroaches (<i>Periplaneta americana</i>): 0.16 g/m ² -0.24 g/m ² (4-6 drops/m ²) Frequency of application: 1 application in 2 weeks. Reapply only once more if the infestation persists. Frequency of treatment: Three months after the infestation's end, treatment may be repeated. Maximum of 6 applications per year.
Category(ies) of users	General public (non-professional)
Pack sizes and packaging material	LDPE plastic syringes of 1, 2, 3, 4, 5, 8, 10, 15 grams of gel bait.

4.1.1 Use-specific instructions for use

Ready-to-use gel bait to be applied as drops deposited with the syringe. Drops adhere to non-porous surfaces.
Apply only in crack and crevices. .
Use indoors in private houses.
Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel.
Apply the biocidal product only as spot application in crack and crevices where cockroaches may be present out of the reach of children and animals:
- In inaccessible places.
- In dark places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: cracks and crevices behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.
Do not use on wood or porous surfaces.
Avoid contact with treated surfaces.
Do not expose bait drops to sunlight or heat source (i.e. radiator).
Do not mix with other chemicals or in areas recently treated with another insecticide.

4.1.2 Use-specific risk mitigation measures

Avoid contact with treated spots.
Avoid contact with eyes and skin.
The product should not be applied in a zone accessible to children
The treatment must be restricted to areas out reach of animals.

4.1.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.1.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product and other waste generated during the treatment are considered hazardous waste. Dispose of in accordance with current regulations
Do not release to soil, ground, surface water or any kind of sewer.

4.1.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

4.2 Use description

Use 2 - Use # 2 – Indoors, cracks and crevices - Gel bait applied as drops - Professional users.

Product type	PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)
Where relevant, an exact description of the authorised use	Insecticide and growth regulator against cockroaches.
Target organism(s) (including development stage)	Scientific name: <i>Periplaneta americana</i> Common name: Cucaracha americana Development stage: adults, nymphs Scientific name: <i>Blatta orientalis</i> Common name: Cucaracha oriental Development stage: adults, nymphs Scientific name: <i>Blattella germanica</i> Common name: Cucaracha alemana Development stage: adults, nymphs
Field(s) of use	Indoor Indoors, crack and crevices. Houses and commercial buildings.
Application method(s)	Method: Bait application Detailed description: Open application of a gel bait applied as drops from a syringe.
Application rate(s) and frequencies	Application Rate: 1 application in 2 weeks. Reapply only once more if the infestation persists. Dilution (%): 0 Number and timing of application: Dose: depends on the level of infestations and species of cockroaches (1 drop = 0.04 g). German cockroaches (<i>Blattella germanica</i>): 0.12-0.16 g/m ² (3-4 drops/m ²).

Oriental cockroaches (*Blatta orientalis*): 0.16-0.24 g/m² (4-6 drops/m²)
American cockroaches (*Periplaneta americana*): 0.16 g/m² -0.24 g/m² (4-6 drops/m²)
Frequency of application:
1 application in 2 weeks. Reapply only once more if the infestation persists.
Frequency of treatment: Three months after the infestation's end, treatment may be repeated. Maximum of 6 applications per year.

Category(ies) of users

Professional

Pack sizes and packaging material

LDPE plastic syringes of 1, 2, 3, 4, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55 and 60 grams of gel bait.

4.2.1 Use-specific instructions for use

Ready-to-use gel bait to be applied as drops deposited with the syringe. Drops adhere to non-porous surfaces.
Apply only in cracks and crevices.
Use indoors in private houses.
Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel.
Apply the biocidal product only as spot application in cracks and crevices where cockroaches may be present out of the reach of children and animals:
- In inaccessible places.
- In dark places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: cracks and crevices behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.
Do not use on wood or porous surfaces.
Avoid contact with treated surfaces.
Do not expose bait drops to sunlight or heat source (i.e. radiator).
Do not mix with other chemicals or in areas recently treated with another insecticide.

4.2.2 Use-specific risk mitigation measures

Avoid contact with treated spots.
Avoid contact with eyes and skin.
The product should not be applied in a zone accessible to children
The treatment must be restricted to areas out reach of animals.

4.2.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.2.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product and other waste generated during the treatment are considered hazardous waste. Dispose of in accordance with current regulations.
Do not release to soil, ground, surface water or any kind of sewer.

4.2.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

4.3 Use description

Use 3 - Use # 3 – Indoors, cracks and crevices - Gel bait applied as drops - Trained professional users.

Product type

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Where relevant, an exact description of the authorised use

Insecticide and growth regulator product against cockroaches.

Target organism(s) (including development stage)

Scientific name: *Periplaneta americana*
Common name: Cucaracha americana
Development stage: adults, nymphs

Scientific name: *Blatta orientalis*
Common name: Cucaracha oriental
Development stage: adults, nymphs

Scientific name: *Blattella germanica*
Common name: Cucaracha alemana
Development stage: adults, nymphs

Field(s) of use

Indoor

Indoors, crack and crevices.
Private houses and commercial buildings.

Application method(s)

Method: Bait application
Detailed description:
Open application of a gel bait applied as drops from a syringe

Application rate(s) and frequencies

Application Rate: 1 application in 2 weeks. Reapply only once more if the infestation persists.
Dilution (%): 0
Number and timing of application:
Dose: depends on the level of infestations and species of cockroaches (1 drop = 0.04 g).
German cockroaches (*Blattella germanica*): 0.12-0.16 g/m² (3-4 drops/m²).
Oriental cockroaches (*Blatta orientalis*): 0.16-0.24 g/m² (4-6 drops/m²)
American cockroaches (*Periplaneta americana*): 0.16 g/m² -0.24 g/m² (4-6 drops/m²)
Frequency of application:
1 application in 2 weeks. Reapply only once more if the infestation persists.
Frequency of treatment: Three months after the infestation's end, treatment may be repeated. Maximum of 6 applications per year.

Category(ies) of users

Trained professional

Pack sizes and packaging material

LDPE syringe of 1, 2, 3, 4, 5, 8, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55 and 60 g of product.
LDPE plastic cartridge of 1, 2, 3, 4, 5, 8, 10, 15, 20, 25, 30, 35, 40, 50, 60, 75, 80, 100, 125, 175, 200, 250, 300, 350, 400, 450 and 500g of product.

4.3.1 Use-specific instructions for use

Ready-to-use gel bait to be applied as drops deposited with the cartridge. Drops adhere to non-porous surfaces.
Apply only in cracks and crevices.
Use indoors in houses and commercial buildings.
Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel.
Apply the biocidal product only as spot application in cracks and crevices where cockroaches may be present out of the reach of children and animals:

- In inaccessible places:.
- In dark places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: cracks and crevices behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.

Do not mix with other chemicals or in areas recently treated with another insecticide
Do not use on wood or porous surfaces.
Avoid contact with treated surfaces.
Do not expose bait drops to sunlight or heat source (i.e. radiator).
In storerooms the product may only be applied in the presence of properly packaged food.

4.3.2 Use-specific risk mitigation measures

Avoid contact with treated spots.
Wear appropriate gloves during application.
Avoid contact with eyes and skin.
The product should not be applied in a zone accessible to children.
The treatment must be restricted to areas out of reach of animals.

Adopt integrated pest management methods such as the combination of chemical, physical control methods and other public health measures, taking into account local specificities (climatic conditions, target species, conditions of use, etc.)
Check the efficacy of the product on site: if need be, cause of reduced efficacy must be investigated to ensure that there is no resistance or to identify potential resistance.
Do not use the product in areas where resistance is suspected or established.
Inform the authorisation holder if the treatment is ineffective.

4.3.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.3.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product, washing water, containers and other waste generated during the treatment are considered hazardous waste. Deliver those wastes to a registered establishment or undertaking, in accordance with current regulations.
Code the waste according to Decision 2014/955 / EU.
Do not release to soil, ground, surface water or any kind of sewer.

--

4.3.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

4.4 Use description

Use 4 - Use # 4 – Indoors-Gel bait applied in bait stations – General public (non-professional users)

Product type	PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)
Where relevant, an exact description of the authorised use	Insecticide and growth regulator against cockroaches.
Target organism(s) (including development stage)	Scientific name: <i>Periplaneta americana</i> Common name: Cucaracha americana Development stage: adults, nymphs Scientific name: <i>Blatta orientalis</i> Common name: Cucaracha oriental Development stage: adults, nymphs Scientific name: <i>Blattella germanica</i> Common name: Cucaracha alemana Development stage: adults, nymphs
Field(s) of use	Indoor Indoors. Private houses.
Application method(s)	Method: Bait application Detailed description: Ready-to-use bait stations.
Application rate(s) and frequencies	Application Rate: Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists. Dilution (%): 0 Number and timing of application: Dose: depending on the level of infestations: - For low infestations, 0.16 g/m ² - For high infestations, 0.24 g/m ² Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists. Frequency of treatment: Three months after the infestation's end, treatment may be repeated.

Category(ies) of users

General public (non-professional)

Pack sizes and packaging material

Plastic (PET) bait station/bait station capsule with 1, 2, 2.5, 3, 4 and 5 grams of gel bait.

4.4.1 Use-specific instructions for use

Ready-to-use gel bait in bait stations.

Use indoors in private houses.

Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel

Apply the biocidal product by placement of bait stations only in places where cockroaches may be present out of the reach of children and animals:

- In inaccessible places: behind furniture, etc.
- In dark and wet places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.

4.4.2 Use-specific risk mitigation measures

Do not force open the bait station.

Never introduce the fingers through the holes in the bait box.

4.4.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.4.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product and other waste generated during the treatment are considered hazardous waste. Dispose of in accordance with current regulations.

Do not release to soil, ground, surface water or any kind of sewer.

At the end of the treatment, collect bait boxes for disposal.

4.4.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

4.5 Use description

Use 5 - Use # 5 – Indoors - Gel bait applied in bait stations – Professional users.

Product type	PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)
Where relevant, an exact description of the authorised use	Insecticide and growth regulator product against cockroaches.
Target organism(s) (including development stage)	Scientific name: <i>Periplaneta americana</i> Common name: Cucaracha americana Development stage: adults, nymphs) Scientific name: <i>Blatta orientalis</i> Common name: Cucaracha oriental Development stage: adults, nymphs Scientific name: <i>Blattella germanica</i> Common name: Cucaracha alemana Development stage: adults, nymphs
Field(s) of use	Indoor Indoors. Houses and commercial buildings.
Application method(s)	Method: Bait application Detailed description: Ready-to-use bait stations
Application rate(s) and frequencies	Application Rate: Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists. Dilution (%): 0 Number and timing of application: Dose: depending on the level of infestations: - For low infestations, 0.16 g/m ² - For high infestations, 0.24 g/m ² Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists. Frequency of treatment: Three months after the infestation's end, treatment may be repeated.
Category(ies) of users	Professional
Pack sizes and packaging material	Plastic (PET) bait station/bait station capsule with 1, 2, 2.5, 3, 4 and 5 grams of gel bait.

4.5.1 Use-specific instructions for use

Ready-to-use gel bait in bait stations.
 Use indoors in private houses.
 Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel.
 Apply the biocidal product by placement of bait stations only in places where cockroaches may be present out of the reach of children and animals:

- In inaccessible places: behind furniture, etc.
- In dark and wet places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.

4.5.2 Use-specific risk mitigation measures

Do not force open the bait station.
 Never introduce the fingers through the holes in the bait box.

4.5.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.5.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product and other waste generated during the treatment are considered hazardous waste. Dispose of in accordance with current regulations.
 Do not release to soil, ground, surface water or any kind of sewer.
 At the end of the treatment, collect bait boxes for disposal.

4.5.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

4.6 Use description

Use 6 - Use # 6 – Indoors - Gel bait applied bait stations – Trained professional users.

Product type

PT18 - Insecticides, acaricides and products to control other arthropods (Pest control)

Where relevant, an exact description of the authorised use

Insecticide and growth regulator product against cockroaches.

Target organism(s) (including development stage)

Scientific name: *Periplaneta americana*
 Common name: Cucaracha americana
 Development stage: adults, nymphs

Scientific name: *Blatta orientalis*
 Common name: Cucaracha oriental

	<p>Development stage: adults, nymphs</p> <p>Scientific name: <i>Blattella germanica</i> Common name: Cucaracha alemana Development stage: adults, nymphs</p>
Field(s) of use	<p>Indoor</p> <p>Indoors. Houses and commercial buildings</p>
Application method(s)	<p>Method: Bait application Detailed description: Ready-to-use bait stations.</p>
Application rate(s) and frequencies	<p>Application Rate: Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists. Dilution (%): 0 Number and timing of application: Dose: depending on the level of infestations: - For low infestations, 0.16 g/m² - For high infestations, 0.24 g/m²</p> <p>Frequency of application: After about 4 weeks, bait stations should be replaced with fresh ones if the infestation persists.</p> <p>Frequency of treatment: Three months after the infestation's end, treatment may be repeated.</p>
Category(ies) of users	<p>Trained professional</p>
Pack sizes and packaging material	<p>Plastic (PET) bait station/bait station capsule with 1, 2, 2.5, 3, 4, 5, 8, 10, 15, 20, 25 and 30grams of gel bait.</p>

4.6.1 Use-specific instructions for use

Ready-to-use gel bait in bait stations.
Use indoors in houses and commercial buildings.
Before treatment, remove all natural source of food from the infested area to encourage the ingestion of the gel.
Apply the biocidal product by placement of bait stations only in places where cockroaches may be present out of the reach of children and animals:

- In inaccessible places: behind furniture, etc.
- In dark and wet places: under the sink, behind the toilet, near the drain, etc.
- In warm temperature places: behind the engines, refrigerators, washing machines, dishwashers, etc.
- In places with food waste or organic materials: under or behind kitchen cupboards, near the dustbin, in storerooms, in cellars, in courtyards, etc.

4.6.2 Use-specific risk mitigation measures

Do not force open the bait station.
Never introduce the fingers through the holes in the station.
Adopt integrated pest management methods such as the combination of chemical, physical control methods and other public health

measures, taking into account local specificities (climatic conditions, target species, conditions of use, etc.)
Check the efficacy of the product on site: if need be, cause of reduced efficacy must be investigated to ensure that there is no resistance or identify potential resistance.
Do not use the product in areas where resistance is suspected or established.
Inform the authorisation holder if the treatment is ineffective.

4.6.3 Where specific to the use, the particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

See Directions of use

4.6.4 Where specific to the use, the instructions for safe disposal of the product and its packaging

Empty containers, unused product, washing water, containers and other waste generated during the treatment are considered hazardous waste. Deliver those wastes to a registered establishment or undertaking, in accordance with current regulations.
At the end of the treatment, collect bait boxes for disposal.
Code the waste according to Decision 2014/955 / EU.
Do not release to soil, ground, surface water or any kind of sewer.

4.6.5 Where specific to the use, the conditions of storage and shelf-life of the product under normal conditions of storage

See Directions of use

5. General directions for use

5.1. Instructions for use

Always read the label or leaflet before use and respect all the instructions provided.
Make an inspection before applying the product to check the level of infestation and affected areas.
Use only indoors out of the reach of children and animals.

5.2. Risk mitigation measures

This product should be used in alternation with other products not containing the same a.s. to avoid resistant populations.
Do not use directly on or near food, feed or drinks, or on surfaces or utensils likely to be in direct contact with food, feed, drinks and livestock. The product should be reapplied when finished only until the pest is controlled.
Use products at recommended doses and intervals.
To optimise the treatment efficacy, respect good hygiene practices: remove or prevent access to all source of food. The bait must be the main source of food available for the cockroaches.
To optimise the efficacy, check the bait once a week and replace/replenish bait if they are damaged or soiled.
Product must be securely applied in a way so as to minimize the risk of consumption by other animals or children.
Use only in concealed areas difficult to access and kept away from water.
Dispose of unused product, its packaging and all other waste (i.d. dead insects) in accordance with local regulations.

5.3. Particulars of likely direct or indirect effects, first aid instructions and emergency measures to protect the environment

Basic First aid procedures:
IF ON SKIN:
Wash it before reuse. Wash skin with water. If skin irritation or rash occur: Get medical advice.
IF IN EYES:
If symptoms occur rinse with water. Remove contact lenses, if present and easy to do. Call a POISON CENTRE or a doctor.
• If necessary take person to a hospital and show the label or packaging when possible. Do not leave poisoned person alone.

Medical advice for doctors and sanitary staff
• Symptomatic and supportive treatment.

IF MEDICAL ADVICE IS NEEDED, KEEP THE LABEL OR CONTAINER HANDY AND CONSULT THE MEDICAL SERVICE FOR TOXICOLOGICAL INFORMATION

5.4. Instructions for safe disposal of the product and its packaging

See the specific sections 4.1.4, 4.2.4, 4.3.4., 4.5.4.and 4.6.4.

5.5. Conditions of storage and shelf-life of the product under normal conditions of storage

The storage stability of this product in its original container is 4 years under normal condition of storage.

Protect from frost.

Store in the original container tightly closed.

Store in a dry, cool and well ventilated place.

It is recommended to store the product at a temperature preferably between 5°C and 45°C.

Store away from light.

Do not store near food, drink and animal feedingstuff.

6. Other information

Definitions:

Trained professional: pest control operators, having received specific training in insecticide control according to the national legislation in force.

Professional: User applying biocidal products in the workplace. This user has some knowledge and skills in the handling of chemicals, and is able to correctly use personal protective equipment (PPE) if necessary.

General public (non-professional user): Users who are not professionals and who apply the product in the context of their private life.

The product contains a bitter substance that makes it repulsive to people or pets.

The applicant must ensure that the general public can understand the difference between species and the level of infestation for correct use of the dose.

The drops should have 0.04 g of gel bait. The applicant should indicate in the label the size of a drop with 0.04 g of product.

The number of bait stations should be indicated in the label depending on the amount of product included in the bait station. E.g. 0.16 g/m² = 4 bait station with 1 g in a room of 25 m².